

# RANDOM OUTER AUTOMORPHISMS OF FREE GROUPS: ATTRACTING TREES AND THEIR SINGULARITY STRUCTURES

ILYA KAPOVICH, JOSEPH MAHER, CATHERINE PFAFF, AND SAMUEL J. TAYLOR

*Catherine Pfaff would like to dedicate this paper to her father Dr. Roland Pfaff, zikhrono livrakha.*

ABSTRACT. We prove that a “random” free group outer automorphism is an ageometric fully irreducible outer automorphism whose ideal Whitehead graph is a union of triangles. This implies that its attracting tree is a nongeometric  $\mathbb{R}$ -tree all of whose branch points are trivalent. Among the applications, we answer in the affirmative a question of Handel and Mosher who asked if it is true that for a “generic”  $\varphi \in \text{Out}(F_r)$  we have that  $\varphi$  is fully irreducible with the attracting trees for both  $\varphi$  and  $\varphi^{-1}$  being nongeometric, i.e. not the dual  $\mathbb{R}$ -tree to a foliation on a finite simplicial 2-complex.

## 1. INTRODUCTION

Let  $\text{Out}(F_r)$  denote the outer automorphism group of a fixed free group of rank  $r \geq 3$ . The study of this group has fundamental contributions dating back to Nielsen, starting in 1915, and Whitehead in the 1930’s to 1950’s. Recently, substantial progress has been made utilizing the fact that just as the mapping class group encodes all (equivalence classes of) homeomorphisms of a surface,  $\text{Out}(F_r)$  encodes all (equivalence classes of) homotopy equivalences of graphs of a fixed rank. In particular, this gives a powerful analogue of Teichmüller space, namely Outer space (denoted  $\text{CV}$ ), the deformation space of marked metric graphs. Furthermore, this correspondence gives rise to related analogues of the curve complex, the train track theory, and the study of asymptotic conjugacy class invariants. We focus on determining generic properties of such invariants.

There are several ways of formalizing the notion of a “random” or “generic” element of a discrete group  $G$ . One natural approach involves considering an element  $w_n \in G$  given by a long (with large  $n$ ) random walk on  $G$  with respect to some discrete probability measure  $\mu$  on  $G$ . Then one says that a particular property holds for a “random” element of  $G$  if the probability that  $w_n \in G$  has this property tends to 1 as  $n \rightarrow \infty$ . In this context “random” group elements have been used to analyze various boundaries associated to  $G$ , to produce existence results, to study analytic properties of groups, to study the complexity of various group-theoretic algorithms on “random” inputs, etc.

Recent work of Maher and Tiozzo [MT14] shows that, for a group  $G$  acting isometrically on a Gromov-hyperbolic space  $X$ , under some mild assumptions on the measure  $\mu$ , a “random” element  $w_n \in G$  acts loxodromically on  $X$ . This result has numerous useful applications. For example, by considering the action of the mapping class group  $\text{MCG}(\Sigma)$  on the curve complex  $\mathcal{C}(S)$  one concludes that a “random” element of  $G$  is pseudo-Anosov. Originally this was established by Rivin using homological methods, [Riv08], with stronger restrictions on  $\mu$ . See also [Hor14, KM98, CM15, Mah11, Sis11, MS17] for related results. Similarly, by considering the action of  $\text{Out}(F_r)$  on

---

2010 *Mathematics Subject Classification.* Primary 20F65, Secondary 57M, 37B, 37D.

*Key words and phrases.* Free group, random walk, Outer space, free group automorphisms, train track maps.

The first named author was supported by the individual NSF grants DMS-1405146 and DMS-1710868. The second named author was supported by Simons Foundation and PSC-CUNY. The fourth named author was partially supported by NSF grant DMS-1744551. All authors acknowledge support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 “RNMS: GEometric structures And Representation varieties” (the GEAR Network).

the free factor complex  $\mathcal{FF}$ , and applying the result of Maher and Tiozzo mentioned above, one concludes that a “random” element of  $\text{Out}(F_r)$  is fully irreducible. One can then ask deeper and finer questions about the properties of such “random” pseudo-Anosovs and fully irreducibles, such as, for example, their “index” or “singularity” properties.

Recently, Gadre and Maher [GM17] provided such finer information in the mapping class group setting. They showed for a random walk  $(w_n)$  on  $\text{MCG}(\Sigma)$  defined by a measure  $\mu$  (under appropriate natural assumptions on  $\mu$  and  $\Sigma$ ) that, with probability going to 1 as  $n \rightarrow \infty$ , the mapping class  $w_n$  is a principle pseudo-Anosov in the sense that its invariant foliations have only 3-pronged singularities (or equivalently that its axis lies in the principal stratum). Their result uses the local structure of the cotangent space of Teichmüller space in an essential way. No analogous local properties hold in Outer space, and instead, we make use of the (non-local) properties of elements of  $\text{Out}(F_r)$  with lone axes, as we discuss below in more detail.

In this paper we study similar questions for “random” elements of  $\text{Out}(F_r)$ . An element  $\varphi \in \text{Out}(F_r)$  is called *fully irreducible* if no positive power of  $\varphi$  leaves invariant a nontrivial proper free factor of  $F_r$ . Fully irreducibles are the dynamically minimal elements of  $\text{Out}(F_r)$  and serve as the main counterparts to pseudo-Anosovs in the  $\text{Out}(F_r)$  setting. We study two highly related asymptotic conjugacy class invariants for a fully irreducible  $\varphi \in \text{Out}(F_r)$ , namely its attracting tree  $T_+^\varphi$  and its index  $i(\varphi)$ .

We define  $T_+^\varphi$  in §3.3, only remarking here that the attracting tree for a pseudo-Anosov is the  $\mathbb{R}$ -tree dual to its invariant foliation and hence has a  $\pi_1$ -orbit of valence- $k$  branch points for each  $k$ -pronged foliation singularity and vice versa. As in [LP97], we call an  $\mathbb{R}$ -tree geometric that is dual to a (singular) foliation on a finite 2-complex. If a fully irreducible  $\varphi$  is *geometric*, i.e. is induced by pseudo-Anosov homeomorphisms of compact surfaces with one boundary component and with fundamental group isomorphic to  $F_r$ , then  $T_+^\varphi$  is geometric. It is possible that  $\varphi$  is nongeometric, but that  $T_+^\varphi$  is still geometric, in which case  $\varphi$  is called *parageometric* (a circumstance studied, for example, in [HM07]). In the remaining case, a fully irreducible is called *ageometric*, a circumstances long conjectured to be generic (with the question formally stated by Handel and Mosher in [HM07]), but with a highly elusive proof. We prove this in Theorem A.

There exists an equivalent “ageometric” definition in terms of the index  $i(\varphi)$ . The index theory of  $\text{Out}(F_r)$  dates back to the 1990s [GL95, GJLL98] and turns out to be considerably more complicated than in the mapping class group setting. One of the key differences between the  $\text{MCG}(\Sigma)$  and the  $\text{Out}(F_r)$  case is that the value of  $i(\varphi)$  is not fixed as a function of the rank  $r$  and in general one only has the inequality  $0 > i(\varphi) \geq 1 - r$ . For geometric and parageometric fully irreducibles  $\varphi \in \text{Out}(F_r)$  one has  $i(\varphi) = 1 - r$ . Ageometric fully irreducibles are those for which  $i(\varphi) > 1 - r$ .

A testament to the depth of  $\text{Out}(F_r)$  invariant theory is the level of refinement of the index. The most detailed such refinement is the ideal Whitehead graph  $IW(\varphi)$  defined by Handel and Mosher in [HM11] and proved to relate intimately to CV geodesics in [HM11], [MP16], [AKKP18], [AKP17], for example. The lamination leaves for a fully irreducible  $\varphi$  are obtained by iteratively applying (a representative of)  $\varphi$  to a suitable conjugacy class in  $F_r$ , and taking an appropriate limit, see [BFH97] for further details. The ideal Whitehead graph records how the lamination leaves pass through the branch points of  $T_+^\varphi$ , and one can view branch points as extending the notion of a foliation singularity to a nongeometric  $\mathbb{R}$ -tree. In particular, if the lamination leaf passes through a valence- $k$  branch point  $p$  in every possible manner, then there is a corresponding component of the the ideal Whitehead graph that is the complete graph on  $k$  vertices.

Our main result, explained in more detail below, shows that for a “random” fully irreducible  $\varphi \in \text{Out}(F_r)$  the attracting tree  $T_+^\varphi$  is trivalent, that is, all branch points of  $T_+^\varphi$  have valency three.

**1.1. Main results.** Before stating our main results, we introduce a few more definitions (even further notation and explanations follow the statement). For  $r \geq 3$ , let  $\Delta_r$  denote the graph which

is the disjoint union of  $2r - 3$  triangles. We call an ageometric fully irreducible  $\varphi \in \text{Out}(F_r)$  *principal* if  $IW(\varphi) \cong \Delta_r$  and *triangular* if  $IW(\varphi)$  is isomorphic to the union of several components of  $\Delta_r$ , that is, if  $IW(\varphi)$  is the disjoint union of  $\leq 2r - 3$  triangles. Note that for a principal  $\varphi$  we have  $i(\varphi) = \frac{3}{2} - r$  and for a triangular  $\varphi$  we have  $i(\varphi) \geq \frac{3}{2} - r$ . We can now state our main result (formally answering the Handel-Mosher question, but with substantially stronger information):

**Theorem A.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ . Let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose  $\langle \text{supp}(\mu) \rangle_+$  contains  $\varphi^{-1}$  for some principal fully irreducible  $\varphi \in \text{Out}(F_r)$ . Then  $w_n$  is an ageometric triangular fully irreducible outer automorphism with probability going to 1 as  $n \rightarrow \infty$ .*

Here  $\mathcal{FF}$  is the *free factor graph* for  $F_r$ , the free group analog of the curve graph (see §6). The key points relevant here are that  $\mathcal{FF}$  is Gromov-hyperbolic and that  $\psi \in \text{Out}(F_r)$  acts loxodromically on  $\mathcal{FF}$  if and only if  $\psi$  is fully irreducible, as shown by Bestvina and Feighn [BF14]. Above,  $\langle \text{supp}(\mu) \rangle_+$  denotes the subsemigroup of  $G$  generated by the support of  $\mu$ . The action on  $\mathcal{FF}$  is considered *nonelementary* if  $\langle \text{supp}(\mu) \rangle_+$  contains two independent loxodromic isometries of  $\mathcal{FF}$ .

Theorem A has several important consequences:

**Corollary B.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ , and let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose that  $\langle \text{supp}(\mu) \rangle_+$  contains  $\varphi^{-1}$  for some principal fully irreducible  $\varphi \in \text{Out}(F_r)$ .*

*Then, with probability going to 1 as  $n \rightarrow \infty$ ,  $w_n$  is an ageometric fully irreducible outer automorphism  $\varphi_n$  satisfying:*

- (1) *each component of the ideal Whitehead graph  $IW(\varphi)$  is a triangle;*
- (2) *each branchpoint of  $T_+^{\varphi_n}$  is trivalent;*
- (3)  *$i(\varphi_n) > 1 - r$  and  $\text{ind}_{\text{geom}}(T_+^{\varphi_n}) < 2r - 2$  (here  $i(\varphi_n)$  is the rotationless index of  $\varphi_n$  and  $\text{ind}_{\text{geom}}(T_+^{\varphi_n})$  is the geometric index of  $\varphi_n$ );*
- (4) *the tree  $T_+^{\varphi_n}$  is nongeometric.*

In fact, our arguments apply equally well to the inverse of a random outer automorphism:

**Corollary C.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ . Let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose that  $\langle \text{supp}(\mu) \rangle_+$  contains a pair of elements  $\varphi$  and  $\psi$ , such that both  $\varphi$  and  $\psi^{-1}$  are principal fully irreducible outer automorphisms.*

*Then, with probability approaching 1 as  $n \rightarrow \infty$ , both  $w_n$  and  $w_n^{-1}$  are ageometric triangular fully irreducible outer automorphisms. In particular, both fixed trees of  $w_n$  are trivalent.*

We remark that it is proved in [AKKP18, Example 6.1] that for each  $r \geq 3$  there exists a principal  $\varphi \in \text{Out}(F_r)$ . Thus the assumption in Theorem A and in Corollary B that  $\langle \text{supp}(\mu) \rangle_+$  contain the inverse of some principal outer automorphism is automatically satisfied if, for example,  $\langle \text{supp}(\mu) \rangle_+ = \text{Out}(F_r)$  or if  $\langle \text{supp}(\mu) \rangle_+$  is a subgroup of finite index in  $\text{Out}(F_r)$ . Similarly, the assumption in Corollary C, that  $\langle \text{supp}(\mu) \rangle_+$  contains a pair of elements  $\varphi$  and  $\psi$  such that both  $\varphi$  and  $\psi^{-1}$  are principal fully irreducible outer automorphisms, is also automatically satisfied in the case where either  $\langle \text{supp}(\mu) \rangle_+ = \text{Out}(F_r)$  or  $\langle \text{supp}(\mu) \rangle_+$  is a subgroup of finite index in  $\text{Out}(F_r)$ .

Corollary C is where we directly answer in the affirmative the question of Handel and Mosher [HM07, p. 3158] as to whether it is true that for a “generic”  $\varphi \in \text{Out}(F_r)$  (where  $r \geq 3$ ) we have that  $\varphi$  is fully irreducible with both trees  $T_+^\varphi, T_+^{\varphi^{-1}}$  nongeometric (i.e. that both  $\varphi$  and  $\varphi^{-1}$  are ageometric fully irreducibles).

**1.2. Elements of proof.** While much powerful random walk machinery has been developed over the past decade for the mapping class group and related groups, there are significant structural

obstacles to the development and application of these techniques in our setting. For example, the Gadre-Maher work heavily utilizes the local Teichmüller space structure, and there is no direct analogue of this in Outer Space CV. In particular, they use the fact that the principal stratum is open in the space of quadratic differentials, which is equal to the cotangent space to Teichmüller space. Therefore a sequence of quadratic differentials with limit in the principal stratum is eventually contained in the principal stratum. We overcome this and other obstacles by combining the stable strata work of [AKKP18] with the lone axis work of [MP16], and general axis bundle constructions.

One result we obtain, using the work of [BF14, DT17, BR15, DH17], is an adapted version of the bounded geodesic image (BGI) property, as described in §7.2. For symmetric metric spaces, the BGI property is equivalent to the strongly contracting property, but the natural metric on outer space CV is asymmetric. We make essential use of properties of the projection  $\pi$  from CV to the free factor complex  $\mathcal{FF}$ , as developed in [BR15, DT17], dealing carefully with the fact that  $\pi$  is far less controlled than in the analogous mapping class group setting.

Our proof can be summarized as follows. Fully irreducible  $\varphi \in \text{Out}(F_r)$  have invariant axes in CV, that project under  $\pi$  to axes in  $\mathcal{FF}$ . By [AKKP18] we know that maximally triangular  $\varphi$  exist in each  $\text{Out}(F_r)$  and have the stable strata property: A maximally triangular  $\varphi$  has the stable strata property if for any other fully irreducible element  $\varphi'$ , if the axis of  $\varphi'$  fellow travels with the axis of  $\varphi$ , closely enough, and for long enough, then the ideal Whitehead graph of  $\varphi'$  is triangular. By Maher-Tiozzo [MT14], we know that a generic sample path contains arbitrarily long subsegments which stay close to maximally triangular axes in  $\mathcal{FF}$ . Since  $\pi$  is coarsely Lipschitz, we may use the BGI property to say that our lifted sample path and maximally triangular axes stay a bounded distance apart for an arbitrary length. The lone axis property promotes the “bounded distance” to a necessary  $\varepsilon$ , allowing us to finally use the stable strata property of [AKKP18].

**1.3. Further questions.** Theorem A naturally raises the questions of whether the properties of being principal or lone axis are typical, in the random walk sense, for the elements of  $\text{Out}(F_r)$ . We believe that the answer to both of these questions is probably negative, but that principal outer automorphisms do occur with asymptotically positive probability. Note that, as a consequence of the results of [MP16], for a triangular automorphism  $\varphi$  it follows that  $\varphi$  is lone axis if and only if  $\varphi$  is principal, i.e. if and only if  $i(\varphi) = \frac{3}{2} - r$ . Computer experiments, that we conducted using Thierry Coulbois’ SAGE train track package<sup>1</sup>, appear to indicate that several possibilities for  $i(\varphi)$  (and not just  $i(\varphi) = \frac{3}{2} - r$ ) happen with asymptotically positive probability. Moreover, specific examples exhibited in [AKKP18] demonstrate a specific reason why a loss of several components of  $IW(\varphi)$  may occur, resulting in the loss of the lone axis property. Roughly, a periodic vertex in a train track representative of  $\varphi$ , which contributes to  $IW(\varphi)$ , may become nonperiodic for a “nearby” outer automorphism  $\psi$  and thus no longer contribute to  $IW(\psi)$ . Heuristic considerations suggest that this behavior will have detectable probabilistic effect.

However, this assumption in Theorem A cannot, in general, be removed. In [KP15] Kapovich and Pfaff constructed a “train track directed” random walk on  $\text{Out}(F_r)$ . While that walk is defined by an irreducible finite state Markov chain, rather than by a sequence of i.i.d. random variables distributed according to  $\mu$ , qualitatively the walk behaves similarly to the ones considered in the present paper. However, with asymptotically positive probability, for the outer automorphisms  $\varphi_n$  produced by that walk after  $n$  steps, the ideal Whitehead graph  $IW(\varphi_n)$  is the complete graph on  $2r - 1$  vertices. Thus  $\varphi_n$  is neither principal nor triangular in that case. The walk considered in [KP15] is constructed in such a way that it never encounters principal outer automorphisms.

---

<sup>1</sup>The Coulbois computer package is available at [Cou14].

**Acknowledgements.** We are grateful to Lee Mosher and Chris Leininger for useful conversations. We also thank Yael Algom-Kfir, who contributed to early discussions of the problems considered in this paper.

## 2. METRIC SPACES, RANDOM WALKS, AND ORIENTED MATCHES

**2.1. Metric space notation.** In this section we collect together some basic definitions about (possibly asymmetric) metric spaces. The main two examples we shall consider are Outer space with the Lipschitz metric  $(CV, d_{CV})$ , an asymmetric geodesic metric space which is locally compact, and the complex of free factors  $(\mathcal{FF}, d_{\mathcal{FF}})$ , a symmetric Gromov hyperbolic space which is not locally compact. See §3.1 and §6 for further details.

An (asymmetric) metric space is a pair  $(X, d_X)$ , where  $X$  is a topological space, and  $d_X$  is a function  $d_X: X \times X \rightarrow \mathbb{R}_{\geq 0}$  which satisfies  $d_X(x, x) = 0$  for all points  $x \in X$ , and  $d_X(x, y) \leq d_X(x, z) + d_X(z, y)$ , for all points  $x, y, z \in X$ . We say that  $(X, d_X)$  is *symmetric* if  $d_X(x, y) = d_X(y, x)$  for all  $x, y \in X$ .

We say that the (possibly asymmetric) metric space  $(X, d_X)$  is *geodesic* if for each pair of points  $x, y \in X$  there is a subinterval  $I = [0, d_X(x, y)] \subset \mathbb{R}$  and a path  $\gamma: I \rightarrow X$  such that  $d_X(x, \gamma(t)) = t$ . Let  $I$  be a connected subset of  $\mathbb{R}$ , or a set of consecutive integers in  $\mathbb{Z}$ , and let  $Q$  be a non-negative constant. We say that a map  $\gamma: I \rightarrow (X, d_X)$  is a *Q-quasigeodesic* if for all  $s < t$  in  $I$ ,

$$\frac{1}{Q}|s - t| - Q \leq d_X(\gamma(s), \gamma(t)) \leq Q|s - t| + Q.$$

We say that  $X$  is *Gromov hyperbolic*, or  $\delta$ -*hyperbolic*, if  $(X, d_X)$  is a symmetric geodesic metric space, and there is a constant  $\delta \geq 0$  such that for any geodesic triangle in  $X$ , each side is contained in a  $\delta$ -neighbourhood of the other two.

The *Gromov product* of three points  $x, y, z \in X$  is defined to be

$$(y|z)_x = \frac{1}{2}(d_X(x, y) + d_X(x, z) - d_X(y, z)).$$

If  $X$  is  $\delta$ -hyperbolic, then  $(y|z)_x$  is equal to the distance from  $x$  to a geodesic from  $y$  to  $z$ , up to an error of at most  $2\delta$ . Given two points  $x$  and  $y$  in  $X$ , and a number  $R$ , we define the *shadow*  $S_x(y, R)$  to be

$$S_x(y, R) = \{z \in X \mid (y|z)_x \geq d_X(x, y) - R\}.$$

The *Gromov boundary*  $\partial X$  may be defined to be equivalence classes of quasigeodesic rays  $\gamma: \mathbb{R}_{\geq 0} \rightarrow X$ , where two quasigeodesic rays  $\gamma$  and  $\gamma'$  are equivalent if there exists a value  $K$  such that  $d(\gamma(t), \gamma'(t)) < K$  for all  $t \in \mathbb{R}_{\geq 0}$ . Given  $\delta \geq 0$ , there is a constant  $Q \geq 0$  such that any two distinct points in  $\partial X$  are connected by a bi-infinite  $Q$ -quasigeodesic.

Let  $g$  be an isometry of  $X$ . We say that  $g$  is *loxodromic* if the orbit of  $(g^n x)_{n \in \mathbb{Z}}$  of any point  $x \in X$  is a quasigeodesic in  $X$ . A loxodromic element  $g$  has precisely two fixed points in the boundary, and we call any bi-infinite  $Q$ -quasigeodesic connecting them a *quasi-axis* for  $g$ , though we may always assume the quasi-axis is  $g$ -invariant.

**2.2. Random walks background.** Let  $G$  be a countable group, and let  $\mu$  be a probability distribution on  $G$ . We shall define the *step space* to be the countable product  $(G, \mu)^{\mathbb{Z}}$ , and we shall write  $g_n$  for projection onto the  $n$ -th factor. The random variables  $(g_n)_{n \in \mathbb{Z}}$  then correspond to a bi-infinite sequence of independent  $\mu$ -distributed random variables. A sequence of steps  $\omega \in (G, \mu)^{\mathbb{Z}}$  determines a random walk, whose location at time  $n$  is given by the product of the first  $n$  steps. More precisely, the location of the random walk at time 0 is  $w_0 = 1$ , and the location of the random walk at time  $n$  is given by  $w_{n+1} = w_n g_n$ . We shall write  $\check{\mu}$  for the reflected measure, i.e.  $\check{\mu}(g) = \mu(g^{-1})$ , and we shall write  $\mu^{*n}$  for the  $n$ -fold convolution of  $\mu$  with itself. In particular,  $w_n$

is a random variable on  $(G, \mu)^{\mathbb{Z}}$  with values in  $G$ , whose distribution is given by  $\mu^{*n}$  if  $n > 0$  and by  $\check{\mu}^{*n}$  if  $n < 0$ .

For a group  $G$  with a probability distribution  $\mu$ , the *support*  $\text{supp}(\mu)$  of  $\mu$  is the set of all elements  $g \in G$  such that  $\mu(g) > 0$ . We also denote by  $\langle \text{supp}(\mu) \rangle_+$  the sub-semigroup of  $G$  generated by  $\text{supp}(\mu)$ . Note that for  $g \in G$  we have  $g \in \langle \text{supp}(\mu) \rangle_+$  if and only if there exists an  $n \geq 1$  such that  $\mu^{*n}(g) > 0$ .

Let  $G$  act on a hyperbolic metric space  $X$  by isometries. We say two loxodromic isometries are *independent* if their fixed point sets are disjoint in the Gromov boundary  $\partial X$ . We say that the probability distribution  $\mu$  is *nonelementary* if  $\langle \text{supp}(\mu) \rangle_+$  contains two independent loxodromic elements.

A choice of basepoint  $x_0 \in X$  determines an orbit map from  $G$  to  $X$  given by  $g \mapsto gx_0$ . A random walk  $(w_n)_{n \in \mathbb{Z}}$  on  $G$  then gives rise to a sequence  $(w_n x_0)_{n \in \mathbb{Z}}$  in  $X$ . If  $\mu$  is nonelementary then almost every sequence  $(w_n)_{n \in \mathbb{Z}}$  converges to a point in  $\partial X$  in both the forward and backward directions. Let  $\gamma_\omega$  be a bi-infinite  $Q$ -quasigeodesic in  $X$  connecting the two limit points. We say that  $\mu$  has *bounded support in  $X$*  if the image of the support of  $\mu$  in  $X$  under the orbit map has bounded diameter. If  $\mu$  has bounded support in  $X$ , then there are constants  $\ell > 0$ ,  $K \geq 0$ , and  $c < 1$  such that the following estimates hold. The random walk has positive drift [MT14, Theorem 1.2], i.e. for all  $n \geq 0$ ,

$$(1) \quad \mathbb{P}(d_X(x_0, w_n x_0) \leq \ell n) \leq Kc^n.$$

Furthermore, there is exponential decay for the convolution measures of shadows [Mah12, Lemma 2.10], i.e. for all  $n$ , for all  $y \in X$ , and for all  $R \geq 0$ ,

$$(2) \quad \mu^{*n}(S_{x_0}(y, R)) \leq Kc^{d_X(x_0, y) - R}.$$

The following two estimates follow from exponential decay for shadows. For all  $n \in \mathbb{Z}$  and for all  $r \geq 0$ , we have the following estimate for the probability of a location of the sample path  $w_n x_0$  being far from  $\gamma_\omega$  [MT14, Proposition 5.7]:

$$(3) \quad \mathbb{P}(d_X(w_n x_0, \gamma_\omega) \geq r) \leq Kc^r.$$

Finally, for Gromov products we have the following estimate for all  $0 \leq i \leq n$  and all  $r \geq 0$ :

$$(4) \quad \mathbb{P}((x_0 | w_n x_0)_{w_i x_0} \geq r) \leq Kc^r.$$

A weaker form of this estimate is stated in [MT14, Proposition 5.9], but the proof essentially shows the version above. We give the argument below for the convenience of the reader.

*Proof of (4).* If  $(x_0 | w_n x_0)_{w_i x_0} \geq r$ , then  $x$  lies in a shadow  $S_{w_i x_0}(w_n x_0, R)$ , with  $d(w_i x_0, w_n x_0) - R \geq r + O(\delta)$ . The random variables  $w_i$  and  $w_i^{-1} w_n$  are independent, so by exponential decay of shadows (2), this occurs with probability at most  $Kc^{r+O(\delta)}$ .  $\square$

We remark that *a priori* the constants  $K$  and  $c$  in (1)–(4) above may all be different, but without loss of generality, we may choose  $K$  and  $c$  to be the largest of any of these constants, and then (1)–(4) hold for the same  $K$  and  $c$ .

**2.3. Oriented matches.** Given  $\kappa > 0$ , a quasigeodesic segment  $\gamma: J \rightarrow X$ , and a quasigeodesic  $\gamma': I \rightarrow X$ , we follow [DH] and say that  $\gamma'$  *crosses  $\gamma$  up to distance  $\kappa$*  if there exists an increasing map  $\theta: J \rightarrow I$  such that  $d_X(\gamma(t), \gamma'(\theta(t))) \leq \kappa$  for all  $t \in J$ .

**Proposition 2.1.** [DH17, Proposition 2.5] *Given non-negative constants  $\delta$  and  $Q$ , there is a constant  $\kappa > 0$  such that if  $G$  is a group acting by isometries on a  $\delta$ -hyperbolic space  $X$ , and if  $\mu$  is a nonelementary probability measure on  $G$  with bounded support in  $X$ , then for almost every sample path  $\omega \in (G, \mu)^{\mathbb{Z}}$  of the random walk, and all  $0 < \varepsilon < 1$ , there exists an integer  $N \geq 0$  such that for all integers  $n \geq N$ , the element  $w_n$  acts loxodromically on  $X$ , and for all geodesic segments*

$\gamma_n$  from  $x_0$  to  $w_n x_0$ , every  $Q$ -quasiaxis  $\alpha_{w_n}$  for  $w_n$  crosses a subsegment of  $\gamma_n$  of length at least  $(1 - \varepsilon)d_X(x_0, w_n x_0)$  up to distance  $\kappa$ .

This result is stated for  $\mu$  with finite support in [DH17, Proposition 2.5], but their proof only uses finite support to give positive drift with exponential decay, and exponential decay for the measures of shadows, so their result holds for  $\mu$  with bounded support in  $X$ .

Now suppose we have an isometric action  $G \curvearrowright X$  and let  $\gamma$  and  $\gamma'$  be quasigeodesics in  $X$ . We say that  $\gamma$  and  $\gamma'$  have an  $(L, \kappa)$ -oriented match if there is a subpath  $s \subset \gamma$  of diameter at least  $L$  and some  $h \in G$  such that  $h \cdot \gamma'$  crosses  $s$  up to distance  $\kappa$ .

The following proposition is essentially given in [MS17, Proposition 3.2]. However, there it is assumed that the action is acylindrical and the authors were not concerned with the orientation of the match.

**Proposition 2.2.** *Given non-negative constants  $\delta$  and  $Q$ , there is a constant  $\kappa_0$ , such that if  $G \curvearrowright X$  is a nonelementary action on a  $\delta$ -hyperbolic space,  $\mu$  is a nonelementary measure on  $G$  for this action, and  $g \in G$  is a loxodromic in the semi-group generated by the support of  $\mu$ , then for all  $L \geq 0$  and  $\kappa \geq \kappa_0$ , the probability that a  $Q$ -quasiaxis  $\alpha_{w_n}$  for  $w_n$  has an  $(L, \kappa)$ -oriented match with a  $Q$ -quasiaxis  $\alpha_g$  of  $g$  goes to 1 as  $n \rightarrow \infty$ .*

Proposition 2.2 follows from Proposition 2.1 and [MS17, Proposition 3.2.4]. Propositions 3.2.1–3.2.5 in [MS17] are stated for acylindrical actions, but acylindricity is only used in Propositions 3.2.1 and 3.2.2. Proposition 2.2 follows from the proof of [MS17, Proposition 3.2.4], which is stated for a more general definition of matching which allows orientation reversing matchings, but in fact the proof shows this by showing that the probability of an oriented match tends to one. We give the details for the convenience of the reader.

We shall start by showing that for any  $0 < \varepsilon < 1$ , the probability that  $\gamma_n$  has a match of size  $(1 - \varepsilon)|\gamma_n|$  with  $\gamma_\omega$  tends to one as  $n \rightarrow \infty$ .

**Proposition 2.3.** [MS17, Proposition 3.2.3] *Given non-negative constants  $\delta$  and  $Q$ , there is a constant  $\kappa > 0$  such that if  $G$  is a group acting by isometries on a  $\delta$ -hyperbolic space  $X$ , and  $\mu$  is a nonelementary probability measure on  $G$  with bounded support in  $X$ , then for almost every sample path  $\omega \in (G, \mu)^{\mathbb{Z}}$  of the random walk, and all  $0 < \varepsilon < 1$ , the probability that  $\gamma_n$  has a subsegment of length  $(1 - \varepsilon)|\gamma_n|$  contained in a  $\kappa$ -neighbourhood of  $\gamma_\omega$  tends to one as  $n \rightarrow \infty$ .*

*Proof.* Let  $\kappa_1$  be a Morse constant for  $Q$ -quasigeodesics in  $X$ , i.e. any geodesic connecting points of a  $Q$ -quasigeodesic is contained in a  $\kappa_1$ -neighbourhood of the  $Q$ -quasigeodesic.

As  $\mu$  has bounded support in  $X$ , there are constants  $\ell > 0$ ,  $K \geq 0$ , and  $c < 1$  such that the estimates for positive drift (1) and distance from  $\gamma_\omega$  (4) imply that the probability that both

$$(5) \quad d_X(x_0, \gamma_\omega) \leq \varepsilon_1 |\gamma_n| \text{ and } d_X(w_n x_0, \gamma_\omega) \leq \varepsilon_1 |\gamma_n|$$

is at least  $1 - 2Kc^{\varepsilon_1 \ell n} + Kc^n$ . Thus the probability that (5) holds tends to one as  $n \rightarrow \infty$ . If (5) holds, then by thin triangles,  $\gamma_n$  has a subgeodesic of length at least  $(1 - 2\varepsilon_1)|\gamma_n|$  contained in a  $(\kappa_1 + 2\delta)$ -neighbourhood of  $\gamma_\omega$ . Therefore the result holds with  $\varepsilon = 2\varepsilon_1$  and  $\kappa = \kappa_1 + 2\delta$ .  $\square$

We may now complete the proof of Proposition 2.2.

*Proof of Proposition 2.2.* Let  $\gamma_n$  be a geodesic from  $x_0$  to  $w_n x_0$ , let  $\alpha_{w_n}$  be a  $Q$ -quasiaxis for  $w_n$ , and let  $\gamma_\omega$  be a bi-infinite  $Q$ -quasigeodesic determined by the forward and backward limit points of a bi-infinite random walk  $\omega$  generated by  $\mu$ . Combining Propositions 2.1 and 2.3, there is a constant  $\kappa \geq 0$  such that for any  $0 < \varepsilon < 1$  the probability that  $\gamma_n$  has a subsegment of length  $(1 - \varepsilon)|\gamma_n|$  which is contained in a  $\kappa$ -neighbourhood of  $\alpha_{w_n}$ , and also in a  $\kappa$ -neighbourhood of  $\gamma_\omega$ , tends to one as  $n \rightarrow \infty$ .

Let  $g$  be a loxodromic element which lies in the semi-group generated by the support of  $\mu$ , which we shall denote  $\langle \text{supp}(\mu) \rangle_+$ , and let  $\alpha_g$  be a  $Q$ -quasi-axis for  $g$ . We now show that the probability that a subsegment of  $\gamma_\omega$  has a large match with  $\alpha_g$  tends to one as the length of the subsegment tends to infinity.

We shall write  $\nu$  for the harmonic measure on  $\partial X$ , and  $\check{\nu}$  for the reflected harmonic measure, i.e the harmonic measure arising from the random walk generated by the probability distribution  $\check{\mu}(g) = \mu(g^{-1})$ . By assumption, the group element  $g$  lies in  $\langle \text{supp}(\mu) \rangle_+$ , and so the group element  $g^{-1}$  lies in  $\langle \text{supp}(\check{\mu}) \rangle_+$ . By [MT14, Proposition 5.4], there is a constant  $R_0$  such that  $\nu(S_{x_0}(gx_0, R_0)) > 0$  for all  $g \in \langle \text{supp}(\mu) \rangle_+$ , and so also  $\check{\nu}(S_{x_0}(gx_0, R_0)) > 0$  for all  $g \in \langle \text{supp}(\check{\mu}) \rangle_+$ .

Given  $\delta$  and  $Q$ , there is a constant  $\kappa_0 \geq 0$  so that for any constants  $\kappa \geq \kappa_0$  and  $L \geq 0$ , there is an  $m$  sufficiently large such that any oriented  $Q$ -quasigeodesic  $\gamma$  from  $S_{x_0}(g^{-m}x_0, R_0)$  to  $S_{x_0}(g^m x_0, R_0)$  has a subsegment of length  $L$  which  $\kappa$ -fellow travels with the  $Q$ -quasi-axis  $\alpha_g$ , and whose orientation agrees with the orientation of  $\alpha_g$ . In particular, any such  $\gamma$  has an  $(L, \kappa)$ -match with  $\alpha_g$ .

As  $\nu(S_{x_0}(g^m x_0, R_0)) > 0$  and  $\check{\nu}(S_{x_0}(g^{-m}x_0, R_0)) > 0$ , there is a positive probability  $p$  so that  $\gamma_\omega$  has an oriented  $(L, \kappa)$ -match with  $\alpha_g$ . Ergodicity now implies that for any  $\varepsilon > 0$  the proportion of times  $\varepsilon n \leq k \leq (1 - \varepsilon)n$  for which  $\gamma_\omega$  has a subsegment of length at least  $L$  which lies in a  $K$ -neighbourhood of  $w_k \gamma_g$  tends to  $p$  as  $n \rightarrow \infty$ , for almost all sample paths  $\omega$ .

The final step is to relate the locations  $w_k x_0$  of the random walk to the geodesic  $\gamma_n$ , by showing that all of the  $w_k x_0$  for  $\varepsilon n \leq k \leq (1 - \varepsilon)n$  have nearest point projections to  $\gamma_n$  that are far from the endpoints, with asymptotic probability one:

**Claim.** For any  $\varepsilon > 0$ , the probability that all  $w_k x_0$  with  $4\varepsilon n \leq k \leq (1 - 4\varepsilon)n$  have nearest point projections to  $\gamma_n$  which lie distance at least  $\varepsilon|\gamma_n|$  from each endpoint tends to one as  $n \rightarrow \infty$ .

*Proof.* Let  $\ell > 0, K \geq 0$ , and  $c < 1$  be the constants from the estimates in (1)–(4). Exponential decay for Gromov products (4) implies that the probability that  $(x_0|w_n x_0)_{w_k x_0} \leq \lambda \log n$  for all  $1 \leq k \leq n$  is at least  $1 - nKc^{\lambda \log n}$ . For  $\lambda$  sufficiently large, this  $\rightarrow 1$ , as  $n \rightarrow \infty$ . Positive drift with exponential decay (1) implies that the probability that  $d_X(x_0, w_k x_0) \geq 2\varepsilon|\gamma_n|$  and  $d_X(w_k x_0, w_n x_0) \geq 2\varepsilon|\gamma_n|$  for all  $4\varepsilon n \leq k \leq (1 - 4\varepsilon)n$  is at least  $1 - 2nKc^{4\varepsilon n}$ , which  $\rightarrow 1$  as  $n \rightarrow \infty$ .

If the Gromov product  $(x_0|w_n x_0)_{w_k} \leq \varepsilon|\gamma_n|$ , and the nearest point projection of  $w_k x_0$  to  $\gamma_n$  is within distance  $\varepsilon|\gamma_n|$  of  $x_0$ , then  $d_X(x_0, w_k x_0) \leq 2\varepsilon|\gamma_n| + O(\delta)$ , and similarly for the other endpoint  $w_n x_0$ . Therefore, as the probability that the Gromov products  $(x_0|w_n x_0)_{w_k x_0} \leq \lambda \log n \leq \varepsilon|\gamma_n|$  for all  $k$  tends to one, this implies that the probability that all of the nearest point projections of  $w_k x_0$  to  $\gamma_n$ , for  $4\varepsilon n \leq k \leq (1 - 4\varepsilon)n$ , lie distance at least  $\varepsilon|\gamma_n|$  from the endpoints,  $\rightarrow 1$  as  $n \rightarrow \infty$ .  $\square$

This completes the proof of Proposition 2.2.  $\square$

### 3. OUTER SPACE AND THE ATTRACTING TREE FOR A FULLY IRREDUCIBLE

We assume throughout that  $r \geq 3$  is an integer, that  $F_r$  is the rank- $r$  free group, and that  $\text{Out}(F_r)$  is its outer automorphism group. Recall that  $\varphi \in \text{Out}(F_r)$  is *fully irreducible* if no positive power of  $\varphi$  fixes the conjugacy class of a nontrivial proper free factor of  $F_r$ . It is proved in [BH92] that fully irreducible outer automorphisms have *train track representatives*.

**3.1. Outer space CV.** Culler–Vogtmann Outer space was first defined in [CV86]. We refer the reader to [FM11, Bes14, Vog15] for background on Outer space and give only abbreviated discussion here. For  $r \geq 2$  we denote the (volume-one normalized) Outer space for  $F_r$  by CV. We think of points of CV in three different ways. First, points of CV are equivalence classes of volume-one marked metric graphs  $h : R_r \rightarrow \Gamma$  where  $R_r$  is the  $r$ -rose, where  $\Gamma$  is a finite volume-one metric graph with betti number  $b_1(\Gamma) = r$  and with all vertices of degree  $\geq 3$ , and where  $h$  is a homotopy equivalence called a *marking*. Second, a point of CV can be viewed as a minimal free and discrete



isometric action of  $F_r$  on an  $\mathbb{R}$ -tree  $T$  with the quotient metric graph of volume one (where the tree  $T$ , in the previous picture, is obtained as the universal cover of the metric graph  $\Gamma$ ). In this paper, by an  $F_r$ -tree we will mean an  $\mathbb{R}$ -tree  $T$  endowed with a minimal nontrivial isometric action of  $F_r$ , or sometimes the projective class of such an action (when talking about points of compactified Outer space). Third, a point  $T \in \text{CV}$  can be viewed as a length function  $\|\cdot\|_T : F_r \rightarrow \mathbb{R}_{\geq 0}$  where for  $w \in F_r$  the number  $\|w\|_T = \inf_{x \in T} d(x, wx)$  is the *translation length* of  $w$  in  $T$ . Each of these three descriptions, in the appropriate sense, uniquely defines a point of CV. One can also think of CV as the union of the open simplices obtained by varying the lengths of edges of  $\Gamma$  for a given  $h : R_r \rightarrow \Gamma$ . By abuse of notation, we will usually denote by  $\Gamma$  a point of CV given by  $h : R_r \rightarrow \Gamma$  and suppress the mention of the marking  $h$ .

Outer space CV admits a natural compactification  $\overline{\text{CV}} = \text{CV} \cup \partial\text{CV}$ . Points of  $\overline{\text{CV}}$  are projective (homothety) classes of minimal *very small*  $F_r$ -actions on  $\mathbb{R}$ -trees, [BF94],[CL95]. See [Pau89] for descriptions of several equivalent topologies on  $\overline{\text{CV}}$ . The space CV naturally embeds in  $\overline{\text{CV}}$  as an open dense  $\text{Out}(F_r)$ -invariant subset, where a point of CV is identified with its projective class.

We consider CV with the asymmetric Lipschitz metric (see, for example, [FM11] for background information). We briefly recall the definition here for completeness. If  $h_1 : R_r \rightarrow \Gamma_1$  and  $h_2 : R_r \rightarrow \Gamma_2$  are two points of CV, a continuous map  $f : \Gamma_1 \rightarrow \Gamma_2$  is called a *difference of markings* from  $\Gamma_1$  to  $\Gamma_2$  if  $f$  is freely homotopic to  $f_2 \circ f_1^{-1}$ . The *Lipschitz distance*  $d_{\text{CV}}(\Gamma_1, \Gamma_2)$  is defined as

$$d_{\text{CV}}(\Gamma_1, \Gamma_2) := \inf_f \log \text{Lip}(f)$$

where  $\text{Lip}(f)$  is the Lipschitz constant of  $f$  and where the infimum is taken over differences in markings  $f$  from  $\Gamma_1$  to  $\Gamma_2$ . It is known that  $d_{\text{CV}}$  is an asymmetric metric, in the sense of §2.1.

By a *geodesic* in CV we always mean a directed geodesic with respect to  $d_{\text{CV}}$ , i.e. a path  $\gamma : J \rightarrow \text{CV}$  (where  $J \subseteq \mathbb{R}$  is a subinterval) such that for any  $s < t$  in  $J$  we have  $d_{\text{CV}}(\gamma(s), \gamma(t)) = t - s$ . It is known that  $(\text{CV}, d_{\text{CV}})$  is a directed geodesic space, i.e. for any  $\Gamma_1, \Gamma_2 \in \text{CV}$ , there exists a geodesic (in the above sense) from  $\Gamma_1$  to  $\Gamma_2$  in CV.

It is sometimes convenient to symmetrize the metric  $d_{\text{CV}}$  to obtain a (nongeodesic) metric  $d_{\text{sym}}(x, y) = (d_{\text{CV}}(x, y) + d_{\text{CV}}(y, x))$ . We remark that both metrics  $d_{\text{CV}}$  and  $d_{\text{sym}}$  define the standard topology on CV [FM11], and, in particular,  $(\text{CV}, d_{\text{sym}})$  is locally compact.

**3.2. Train track maps.** We adopt the same conventions regarding train track maps and train track representatives of elements of  $\text{Out}(F_r)$  as in [AKKP18]. In particular, for a graph  $\Gamma$  (which we do not yet assume to be finite) and a vertex  $x$  of  $\Gamma$ , the *directions* at  $x$  in  $\Gamma$  are germs of initial segments of edges emanating from  $x$ . We refer the reader to [BH92, Bog08, Bes11, DKL15] for more detailed background on train track maps in the  $\text{Out}(F_r)$  context.

**3.3. Attracting and repelling trees**  $T_+^\varphi, T_-^\varphi$ . Let  $\varphi \in \text{Out}(F_r)$  be a fully irreducible outer automorphism. Then  $\varphi$  acts on  $\overline{\text{CV}}$  with “North-South dynamics” (see [LL03]). As elements of  $\partial\text{CV}$ , the attractor and repeller for the  $\text{Out}(F_r)$ -action are  $F_r$ -trees, we denote them respectively by  $T_+^\varphi$  and  $T_-^\varphi$ .

We recall from [GJLL98] a concrete construction of  $T_+^\varphi$ . Let  $g : \Gamma \rightarrow \Gamma$  be a train track representative of  $\varphi$  and  $\tilde{\Gamma}$  the universal cover of  $\Gamma$  equipped with a distance function  $\tilde{d}$  lifted from  $\Gamma$ . The fundamental group,  $F_r$ , acts by deck transformations, hence isometries, on  $\tilde{\Gamma}$ . A lift  $\tilde{g}$  of  $g$  is associated to the unique automorphism  $\Phi$  representing  $\varphi$  satisfying that, for each  $w \in F_r$  and  $x \in \tilde{\Gamma}$ , we have  $\Phi(w)\tilde{g}(x) = \tilde{g}(wx)$ . Define the pseudo-distance  $d_\infty$  on  $\tilde{\Gamma}$  by  $\lim_{k \rightarrow +\infty} d_k$ , where

$$d_k(x, y) = \frac{d(\tilde{g}^k(x), \tilde{g}^k(y))}{\lambda^k}$$

for each  $x, y \in \tilde{\Gamma}$ . Then  $T_+$  is the  $F_r$ -tree defined by identifying each pairs of points  $x, y \in \tilde{\Gamma}$  such that  $d_\infty(x, y) = 0$ . It is known from [BF94], for example, that  $T_+^\varphi$  is the  $F_r$ -equivariant Gromov limit of the trees  $\tilde{\Gamma}_k$  obtained from  $\tilde{\Gamma}$  by identifying distance-0 points in  $\tilde{\Gamma}$  endowed with the respective pseudo-distance  $d_k$ .

**Remark 3.1** (Left and right actions on CV and the direction of the folding lines). The standard action of  $\text{Out}(F_r)$  on CV and on  $\overline{\text{CV}}$  is a right action. At the level of length functions, this action is described as follows. For a tree  $T$ , an outer automorphism  $\varphi \in \text{Out}(F_r)$ , and an element  $w \in F_r$  we have  $\|w\|_{T\varphi} := \|\varphi(w)\|_T$ . In the language of actions on trees, for an  $F_r$ -tree  $T$  and an outer automorphism  $\varphi \in \text{Out}(F_r)$ , the  $F_r$ -tree  $T\varphi$  is defined as the tree  $T$  with the action of  $F_r$  twisted via  $\varphi$ : For  $x \in T$  and  $w \in F_r$  we have  $w \cdot_{T\varphi} x := \varphi(w) \cdot_T x$ .

If  $\varphi \in \text{Out}(F_r)$  is fully irreducible and  $T_+^\varphi \in \overline{\text{CV}}$  is the attracting tree of  $\varphi$ , then, projectively, we have  $\lim_{n \rightarrow \infty} T_0 \varphi^n = T_+^\varphi$  in  $\overline{\text{CV}}$ , for any  $T_0 \in \text{CV}$ . In particular, if  $\gamma$  is a periodic folding axis for  $\varphi$  in CV (see §5.1 below) with  $\lim_{t \rightarrow \infty} \gamma(t) = T_+^\varphi$  (see §5.1 below) then for every  $S \in \gamma$  we have  $\lim_{n \rightarrow \infty} S \varphi^n = T_+^\varphi$ .

There is also a standard way to convert this right action of  $\text{Out}(F_r)$  on CV and on  $\overline{\text{CV}}$  to a left action. Namely, for a tree  $T$  and an element  $\varphi \in \text{Out}(F_r)$  set  $\varphi \cdot T := T\varphi^{-1}$ . We need to work with both these right and left actions. Note that if  $\varphi \in \text{Out}(F_r)$  is fully irreducible with a periodic folding axis  $\gamma$  in CV (see the discussion of axes below) then for any  $S \in \gamma$  we have (in  $\overline{\text{CV}}$ ):

$$\lim_{t \rightarrow \infty} \gamma(t) = \lim_{n \rightarrow \infty} \varphi^{-n} S = \lim_{n \rightarrow \infty} S \varphi^n = T_+^\varphi.$$

**3.4. Geometric fully irreducible outer automorphisms and PNPs.** As in [LP97, Definition 0.1], we call an  $\mathbb{R}$ -tree *geometric* if it arises as the dual to a measured foliation on some finite simplicial 2-complex. A fully irreducible  $\varphi \in \text{Out}(F_r)$  is *geometric* if  $T_+^\varphi$  is not geometric. If  $T_+^\varphi$  is geometric and  $\varphi$  is induced by a surface homeomorphism, then  $\varphi$  is called *geometric*. One otherwise calls a fully irreducible outer automorphism  $\varphi$  *parageometric*, i.e. when  $T_+^\varphi$  is geometric, but  $\varphi$  is not induced by a surface homeomorphism.

While we do not directly use the definition of a (*periodic*) *Nielsen path*, or *PNP*, it will be relevant that each rotationless power of each geometric fully irreducible outer automorphism has a PNP-free train track representative. In particular, by [BF94, Theorem 3.2], for a fully irreducible  $\varphi \in \text{Out}(F_r)$ , we have that  $T_+^\varphi$  is geometric if and only if the [BH92] “stable” train track representative of  $\varphi$  contains a PNP. Using the “stable” PNP-free train track representative for an geometric fully irreducible outer automorphism allows us, for example, to use the simplified definition of a principal vertex stated in §4.2 and the simplified ideal Whitehead graph definition of §4.4.

#### 4. LAMINATIONS AND IDEAL WHITEHEAD GRAPHS

**Convention 4.1.** We continue to assume throughout this section that  $r \geq 3$  is an integer, that  $F_r$  is the rank- $r$  free group, and that  $\text{Out}(F_r)$  is its outer automorphism group. We add an assumption that  $\varphi \in \text{Out}(F_r)$  is an geometric fully irreducible outer automorphism and that  $g: \Gamma \rightarrow \Gamma$  is a PNP-free irreducible train track representative of  $\varphi$ . We use  $\tilde{g}: \tilde{\Gamma} \rightarrow \tilde{\Gamma}$  to denote a lift of  $g$  to the universal cover. Finally, for simplicity, we assume that  $\varphi$ , hence also all of its train track representatives, are *rotationless*.

The theory of rotationless (outer) automorphisms can be found in [FH11] or [HM11]. We will use the facts that each positive power of this rotationless  $g$  will itself be rotationless, and that anything that is fixed (in the sense of §4.2) by a positive power  $g^k$  will in fact be fixed by  $g$  itself. Thus, for  $g$  there is no difference between “periodic” and “fixed” for vertices, directions, etc.

We remark that the PNP-free and rotationless assumptions for  $\varphi$  and  $g$  are not restrictive since every fully irreducible outer automorphism has a positive rotationless power, and since (see §3.4) every ageometric fully irreducible outer automorphism admits a PNP-free train track representative.

**4.1. Laminations.** The attracting lamination,  $\Lambda_\varphi$ , for a fully irreducible  $\varphi \in \text{Out}(F_r)$  is defined in [BFH97]. The lamination records the limiting behavior of a cyclic word under repeated application of an automorphism, and hence has intimate connections to tiling and substitution theory, for example (see [ABH<sup>+</sup>06], [CHL08]). We will be most interested in its “singularity structure,” as described in §4.3 and §4.4.

Before proceeding further, we warn the reader of a bit of notational discrepancy in the literature. In most places  $\Lambda_\varphi$  comes endowed with a “+” sign, but in [HM11]  $\Lambda_\varphi$  comes instead endowed with a “−” sign. Since we only use the attracting lamination, we leave the sign out altogether.

The attracting lamination can equally be realized as certain unordered pairs (*leaves*) of distinct boundary points in  $F_r$ , in  $T_+^\varphi$ , or in the boundary of any point of CV whose quotient under the  $F_r$  action carries a train track representative for  $\varphi$  (points that are “train tracks” in the language of [HM11] or [MP16]).

For a definition of  $\Lambda_\varphi$  in terms of a train track representative  $g: \Gamma \rightarrow \Gamma$ , and most closely relevant to our context, we refer the reader to [HM11, pg. 36]. We give a brief description here. The *realization*  $\Lambda(g)$  of  $\Lambda_\varphi$  in  $\Gamma$  is the collection of all lines (*leaves*) realized as bi-infinite immersed edge-paths  $\alpha$  in  $\Gamma$

$$\alpha = \dots, e_{-n}, \dots, e_{-1}, e_0, e_1, \dots, e_n, \dots$$

(where  $e_i$  are edges of  $\Gamma$ ) with the property that for every finite subpath  $\beta$  of  $\alpha$  there exists an integer  $k \geq 1$  and an edge  $e$  of  $\Gamma$  such that  $\beta$  is a subpath of  $g^k(e)$ .

The lifts of leaves of  $\Lambda(g)$  to  $\tilde{\Gamma}$  are also referred to as lamination leaves. Every such lift connects a pair of distinct points in  $\partial\tilde{\Gamma}$  and the set of the resulting pairs forms a closed flip-invariant  $F_r$ -invariant subset of  $\partial^2\tilde{\Gamma} = \partial\tilde{\Gamma} \times \partial\tilde{\Gamma} - \Delta$  which gives another way of viewing  $\Lambda(g)$ . Via a natural identification  $\partial\tilde{\Gamma} \cong \partial F_r$  and, correspondingly,  $\partial^2\tilde{\Gamma} \cong \partial^2 F_r$ , this allows one to associate to  $\Lambda(g)$  a subset  $\Lambda_\varphi \subseteq \partial^2 F_r$  that turns out to be independent of the choice of  $g$  and to depend on  $\varphi$  only.

On pg. 29-30 of [HM11], Handel and Mosher define a “direct limit” map  $g_\infty: \tilde{\Gamma} \rightarrow T_+^\varphi$  (there called  $f_g$ ) that is an isometry when restricted to each edge of  $\tilde{\Gamma}$  (when  $\tilde{\Gamma}$  is equipped with the natural “eigenmetric” determined by the transition matrix of  $g$ ). The map  $g_\infty$  is well-defined up to isometric conjugacy of the  $F_r$  action on  $T_+^\varphi$ . By [HM11, Lemma 2.21] we know that the restriction of  $g_\infty$  to each leaf is an isometry and that the collection in  $T_+^\varphi$  of  $g_\infty$ -images of leaves is independent of the choice of train track representative  $g$  of  $\varphi$ . This collection of leaf images will be the *realization of  $\Lambda_\varphi$  in  $T_+^\varphi$* .

**4.2. Principal vertices and lifts.** The notions of principal points and principal lifts were first introduced in [FH11] to mimic the corresponding surface theory notions. We first establish certain relevant terminology.

Since both  $T_+^\varphi$  and  $\tilde{\Gamma}$  are  $\mathbb{R}$ -trees, we freely use  $\mathbb{R}$ -tree terminology. For a general  $\mathbb{R}$ -tree  $T$  we have the following. A point  $p \in T$  is called a *branch point* if  $T \setminus p$  has  $\geq 3$  components. Each of these components is called a *direction* at  $p$ . A *turn* at  $p$  is an unordered pair of directions  $\{d_i, d_j\}$  at  $p$ .

Recall that  $\varphi$  and  $g$  are as in Convention 4.1. A *g-fixed point* (which, since we are in the absence of PNPs, is necessarily a vertex) in  $\Gamma$  is *principal* if it has  $\geq 3$   $g$ -fixed directions. A point  $\tilde{v} \in \tilde{\Gamma}$  is called *principal* if it is a lift of a principal point or, equivalently, if some positive power of the lift  $\tilde{g}$  fixes  $\tilde{v}$  and  $\geq 3$  directions at  $\tilde{v}$ . (See, for example, [HM11, pg. 27-28]). A lift  $\tilde{g}: \tilde{\Gamma} \rightarrow \tilde{\Gamma}$  is *principal* when its boundary extension, denoted  $\hat{g}$ , has  $\geq 3$  nonrepelling fixed points.

**4.3. Local/stable Whitehead graphs.** In [HM11], Handel and Mosher define the ideal Whitehead graph for an arbitrary nongeometric fully irreducible  $\varphi \in \text{Out}(F_r)$ . Here we will restrict our consideration to the case of ageometric fully irreducible outer automorphism. since that is the only case relevant to the present paper. We recall that  $\varphi$  and  $g$  are as in Convention 4.1.

We define the following graphs:

- For a vertex  $v$  of  $\Gamma$ , the *local Whitehead graph*  $LW(g, v)$  has a vertex for each direction at  $v$  and an edge connecting the vertices corresponding to the pair of directions  $\{d_1, d_2\}$  if the turn  $\{d_1, d_2\}$  is taken by an image under some  $g^k$  of some, hence any, edge of  $\Gamma$  (equivalently is taken by a leaf of  $\Lambda(g)$ ).
- The *stable Whitehead graph*  $SW(g, x)$  at a principal vertex  $x$  is then the induced subgraph of  $LW(g, x)$  obtained by restricting to the  $g$ -fixed directions.

For a principal vertex  $\tilde{x} \in \tilde{\Gamma}$ , one similarly defines  $\widetilde{SW}(\tilde{x})$  to have a vertex for each direction at  $\tilde{x}$  fixed by the principal lift  $\tilde{g}: \tilde{\Gamma} \rightarrow \tilde{\Gamma}$  fixing  $\tilde{x}$  and an edge connecting the corresponding pair of vertices for each turn taken by a lamination leaf. See §3 of [HM11] for the detailed definition of  $\widetilde{SW}(\tilde{x})$ . Note, however, that the graph  $\widetilde{SW}(\tilde{x})$  is naturally isomorphic to  $SW(g, x)$ , where  $\tilde{x}$  is a lift of  $x$ .

**4.4. Ideal Whitehead graphs.** In §3 of [HM11] Handel and Mosher define the ideal Whitehead graph  $IW(\varphi)$  and give several equivalent descriptions of it. We use their description in the train track language (under our assumption of no PNPs) as the main definition. See [HM11, Lemma 3.1, Corollary 3.2] for additional details. Note that if  $\tilde{g}$  fixes two distinct vertices in  $\tilde{\Gamma}$  then the geodesic between them projects to a PNP in  $\Gamma$  [HM11, Lemma 3.1], contradicting our no PNP assumption on  $g$ . Thus the no PNPs assumption on  $g$  means that each principal lift  $\tilde{g}$  of  $g$  fixes a unique principal vertex  $\tilde{x}$  in  $\tilde{\Gamma}$ , which we call the *center* of  $\tilde{g}$ . We will also sometimes say that  $\tilde{g}$  is *centered* at  $\tilde{x}$ . In this case, we define<sup>2</sup>

$$W(\tilde{g}) := \widetilde{SW}(\tilde{x}).$$

Recall that two principal lifts  $\tilde{g}, \tilde{g}'$  of  $g$  are considered equivalent,  $\tilde{g} \sim \tilde{g}'$ , if there exists some  $w \in F_r$  such that  $\tilde{g}' = w \circ \tilde{g} \circ w^{-1}$  (for the left translation covering action of  $F_r$  on  $\tilde{\Gamma}$ ). In this case (again, under the no PNPs assumption), there is a bijective correspondence between equivalence classes of principal lifts of  $g$  and the set of principal vertices of  $\Gamma$ , see §2 of [HM11] for the details.

We now state the definition of the ideal Whitehead graph of  $\varphi$ , following the discussion on p. 42 in §3.1 of [HM11].

**Proposition-Definition 4.2** (Ideal Whitehead graph). We define the *ideal Whitehead graph* as follows.

- (1) Let  $\varphi$  and  $g$  be as in Convention 4.1.

We define the *ideal Whitehead graph*  $IW(\varphi)$  of  $\varphi$  as the disjoint union of the  $W(\tilde{g})$  taken over the equivalence classes of all principal lifts  $\tilde{g}$  of  $g$ .

In view of the discussion above and using [HM11, Lemma 3.1, Corollary 3.2], under our assumption of no PNPs, we have

$$IW(\varphi) = \sqcup SW(g, x)$$

where the union is taken over all principal vertices  $x \in \Gamma$ .

- (2) Handel and Mosher (see also [Pfa12]) prove that this definition of  $IW(\varphi)$  does not depend on the choice of  $g$ , is an invariant of the conjugacy class of  $\varphi$  in  $\text{Out}(F_r)$ , and is preserved under taking positive powers. Recall that each ageometric fully irreducible  $\varphi \in \text{Out}(F_r)$

<sup>2</sup>[HM11] give a different definition of  $W(\tilde{g})$  which in our setting is equivalent to the one given here.

admits an expanding irreducible train track representative with no PNPs, and therefore the above definition applies to all rotationless ageometric fully irreducible outer automorphisms.

- (3) Let  $\varphi \in \text{Out}(F_r)$  be an arbitrary ageometric fully irreducible outer automorphism. Then there exists some integer  $k \geq 1$  such that  $\varphi^k$  is rotationless. Set  $IW(\varphi) := IW(\varphi^k)$ .

**4.5. Ideal Whitehead graph and branch points in the attracting tree.** In [HM11], for a nongeometric fully irreducible  $\varphi \in \text{Out}(F_r)$ , Handel and Mosher also give an equivalent description of  $IW(\varphi)$  in terms of directions at branch points in  $T_+^\varphi$  and the turns “taken” by  $\Lambda_\varphi$  there. We will not need the full strength of this description here and will only state the simplified versions of their results required for our purposes. Recall that  $g_\infty$  is the map  $f_g$  of [HM11, pg. 29-30].

**Proposition 4.3.** *Let  $\varphi \in \text{Out}(F_r)$  be a rotationless ageometric fully irreducible outer automorphism represented by an expanding irreducible train track map  $g : \Gamma \rightarrow \Gamma$  with no PNPs. Let  $g_\infty : \tilde{\Gamma} \rightarrow T_+^\varphi$  be the  $F_r$ -equivariant “direct limit” map. Then the following hold:*

- (1) *Let  $\tilde{g}$  be a principal lift of  $g$  centered at a principal vertex  $\tilde{x}$  of  $\tilde{\Gamma}$ . Then  $b = g_\infty(\tilde{x})$  is a branch point of  $T_+^\varphi$  and the map  $g_\infty$  induces a bijection between the set of  $\tilde{g}$ -fixed directions at  $\tilde{x}$  in  $\tilde{\Gamma}$  (which are precisely the vertices of  $\widetilde{SW}(\tilde{x})$ ) and the set of directions (i.e. connected components of  $T_+^\varphi - \{b\}$ ) of  $T_+^\varphi$  at  $b$ .*
- (2) *Every branch point  $b \in T_+^\varphi$  arises as in part (1).*

*Proof.* Part (1) follows from [HM11, Lemma 3.4]. Part (2) follows from [HM11, Lemma 2.16(1)].  $\square$

**4.6. The rotationless index and an alternate ageometric characterization.** The notion of an index  $\text{ind}(\varphi)$  for an outer automorphism  $\varphi \in \text{Out}(F_r)$  was first introduced in [GJLL98]. This notion is not in general invariant under taking powers. [HM11] introduces the notion of a rotationless index (there just called the index sum)  $i(\varphi)$  for a fully irreducible  $\varphi \in \text{Out}(F_r)$ . Let  $\varphi$  be a nongeometric fully irreducible outer automorphism. For each component  $C_i$  of  $IW(\varphi)$ , let  $k_i$  denote the number of vertices of  $C_i$ . Then the *rotationless index* is defined as  $i(\varphi) := \sum 1 - \frac{k_i}{2}$ . It follows from [HM11, Lemma 3.4] that for a rotationless nongeometric fully irreducible  $\varphi \in \text{Out}(F_r)$ , the two notions differ only by a change of sign.

The rotationless index can be used to determine whether a fully irreducible outer automorphism is ageometric. In particular, a fully irreducible  $\varphi \in \text{Out}(F_r)$  is ageometric if and only if  $0 > i(\varphi) > 1 - r$ . While the characterization follows fairly easily from a sequence of previous results, it is finally more or less stated in [KP15] as Corollary 2.36. A justification and explanation of the fact can be found sprinkled throughout [KP15] or in [MP16, §2.9].

## 5. FOLD LINES, AXIS BUNDLES, AND GEODESICS IN CV

The proof of our main theorem requires us to understand the behavior of an axis of a random outer automorphism. As such, we need a detailed account of these axes.

**5.1. Fold lines and axis bundles.** Let  $\widehat{CV}$  denote the “unprojectivized” version of CV and  $p : \widehat{CV} \rightarrow CV$  the projection map. Then a *fold line* in CV is the image under  $p$  of a continuous, injective, proper function from  $\mathbb{R}$  to  $\widehat{CV}$  that is defined by a continuous 1-parameter family of marked graphs  $t \rightarrow \Gamma_t$  and a family of homotopy equivalences  $h_{ts} : \Gamma_s \rightarrow \Gamma_t$  defined for  $s \leq t \in \mathbb{R}$ , each marking-preserving, and satisfying:

- $h_{ts}$  is a local isometry on each edge for all  $s \leq t \in \mathbb{R}$  and
- $h_{ut} \circ h_{ts} = h_{us}$  for all  $s \leq t \leq u \in \mathbb{R}$  and  $h_{ss} : \Gamma_s \rightarrow \Gamma_s$  is the identity for all  $s \in \mathbb{R}$ .

A fold line in Outer space  $\mathbb{R} \rightarrow CV$  is said to be *simple* if there exists a subdivision of  $\mathbb{R}$  by points  $(t_i)_{i \in \mathbb{Z}}$

$$\dots t_{i-1} < t_i < t_{i+1} \dots$$

such that  $\lim_{i \rightarrow \infty} t_i = \infty$ ,  $\lim_{i \rightarrow -\infty} t_i = -\infty$ , and the following holds:

For each  $i \in \mathbb{Z}$  there exist distinct edges  $e, e'$  in  $\Gamma_{t_i}$ , with a common initial vertex, such that: For each  $s \in (t_i, t_{i+1}]$  the map  $h_{st_i}: \Gamma_{t_i} \rightarrow \Gamma_s$  identifies an initial segment of  $e$  with an initial segment of  $e'$ , with no other identifications (that is,  $h_{st_i}$  is injective on the complement of those two initial segments in  $\Gamma_{t_i}$ ).

**Remark 5.1.** By [AKKP18, Lemma 2.27], simple periodic fold lines in CV, defined by expanding irreducible train track representatives of outer automorphisms  $\varphi \in \text{Out}(F_r)$ , are geodesics in CV.

In [HM11], Handel and Mosher define the axis bundle of a fully irreducible, in analogy with the Teichmüller axis for a pseudo-Anosov:

Suppose that  $\varphi \in \text{Out}(F_r)$  is a nongeometric fully irreducible outer automorphism. Then the *axis bundle*  $\mathcal{A}_\varphi$  is the union of the images of all fold lines  $\gamma: \mathbb{R} \rightarrow \text{CV}$  such that  $\gamma(t)$  converges in  $\overline{\text{CV}}$  to  $T_-^\varphi$  as  $t \rightarrow -\infty$  and to  $T_+^\varphi$  as  $t \rightarrow +\infty$ .

We call the fold lines of the axis bundle *axes*. An axis  $\gamma$  in CV for  $\varphi$  is *periodic* if there exists an  $R > 0$  such that for every  $t \in \mathbb{R}$  we have  $\gamma(t + R) = \gamma(t)\varphi$ . Thus in this case  $\gamma$  is  $\varphi$ -invariant and for each  $x_0 \in \gamma$  we have  $\lim_{n \rightarrow \infty} x_0\varphi^n = T_+^\varphi$  and  $\lim_{n \rightarrow \infty} x_0\varphi^{-n} = T_-^\varphi$  in  $\overline{\text{CV}}$ . (We recall the conventions regarding right and left actions of  $\text{Out}(F_r)$  explained in Remark 3.1). In more generally understanding “periodic fold lines,” of particular relevance here are descriptions given in [AKKP18, Definitions 2.18-2.20], these notions date back to Stallings [Sta83] and Skora [Sko89].

In the circumstance where  $\mathcal{A}_\varphi$  is formed from only a single fold line, we call this fold line a *lone axis* and we say that  $\varphi$  is a *lone axis* fully irreducible outer automorphism. It is noteworthy that the lone axis will always be a periodic fold line for  $\varphi$ .

**Remark 5.2** (Fully irreducible outer automorphisms have simple axes). While not every periodic axis of a fully irreducible outer automorphism is simple, the axes constructed from Stallings fold decompositions, as explained in [AKKP18, Definitions 2.18-2.20] for example, are simple. In particular, every fully irreducible outer automorphism has at least one simple axis.

Mosher–Pfaff [MP16] gives a necessary and sufficient condition for an ageometric fully irreducible outer automorphism to have a lone axis.

**Theorem 5.3** ([MP16] Theorem 4.7). *The axis bundle of an ageometric fully irreducible outer automorphism  $\varphi \in \text{Out}(F_r)$  is a unique axis precisely if both of the following two conditions hold:*

- (1) *the rotationless index satisfies  $i(\varphi) = \frac{3}{2} - r$  and*
- (2) *no component of the ideal Whitehead graph  $IW(\varphi)$  has a cut vertex.*

One may note that, using the definition of §4.6, the first condition can be replaced by an ideal Whitehead graph condition.

**Remark 5.4** (Lone axes are simple greedy fold lines). It will be relevant throughout this paper that the lone axis of a lone axis fully irreducible outer automorphism is always a “greedy fold line” (also called a “fast folding path” in [FM11, Definition 5.7]). One can find the full definition of a greedy fold path in §2 of [BF14] and on pg. 33 of [DH17]. We omit the precise definition here but the idea is that at every time  $t$  of the folding process one folds all the turns that can be folded then, at uniform speed. The lone axis of a lone axis fully irreducible outer automorphism is greedy because at every time  $t$  there is a unique turn to fold. By [FM11, Theorem 5.6], greedy fold lines are geodesics in CV, and hence lone axes of lone axis fully irreducible outer automorphisms are geodesics as well. This fact provides another explanation of these axes being geodesic, alternative to the use of [AKKP18, Lemma 2.27] mentioned in Remark 5.1 above. The lone axis of a lone axis fully irreducible outer automorphism is also always a simple fold line.

The following lemma states that any geodesic which fellow travels an axis has the same forward and backward limits. It will be important later.

**Lemma 5.5.** *Suppose that  $\gamma$  is any geodesic axis of a nongeometric fully irreducible  $\varphi \in \text{Out}(F_r)$  and that a bi-infinite geodesic  $\gamma'$  satisfies that, for some  $\rho > 0$ , we have  $d_{\text{sym}}(\gamma(t), \gamma'(t)) \leq \rho$  for all  $t \in \mathbb{R}$ . Then  $\gamma'(t) \rightarrow T_+^\varphi$  as  $t \rightarrow +\infty$  and  $\gamma'(t) \rightarrow T_-^\varphi$  as  $t \rightarrow -\infty$ .*

*Proof.* Suppose that  $\lim_{t \rightarrow \infty} \gamma'(t) \neq T_+^\varphi$  in  $\overline{\text{CV}}$ . Since  $\overline{\text{CV}}$  is compact, there exists some  $T' \in \partial \text{CV}$ , with  $T' \neq T_+^\varphi$ , and a sequence  $t_i \geq 0$ , with  $\lim_{i \rightarrow \infty} t_i = \infty$ , such that  $\lim_{i \rightarrow \infty} \gamma'(t_i) = T'$  in  $\overline{\text{CV}}$ . By the assumptions on  $\gamma'$ , for each  $i \geq 1$ , we have that  $d_{\text{sym}}(\gamma'(t_i), \gamma(t_i)) \leq \rho$ . Since  $\lim_{i \rightarrow \infty} t_i = \infty$ , we have that  $\lim_{i \rightarrow \infty} \gamma(t_i) = T_+^\varphi$ . Because of the condition that  $d_{\text{sym}}(\gamma(t_i), \gamma'(t_i)) \leq \rho$ , it follows that for the (unprojectivized)  $\mathbb{R}$ -trees  $T_+^\varphi$  and  $T'$  there exists some  $c \geq 1$  such that for every  $w \in F_r$

$$\frac{1}{c} \|w\|_{T_+^\varphi} \leq \|w\|_{T'} \leq c \|w\|_{T_+^\varphi}.$$

Hence  $L^2(T_+^\varphi) = L^2(T')$  where  $L^2(\cdot)$  is the ‘‘dual lamination’’ of a tree in  $\overline{\text{CV}}$ , see [KL10] for the precise definition. Therefore, by [KL10, Theorem 1.3(2)], we have  $T' = T_+^\varphi$  in  $\overline{\text{CV}}$ . This contradicts the assumption that  $T' \neq T_+^\varphi$ . Thus  $\lim_{t \rightarrow \infty} \gamma'(t) = T_+^\varphi$ , as claimed. The argument for  $\lim_{t \rightarrow -\infty} \gamma'(t) = T_-^\varphi$  is similar.  $\square$

## 5.2. The space of geodesics.

**Definition 5.6** (Fellow travelling). Let  $R \geq 0$  and  $\rho \geq 0$ . Let  $\gamma : I \rightarrow \text{CV}$  and  $\gamma' : J \rightarrow \text{CV}$  be parameterized geodesics in  $\text{CV}$  (where  $I, J \subseteq \mathbb{R}$  are some intervals).

(1) Let  $t_0, t'_0 \in \mathbb{R}$  be such that  $[t_0, t_0 + R] \subseteq I$ , and  $[t'_0, t'_0 + R] \subseteq J$ , and for each  $s \in [0, R]$ ,

$$d_{\text{sym}}(\gamma(t_0 + s), \gamma'(t'_0 + s)) \leq \rho.$$

We then say that  $\gamma|_{[t_0, t_0 + R]}$  and  $\gamma'|_{[t'_0, t'_0 + R]}$   $\rho$ -fellow travel.

(2) We say that  $\gamma$  and  $\gamma'$   $\rho$ -fellow travel for length  $R$  if there exist  $t_0, t'_0$  such that  $\gamma|_{[t_0, t_0 + R]}$  and  $\gamma'|_{[t'_0, t'_0 + R]}$   $\rho$ -fellow travel.

(3) For a point  $x = \gamma(t_1)$  on  $\gamma$ , we say that  $\gamma$  and  $\gamma'$   $\rho$ -fellow travel for length  $R$  at a segment centered at  $x$  if there is some  $t'_0 \in \mathbb{R}$  such that  $\gamma|_{[t_1 - \frac{R}{2}, t_1 + \frac{R}{2}]}$  and  $\gamma'|_{[t'_0, t'_0 + R]}$   $\rho$ -fellow travel.

**Definition 5.7** (Space of geodesics  $Fl_r$ ). We denote by  $Fl_r$  the set of all parameterized bi-infinite geodesic fold lines  $\gamma : \mathbb{R} \rightarrow \text{CV}$  in  $\text{CV}$ . For  $\gamma \in Fl_r$ , as in [AKKP18], we let  $N(\gamma, \varepsilon)$  denote the  $\varepsilon$ -neighborhood of  $\gamma$  in  $\text{CV}$  with respect to the symmetrized Lipschitz metric on  $\text{CV}$ .

- (a) For  $R > 0, \varepsilon > 0$ , and  $\gamma \in Fl_r$ , we denote by  $B(\gamma, R, \varepsilon) \subset Fl_r$  the set of all  $\gamma' \in Fl_r$  such that  $\gamma$  and  $\gamma'$   $\varepsilon$ -fellow travel for length  $R$ .<sup>3</sup>
- (b) We topologize  $Fl_r$  by using, for each  $\gamma \in Fl_r$ , the family of sets  $\{B(\gamma, R, \varepsilon)\}_{\varepsilon > 0, R \geq 1}$  as the basis of neighborhoods of  $\gamma$  in  $Fl_r$ . We call this topology  $\tau$ . And we write  $\gamma_i \rightarrow \gamma$  to mean that a sequence of fold lines  $\{\gamma_i\}$  converges to  $\gamma$  in this topology.

The following lemma is useful in terms of understanding the topology  $\tau$  on  $Fl_r$ .

**Lemma 5.8.** *Let  $\gamma \in Fl_r$ , where  $\gamma$  is contained in some thick part of  $\text{CV}$ , and let  $R \geq 1, \varepsilon > 0$  be such that  $R > 4\varepsilon$ . Let  $\gamma' : J \rightarrow \text{CV}$  be a geodesic where  $J \subseteq \mathbb{R}$  is an interval of real numbers. Let  $t_1 < t_2$  and let  $t'_1, t'_2 \in J$  be such that  $t'_2 - t'_1 = R$ , such that  $d_{\text{sym}}(\gamma(t_1), \gamma'(t'_1)) \leq \varepsilon$ , such that  $d_{\text{sym}}(\gamma(t_2), \gamma'(t'_2)) \leq \varepsilon$ , and such that  $\gamma'([t'_1, t'_2]) \subseteq N(\gamma, \varepsilon)$ .*

*Then  $R - 2\varepsilon \leq t_2 - t_1 \leq R + 2\varepsilon$  and there exists a constant  $c > 0$ , depending only on the thickness of  $\gamma$ , and such that  $\gamma'|_{[t'_1, t'_2]}$  has a subsegment of length  $R - c\varepsilon$  which  $c\varepsilon$ -fellow travels a subsegment of  $\gamma$  of length  $c\varepsilon$ .*

<sup>3</sup>Note that this definition of  $B(\gamma, R, \varepsilon)$  is slightly different from the definition of  $B(\gamma, R, \varepsilon)$  given in [AKKP18], where the issue of directions on  $\gamma$  and  $\gamma'$  was ignored due to insufficiently careful phrasing. The definition of  $B(\gamma, R, \varepsilon)$  is the one which is actually meant in [AKKP18] and is needed in the present paper.

In particular, if  $J = \mathbb{R}$  and  $\gamma' \in FL_r$ , then  $\gamma' \in B(\gamma, R - c\varepsilon, c\varepsilon)$ .

*Proof.* Since  $\gamma$  is contained in some thick part of CV, there exists a constant  $c_0 > 0$  depending only on the thickness of  $\gamma$  (see [AKB12, Theorem 24]) such that for any  $s_1, s_2 \in \mathbb{R}$  we have

$$d_{CV}(\gamma(s_1), \gamma(s_2)) \leq d_{\text{sym}}(\gamma(s_1), \gamma(s_2)) \leq c_0|s_1 - s_2|.$$

Using the triangle inequality for  $d_{CV}$  and the fact that  $d_{CV} \leq d_{\text{sym}}$  we see that  $R - 2\varepsilon \leq t_2 - t_1 \leq R + 2\varepsilon$ .

Moreover, for the same reason, if  $s, s' \in [0, R]$  are such that

$$(\ddagger) \quad d_{\text{sym}}(\gamma'(t'_1 + s'), \gamma(t_1 + s)) \leq \varepsilon,$$

then  $|s - s'| \leq 2\varepsilon$ . Hence, by the choice of  $c_0$ , in the case of  $(\ddagger)$  we have

$$(\ddagger\ddagger) \quad d_{\text{sym}}(\gamma'(t'_1 + s'), \gamma(t_1 + s')) \leq 2\varepsilon(1 + c_0).$$

Let

$$Z = \{s' \in [0, R] \mid \exists s \in [0, R] \text{ with } d_{\text{sym}}(\gamma'(t'_1 + s'), \gamma(t_1 + s)) \leq \varepsilon\}.$$

Thus  $Z$  is a closed subset of  $[0, R]$  containing 0 and  $R$ . Set  $\Omega = [0, R] \setminus Z$ . Then  $\Omega$  is an open subset of  $[0, R]$ , not containing 0 or  $R$ , so that  $\Omega$  must be a disjoint union of open intervals. Suppose that  $J = (a', b')$ , with  $a' < b'$ , is one of these intervals. Thus  $a', b' \in Z$ . Hence, by  $(\ddagger\ddagger)$ ,  $d_{\text{sym}}(\gamma'(t'_1 + a'), \gamma(t_1 + a')) \leq 2\varepsilon(1 + c_0)$  and  $d_{\text{sym}}(\gamma'(t'_1 + b'), \gamma(t_1 + b')) \leq 2\varepsilon(1 + c_0)$ .

Recall that, by our assumptions,  $\gamma'([t'_1, t'_2]) \subseteq N(\gamma, \varepsilon)$ . Thus, since  $J \subseteq [0, R] \setminus Z$ , either

$$d_{\text{sym}}(\gamma'(t'_1 + a'), \gamma((-\infty, t_1])) \leq \varepsilon \text{ or } d_{\text{sym}}(\gamma'(t'_1 + a'), \gamma([t_2, \infty))) \leq \varepsilon$$

and, similarly, either

$$d_{\text{sym}}(\gamma'(t'_1 + b'), \gamma((-\infty, t_1])) \leq \varepsilon \text{ or } d_{\text{sym}}(\gamma'(t'_1 + b'), \gamma([t_2, \infty))) \leq \varepsilon.$$

We consider separately 3 cases.

**Case 1.** There exist  $y_a, y_b < t_1$  such that  $d_{\text{sym}}(\gamma(y_a), \gamma'(t'_1 + a')) \leq \varepsilon$  and  $d_{\text{sym}}(\gamma(y_b), \gamma'(t'_1 + b')) \leq \varepsilon$ .

Then  $d_{\text{sym}}(\gamma(y_a), \gamma(t_1 + a')) \leq \varepsilon + 2\varepsilon(1 + c_0)$  and  $d_{\text{sym}}(\gamma(y_b), \gamma(t_1 + b')) \leq \varepsilon + 2\varepsilon(1 + c_0)$ . Hence  $|t_1 + a' - y_a| \leq \varepsilon + 2\varepsilon(1 + c_0)$  and  $|t_2 + b' - y_b| \leq \varepsilon + 2\varepsilon(1 + c_0)$  which implies that  $a', b' \leq \varepsilon + 2\varepsilon(1 + c_0)$ .

**Case 2.** There exist  $y_a, y_b > t_2$  such that  $d_{\text{sym}}(\gamma(y_a), \gamma'(t'_1 + a')) \leq \varepsilon$  and  $d_{\text{sym}}(\gamma(y_b), \gamma'(t'_1 + b')) \leq \varepsilon$ .

Then by a similar argument as in Case 1, we see that  $|R - a'|, |R - b'| \leq \varepsilon + 2\varepsilon(1 + c_0)$ .

**Case 3.** There exist values  $y_1 \leq t_1$  and  $y_2 \geq t_2$  such that either

$$d_{\text{sym}}(\gamma(y_1), \gamma'(t'_1 + a')) \leq \varepsilon \text{ and } d_{\text{sym}}(\gamma(y_2), \gamma'(t'_1 + b')) \leq \varepsilon,$$

or

$$d_{\text{sym}}(\gamma(y_1), \gamma'(t'_1 + b')) \leq \varepsilon \text{ and } d_{\text{sym}}(\gamma(y_2), \gamma'(t'_1 + a')) \leq \varepsilon.$$

Let

$$Q_1 = \{q \in [a', b'] \mid \exists y \leq t_1 \text{ with } d_{\text{sym}}(\gamma(y), \gamma'(t'_1 + q)) \leq \varepsilon\}$$

and

$$Q_2 = \{q \in [a', b'] \mid \exists y \geq t_2 \text{ with } d_{\text{sym}}(\gamma(y), \gamma'(t'_1 + q)) \leq \varepsilon\}.$$

Thus  $Q_1, Q_2$  are nonempty closed subsets of  $[a', b']$  with  $Q_1 \cup Q_2 = [a', b']$ . Since  $[a', b']$  is connected, it follows that there exists some  $q \in Q_1 \cap Q_2$ . For this  $q$  there exist  $y_1 \leq t_1$  and  $y_2 \geq t_2$  such that  $d_{\text{sym}}(\gamma(y_1), \gamma'(t'_1 + q)) \leq \varepsilon$  and  $d_{\text{sym}}(\gamma(y_2), \gamma'(t'_1 + q)) \leq \varepsilon$ . Hence

$$R - 2\varepsilon \leq |t_2 - t_1| \leq |y_2 - y_1| \leq d_{\text{sym}}(\gamma(y_1), \gamma(y_2)) \leq 2\varepsilon,$$

which contradicts our assumption that  $R > 4\varepsilon$ . Thus Case 3 cannot occur.

We have proved that  $[\varepsilon(3+2c_0), R - \varepsilon(3+2c_0)] \subseteq Z$ . By  $(\ddagger\ddagger)$  for every  $s' \in [\varepsilon(3+2c_0), R - \varepsilon(3+2c_0)]$  we have  $d_{\text{sym}}(\gamma'(t'_1 + s'), \gamma(t_1 + s')) \leq 2\varepsilon(1 + c_0)$ .

Thus, with  $c = 3 + 2c_0$ , we see that  $\gamma'|_{[t'_1, t'_2]}$  has a segment of length  $R - c\varepsilon$  that  $c\varepsilon$ -fellow travels a subsegment of length  $R - c\varepsilon$  of  $\gamma$ , as required.  $\square$



We conclude this subsection with the following lemma, which is our main tool for showing that geodesics converge. The reader should note the very important hypothesis that  $\varphi$  has a lone axis.

**Lemma 5.9.** *Suppose  $\varphi \in \text{Out}(F_r)$  is fully irreducible with lone axis  $\gamma$ , that  $x$  is a point on  $\gamma$ , that  $\rho \geq 0$ , and that  $D_i$  is a sequence of positive numbers with  $D_i \rightarrow \infty$ . Suppose further that  $(\gamma_i)_{i \geq 0}$  is a sequence of simple periodic fold lines determined by expanding irreducible train track maps and satisfying that  $\gamma_i$   $\rho$ -fellow travels  $\gamma$  along a length  $D_i$ -interval centered at  $x$ . Then  $\gamma_i \rightarrow \gamma$  in  $FL_r$ .*

*Proof.* We prove convergence by showing that any subsequence  $\{\gamma_{i_j}\}_{j \in \mathbb{N}}$  of  $\{\gamma_i\}$  has a subsequence converging to  $\gamma$ . Since  $(CV, d_{\text{sym}})$  is proper, each  $B(D_{i_j}, \rho)$  is compact, so  $\{\gamma_{i_j}\}_j$  has finer and finer subsequences  $\{\gamma_{i_k}\}_k$  converging pointwise in the respective  $B(D_{i_j}, \rho)$ . This allows us, by diagonalization, to take a subsequence of  $\{\gamma_{i_j}\}_j$  converging to a fold line  $\gamma'$  in  $FL_r$ . (That the limit  $\gamma'$  is indeed a fold line comes from [BR15, Lemma 7.3].) By the assumptions on the  $\gamma_i$ , we know  $\gamma'$  must be a fold line in  $N(\gamma, \rho)$ . Moreover, by Remark 5.1, the  $\gamma_i$  are geodesics and therefore  $\gamma'$  is a geodesic as well. Then Lemma 5.5 implies that  $\gamma'$  shares endpoints with  $\gamma$ . We conclude that  $\gamma'$  is in the axis bundle of  $\varphi$ . Since  $\varphi$  has only a lone axis,  $\gamma' = \gamma$  and the proof is complete.  $\square$

**5.3. Principal and triangular automorphisms.** Our main theorem can be informally stated as saying that a random element of  $\text{Out}(F_r)$  is triangular. We explain this terminology here.

Let  $\Delta_r$  denote the graph consisting of the disjoint union of  $2r - 3$  triangles. As in [AKKP18], we call an ageometric fully irreducible  $\varphi \in \text{Out}(F_r)$  with  $IW(\varphi) \cong \Delta_r$  *principal*.

By the main theorem of [MP16], principal outer automorphisms are lone axis outer automorphisms. We say their axis is in the *principal stratum* of  $Fl_r$ . If for an ageometric fully irreducible  $\varphi \in \text{Out}(F_r)$  the graph  $IW(\varphi)$  is a disjoint union of  $\leq 2r - 3$  triangles, we call  $\varphi$  *triangular* (called *principal basin* in [AKKP18]).

**Remark 5.10.** In the present paper we only defined the notion of  $IW(\varphi)$  for an ageometric fully irreducible  $\varphi \in \text{Out}(F_r)$ . Thus for us principal and triangular fully irreducible outer automorphisms are ageometric by definition. However, in [HM11], Handel and Mosher defined  $IW(\varphi)$  in the more general context of any nongeometric fully irreducible  $\varphi \in \text{Out}(F_r)$ . If  $\varphi \in \text{Out}(F_r)$  is a nongeometric fully irreducible outer automorphism with  $IW(\varphi)$  being a disjoint union of  $\leq 2r - 3$  triangles, then for the rotationless index of  $\varphi$  we have  $i(\varphi) < 1 - r$  and therefore  $\varphi$  is ageometric in this case. Thus we could have omitted the ageometric assumption in the definition of principal and replaced it by “nongeometric”, and the ageometric property for principal and triangular fully irreducible outer automorphisms would still have been automatically satisfied in that case.

It is proved in [AKKP18] that principal outer automorphisms exist in each rank:

**Proposition 5.11** ([AKKP18] Example 6.1). *For each  $r \geq 3$ , there exists a principal  $\varphi \in \text{Out}(F_r)$ .*

Before proceeding, we record a key consequence of being a triangular outer automorphism.

**Proposition 5.12.** *Let  $\varphi \in \text{Out}(F_r)$  be a triangular fully irreducible outer automorphism. Then every branch point in  $T_+^\varphi$  is trivalent.*

*Proof.* By replacing  $\varphi$  by a positive power, we may assume that  $\varphi$  is rotationless (note that this operation does not change  $T_+^\varphi$  or  $IW(\varphi)$ ; in particular the outer automorphism remains triangular). Recall that all triangular outer automorphisms are ageometric fully irreducible. As noted before, by [BF94, Theorem 3.2] every ageometric fully irreducible outer automorphism admits a train track representative without PNPs. Thus we can represent  $\varphi$  by a rotationless expanding irreducible train track map  $g : \Gamma \rightarrow \Gamma$  with no PNPs.

Let  $b \in T_+^\varphi$  be a branch point. Then, by Proposition 4.3(2), we have  $b = g_\infty(\tilde{x})$  for a principal vertex  $\tilde{x} \in \tilde{G}$  that is the center of a principal lift  $\tilde{g}$  of  $g$ . Proposition 4.3(1) then implies that  $g_\infty$

induces a bijection between the set of directions at  $b$  in  $T_+^\varphi$  and the set of  $\tilde{g}$ -fixed directions in  $\tilde{G}$ . If  $\tilde{x}$  is a lift of the (principal vertex)  $x \in \Gamma$  then  $\widetilde{SW}(\tilde{x}) \cong SW(g, x)$  and there is a bijection between the set of  $\tilde{g}$ -fixed directions at  $\tilde{x}$  and the set of  $g$ -fixed directions at  $x$ . By the assumptions on  $\varphi$ ,

$$IW(\varphi) = \sqcup SW(g, v)$$

where the union is taken over all principal vertices  $v$  of  $\Gamma$ . Thus  $SW(g, x)$  is one of the triangular components of  $IW(\varphi)$ , which implies that  $\widetilde{SW}(\tilde{x})$  has three vertices, and therefore  $T_+^\varphi$  has three directions at  $b$ . It follows that  $b$  is trivalent, as required.  $\square$

The following is a consequence of [AKKP18, Theorem A], see Corollary 5.4 in [AKKP18].

**Proposition 5.13** ([AKKP18] Theorem A). [AKKP18, Corollary 5.4] *Suppose that  $r \geq 3$  and  $\gamma$  is the axis in  $\text{CV}$  of a principal  $\varphi \in \text{Out}(F_r)$ . Then there exist constants  $R, \varepsilon > 0$  so that for any simple periodic fold line  $\gamma' \in B(\gamma, R, \varepsilon)$ , we have that  $\gamma'$  is an axis of a triangular fully irreducible  $\varphi' \in \text{Out}(F_r)$ .*

The following is a strengthening of Proposition 5.13 using Lemma 5.9.

**Corollary 5.14.** *Let  $\varphi \in \text{Out}(F_r)$  be a principal (and hence lone axis) fully irreducible outer automorphism with axis  $\gamma$  in  $\text{CV}$ . Let  $\rho \geq 0$  be arbitrary. Then there exists some value  $R \geq 1$  such that, whenever  $\psi$  is a fully irreducible outer automorphism with an axis  $\gamma'$  satisfying that  $\gamma' \in B(\gamma, R, \rho)$ , we have that  $\psi$  is a triangular fully irreducible outer automorphism.*

*Proof.* By Proposition 5.13, there exist values  $\varepsilon \geq 0$  and  $R' \geq 1$  such that whenever a fully irreducible  $\psi \in \text{Out}(F_r)$  has an axis  $\gamma' \in B(\gamma, R', \varepsilon)$ , we have that  $\psi$  is triangular.

We now argue by contradiction. Suppose the conclusion of the corollary fails. Then there exists a sequence of positive real numbers  $R_i \rightarrow \infty$  as  $i \rightarrow \infty$  and a sequence of fully irreducible  $\psi_i \in \text{Out}(F_r)$  with axes  $\gamma_i \in B(\gamma, R_i, \rho)$  such that for every  $i \geq 1$ , we have that  $\psi_i$  is not triangular.

Since  $\gamma$  is  $\varphi$ -periodic and a period of  $\varphi$  is a compact interval, by choosing a specific point  $x \in \gamma$  and replacing each  $\gamma_i$  by some translate by a power of  $\varphi$  (which corresponds to replacing  $\psi_i$  by its conjugate by that power of  $\varphi$ ), we may assume that each  $\gamma_i$   $\rho$ -follows  $\gamma$  along a length  $D_i$ -interval centered at  $x$ , where  $\lim_{i \rightarrow \infty} D_i = \infty$ . Then, by Lemma 5.9, we have  $\gamma_i \rightarrow \gamma$ . This implies that for a sufficiently large index  $i_0 \geq 1$  we have that  $\gamma_{i_0} \in B(\gamma, R', \varepsilon)$ , and therefore  $\psi_{i_0}$  is a triangular fully irreducible outer automorphism, yielding a contradiction.  $\square$

## 6. THE FREE FACTOR GRAPH $\mathcal{FF}$

The *free factor graph*  $\mathcal{FF}$  for  $F_r$  is the graph having a vertex for each conjugacy class of proper nontrivial free factors of  $F_r$ . When either  $A \subsetneq B$  or  $B \subsetneq A$  an edge connects the vertices  $[A]$ ,  $[B]$ . The free factor graph  $\mathcal{FF}$  was proven to be Gromov hyperbolic by Bestvina and Feighn in [BF14] (this is also implied by [KR14] applied to [HM13]).

Recall from Reynolds [Rey12] that, for a point  $T \in \partial\text{CV}$ , a proper free factor  $A$  of  $F_r$  is a *reducing factor* for  $T$  if there exists an  $A$ -invariant subtree  $T'$  in  $T$  such that the action of  $A$  on  $T'$  has a dense orbit. In particular, if  $A$  is a free factor that acts elliptically on  $T$  fixing a point  $p$  in  $T$ , then  $A$  is a reducing factor, by taking  $T' = \{p\}$ . We denote the set of reducing factors of  $T$  by  $R(T)$ . The tree  $T$  is *arational* if it has no reducing free factors.

We let  $\mathcal{AT}$  denote the set of *arational trees*  $T \in \partial\text{CV}$  (see [Rey12]). Bestvina and Reynolds [BR15, Theorem 1.1], and independently Hamenstädt [Ham12], proved that  $\partial\mathcal{FF}$  is homeomorphic to equivalence classes of arational trees. This is more precisely explained in what follows.

## 6.1. The projection map $\pi: \text{CV} \rightarrow \mathcal{FF}$ .

**Definition 6.1** (Projection map  $\pi: \text{CV} \rightarrow \mathcal{FF}$ ). We let  $\pi: \text{CV} \rightarrow \mathcal{FF}$  denote the *projection map* defined by sending  $T$  to the collection of conjugacy classes of proper free factors that act elliptically on a simplicial tree obtained by equivariantly collapsing certain edges in  $T$  to points. In terms of the graph description of  $\text{CV}$ ,  $\pi(T/\mathbb{F}_r)$  is sent to the collection of conjugacy classes of nontrivial free factors that arise as the fundamental group of a proper subgraph of  $T/\mathbb{F}_r$ . We will use extensively the observation that all points in the same open simplex of  $\text{CV}$  have the same projection under  $\pi$ .

Using the main results from Bestvina–Reynolds [BR15], we follow [DH17, Theorem 5.5] and extend  $\pi$  to a map, which we also call  $\pi$  (called  $\psi$  in [DH17]),  $\pi: \overline{\text{CV}} \rightarrow \mathcal{FF} \cup \partial_\infty \mathcal{FF}$  satisfying: There exists a value  $\eta \in \mathbb{R}$  so that

- for each  $T \in \mathcal{AT}$  and sequence  $\{T_n \in \text{CV}\}$  converging to  $T$ , the sequence  $\{\pi(T_n)\}$  converges to  $\pi(T) \in \partial_\infty \mathcal{FF}$  and
- for each  $T \in \overline{\text{CV}} \setminus \mathcal{AT}$  and greedy fold line  $\gamma: \mathbb{R}_+ \rightarrow \text{CV}$ , if  $\gamma(t)$  converges to  $T$  as  $t \rightarrow +\infty$  in  $\overline{\text{CV}}$ , then there exists an  $N \in \mathbb{R}_+$  so that  $d(\pi(\gamma(t)), \pi(T)) \leq \eta$  for each  $t > N$ .

As in [BF14, pg. 115], for  $\Gamma, \Gamma' \in \text{CV}$ , we define

$$(6) \quad d_{\mathcal{FF}}(\Gamma, \Gamma') := \sup\{d_{\mathcal{FF}}(A, A') \mid A \in \pi(\Gamma), A' \in \pi(\Gamma')\}.$$

With these definitions,  $\pi: \text{CV} \rightarrow \mathcal{FF}$  is  $\text{Out}(F_r)$ -equivariant and coarsely  $L_{\text{CV}}$ -Lipschitz [BF14, Proposition 9.3], for some constant  $L_{\text{CV}}$  depending only on the rank of the fixed free group. In the course of establishing hyperbolicity of  $\mathcal{FF}$ , Bestvina–Feighn also showed that  $\pi$ -images in  $\mathcal{FF}$  of geodesics in  $\text{CV}$  are uniform unparametrized quasi-geodesics [BF14, Corollary 6.5]. That is, they can be reparameterized to be  $Q$ -quasigeodesics for some  $Q \geq 1$  depending only on the rank of  $F_r$ .

It follows from the [DH17, Theorem 5.5] construction and [BR15, Lemma 7.1] that if  $T \in \partial \text{CV}$  is not arational, then  $\pi(T)$  is by definition the set of reducing factors  $R(T)$  of  $T$ . By work of Bestvina–Reynolds [BR15], there is a constant  $\mathfrak{d} \geq 0$  such that if  $S, T \in \overline{\text{CV}}$  with  $R(T) \cap R(S) \neq \emptyset$ ,

$$(7) \quad d_{\mathcal{FF}}(R(T), R(S)) \leq \mathfrak{d}.$$

The constants  $L_{\text{CV}}$ ,  $Q$ , and  $\mathfrak{d}$  established here will be important in this paper.

**6.2. Nearest-point projection and  $\text{Pr}_\gamma$ .** Recall from §2.1 the definition of a Gromov product and that, since  $\mathcal{FF}$  is  $\delta$ -hyperbolic, given points  $x, y, z \in \mathcal{FF}$ , we have that

$$(8) \quad |d(x, \mathbf{n}_{[x,y]}(z)) - (x|y)_z| < 2\delta,$$

where  $\mathbf{n}_{[x,y]}(z)$  denotes the nearest-point projection of  $z$  to the geodesic segment  $[x, y]$  connecting  $x$  and  $y$  in  $\mathcal{FF}$  and  $\delta$  is the hyperbolicity constant of  $\mathcal{FF}$ .

Suppose that  $\gamma: I \rightarrow \text{CV}$  is a greedy folding path. As in [DT17], we use  $\mathbf{n}_{\pi \circ \gamma}$  to denote the nearest-point projection to the image of  $\pi \circ \gamma$  in  $\mathcal{FF}$ .

With the knowledge that a greedy folding path is a standard geodesic, the following is a direct consequence of [DT17, Lemma 4.2]. Except for where references are given, the lemma will be the extent of what we need to know about the projection function  $\text{Pr}_\gamma: \text{CV} \rightarrow \gamma(I)$  of [BF14, §6]. Note that [DT17, §4.1] is also dedicated to explaining the function. We need only the following lemma:

**Lemma 6.2** ([DT17] Lemma 4.2). *There exists a constant  $\mathfrak{c} \geq 0$ , depending only on the rank  $r$  of  $F_r$ , so that for any  $T \in \text{CV}$  and a greedy folding path  $\gamma: I \rightarrow \text{CV}$ , we have that*

$$d_{\mathcal{FF}}(\pi(\text{Pr}_\gamma(T)), \mathbf{n}_{\pi \circ \gamma}(\pi(T))) \leq \mathfrak{c}.$$

For this reason, we will at times in statements replace  $\pi(\text{Pr}_\gamma(T))$  with  $\mathbf{n}_{\pi \circ \gamma}(\pi(T))$  or vice versa.

The following lemma, Lemma 6.3, is indirectly stated in the proof of [BR15, Proposition 8.5] and then directly stated as [DH17, Proposition 5.11].

**Lemma 6.3** ([DH17] Proposition 5.11). *There exists a value  $M > 0$  so that for all sufficiently large  $B > 0$ , all  $y_0, y_1 \in \text{CV}$ , all trees  $T \in \overline{\text{CV}}$ , and all sequences  $(T_n)$  in  $\text{Out}(F_r)$  converging to  $T$  we have: If  $(\pi(T_n)) \mid \pi(y_1))_{\pi(y_0)} \geq B$  for all  $n \geq 0$ , then  $(\pi(T)) \mid \pi(y_1))_{\pi(y_0)} \geq B - M$ .*

## 7. THE BOUNDED GEODESIC IMAGE PROPERTY FOR FULLY IRREDUCIBLE AXES

In this section, we show that greedy folding axes of fully irreducible outer automorphism have the bounded geodesic image property in  $\text{CV}$ . Informally, this means that geodesics which are far from the axis have bounded diameter projection to the axis in  $\text{CV}$ . For honest (symmetric) metric spaces, the bounded geodesic image property is equivalent to the strong contracting property [ACT15] (strong contracting was established in [AK11] for axes in  $\text{CV}$ ). However, for the asymmetric metric spaces the contracting property does not appear to directly imply the bounded geodesic image property and additional, situation specific arguments, are required. Here, we rely on recent work of Dahmani and Horbez [DH17].

**Definition 7.1** (Bounded geodesic image property). A greedy folding path  $\gamma$  in  $\text{CV}$  has the *bounded geodesic image property* if there exist constants  $C_1$  and  $C_2$  so that if points  $X, Y \in \text{CV}$  satisfy  $d_{\text{sym}}(\text{Pr}_\gamma(X), \text{Pr}_\gamma(Y)) \geq C_1$  and satisfy  $\text{Pr}_\gamma(X) = \gamma(t_1)$ ,  $\text{Pr}_\gamma(Y) = \gamma(t_2)$  for some  $t_1 < t_2$ , then any geodesic segment  $[X, Y]$  between them contains a subsegment  $[Z_1, Z_2]$  such that

$$d_{\text{sym}}(Z_1, \text{Pr}_\gamma(X)) \leq C_2 \quad \text{and} \quad d_{\text{sym}}(Z_2, \text{Pr}_\gamma(Y)) \leq C_2.$$

**7.1. Progressing and contracting geodesics.** This subsection contains the technical results we need from Dahmani and Horbez [DH17]. First, the space  $\mathcal{UE}$  of *uniquely ergonomic* trees is the subspace of  $\partial\text{CV}$  consisting of trees  $T$  with  $\pi^{-1}(\pi(T)) = T$ .

We begin with a definition from [DH17] (Definition 5.13). While it involves (*full*) *recurrence*, the actual definition of full recurrence will not be necessary for our use. Please notice that we have slightly reformulated the [DH17] definition using Lemma 6.2.

**Definition 7.2** ( $(C: A_0, B_0, C_0)$ -progressing). Suppose that  $(T, T') \in \mathcal{UE} \times \mathcal{UE} \setminus \Delta$ , that  $\gamma: \mathbb{R} \rightarrow \text{CV}$  is a greedy fold line from  $T$  to  $T'$ , and that  $S$  is a point on  $\gamma$ . Suppose further that  $C, A_0, B_0, C_0 > 0$ . Then  $\gamma$  is said to be  $(C: A_0, B_0, C_0)$ -*progressing at  $S$*  if the pair  $(\gamma, S)$  satisfies:

If we have

- a tree  $\tilde{S} \in \gamma$  to the right of  $S$  along  $\gamma$  with  $d_{\mathcal{FF}}(S, \tilde{S}) \leq A_0$ , and
- a tree  $R \in \text{CV}$  such that  $\text{Pr}_\gamma(R)$  is to the right of  $S$  and  $d_{\mathcal{FF}}(\tilde{Q}, \text{Pr}_\gamma(R)) \geq B_0$  for any choice of  $\tilde{Q}$  satisfying the conditions of  $\tilde{S}$  above, and
- a greedy folding path  $\gamma': [a, b] \rightarrow \text{CV}$  from  $S'$  to  $R$ , where  $S'$  is either equal to  $\tilde{S}$  or is in the same simplex as  $\tilde{S}$  and is fully recurrent with respect to  $R$ , and
- a value  $c \in [a, b]$  so that, in the symmetrized metric,  $\text{diam}(\gamma'([a, c])) \geq 3$ .

Then  $\text{diam}_{\mathcal{FF}}(\pi \circ \gamma'([a, c])) \geq C_0$ .

Recall that  $Q \geq 0$  is the constant such that all  $d_{\text{CV}}$ -geodesics in  $\text{CV}$  map to unparameterized  $Q$ -quasigeodesics under the map  $\pi: \text{CV} \rightarrow \mathcal{FF}$ . As in the paragraph preceding [DH17, Definition 3.3], we fix  $\kappa \geq 0$  to be as in the conclusion of Proposition 2.1, where  $Q$  is the constant fixed here.

**Definition 7.3** ( $(B, D)$ -contracting). Suppose that  $\gamma$  is a geodesic in  $\text{CV}$  and  $S$  is a point on  $\gamma$ . Then we say that  $\gamma$  is  $(B, D)$ -*contracting at  $S$*  if the following holds:

Let  $a \in \mathbb{R}$  be such that  $\gamma(a) = S$  and let  $b := \inf\{x \in \mathbb{R} \mid \text{diam}_{\mathcal{FF}}(\gamma|_{[a, x]}) \geq B\}$ . Then for all geodesics  $\gamma' \in \text{CV}$ , if  $\pi(\gamma')$  crosses  $\pi(\gamma|_{[a, b]})$  up to distance  $\kappa$ , then there exists an  $a' \in \mathbb{R}$  so that  $d_{\text{CV}}(\gamma'(a'), S) \leq D$ .

We need the following result of Dahmani–Horbez. Informally, it states that progressing implies contracting.

**Proposition 7.4** ([DH17] Proposition 5.17). *There exist constants  $\alpha_0, \beta_0, \gamma_0$  so that for each triple  $(A_0, B_0, C_0)$  with  $A_0 \geq \alpha_0$ ,  $B_0 \geq \beta_0$ , and  $C_0 \geq \gamma_0$ , there exists a  $B > 0$  satisfying:*

*For each  $C > 0$ , there exists a  $D > 0$  such that for all  $(T, T') \in \mathcal{UE} \times \mathcal{UE} \setminus \Delta$ , all greedy folding lines  $\gamma: \mathbb{R} \rightarrow \text{CV}$  from  $T$  to  $T'$ , and all  $S \in \text{im}(\gamma)$ , if  $\gamma$  is  $(C; A_0, B_0, C_0)$ -progressing at  $S$ , then  $\gamma$  is  $(B, D)$ -contracting at  $S$ .*

**7.2. The bounded geodesic image property for fully irreducible axes.** In this section we will prove Theorem 7.8, establishing that a greedy folding axis in CV of a fully irreducible element of  $\text{Out}(F_r)$  has the bounded geodesic image property.

We will need the following lemma, Lemma 7.5. This lemma could be ascertained from a combination of results in [DH17], but we give a more direct proof in our circumstance here.

**Lemma 7.5.** *Suppose  $\gamma$  is a greedy fold axis of a fully irreducible element of  $\text{Out}(F_r)$ . Fix  $A_0, B_0, C_0 \geq 0$  sufficiently large. Then for each  $S \in \gamma$  there is a  $C \geq 0$  such that  $\gamma$  is  $(C; A_0, B_0, C_0)$ -progressing at  $S$ .*

*Proof of Lemma 7.5.* Suppose for the sake of contradiction that we have:

- (1) a sequence of  $\tilde{S}_n \in \gamma$  each lying to the right of  $S$  along  $\gamma$  and satisfying  $d_{\mathcal{FF}}(S, \tilde{S}_n) \leq A_0$ ,
- (2) a sequence  $R_n \in \text{CV}$  such that  $\text{Pr}_\gamma(R_n)$  is to the right of  $\tilde{S}_n$  and  $d_{\mathcal{FF}}(\tilde{S}_n, \text{Pr}_\gamma(R_n)) \geq B_0$ ,
- (3) a sequence of greedy fold paths  $\gamma'_n: [a, b] \rightarrow \text{CV}$  from  $S'_n$  to  $R_n$ , where  $S'_n$  is either equal to  $\tilde{S}_n$  or is in the same simplex as  $\tilde{S}_n$  and is fully recurrent with respect to  $R_n$

so that the greedy fold paths  $\gamma'_n$  further satisfy: if  $\sigma_n$  is the smallest initial segment  $\gamma'_n$  whose endpoints have symmetric distance at least  $n$ , then  $\text{diam}_{\mathcal{FF}}(\sigma_n) \leq C_0$ .

Note that  $\text{diam}_{\text{CV}}(\sigma_n) \rightarrow \infty$  as  $n \rightarrow \infty$ , even though their projections to  $\mathcal{FF}$  remain bounded.

We pass to a subsequence so that the  $\{S'_n\}$  converge in  $\overline{\text{CV}}$  to an  $F_r$ -tree  $S'$ . As in [DH17, Lemma 5.20], we prove:

**Claim 7.6.**  $S' \in \text{CV}$  (as opposed to  $\in \partial\text{CV}$ ).

*Proof of Claim.* Observe that the  $\tilde{S}_n$  vary in a compact subset of  $\gamma$  since  $\gamma$  projects to a quasi-geodesic in  $\mathcal{FF}$ . Hence, we can pass to a subsequence so that they are all in the same simplex – hence so are the  $S'_n$ . Now if  $S' \in \partial\text{CV}$ , then by [BR15, Lemma 7.3] each element of  $F_r$  acting elliptically on  $S'$  also acts elliptically on  $R$ . Since the trees  $S'_n$  live in the same open simplex in CV, this means that  $S'$  has a primitive elliptic element  $a$ , where  $a$  comes from a closed loop in the underlying graph of the open simplex containing all the  $S'_n$ . (In particular, it is in the subgraph whose edge-lengths shrink to zero as  $n \rightarrow \infty$ .) Then  $\langle a \rangle$  is a common reducing factor for the trees  $R$  and  $S'$  and, by the definition of the map  $\pi$ , we have that, coarsely speaking,  $\pi(R) = \pi(S) = \langle a \rangle$ . The choice of  $a$  also implies that  $\pi(S'_n) = \langle a \rangle$ .

Let  $M$  be the constant provided by Lemma 6.3. Define  $C$  to be the constant replacing the  $2\delta$  in Equation 8 if  $[x, y]$  were instead a quasi-geodesic. And let  $\mathfrak{c}$  be the constant of Lemma 6.2. Assume now that  $B_0 \geq 10M + C + \mathfrak{c}$ . We reach a contradiction by applying Lemma 6.3 as follows.

Since all  $S'_n$  live in the same open simplex, we have  $\pi(S'_n) = \pi(S'_1)$  for all  $n \geq 1$ . So we insert into Lemma 6.3:  $y_0 = S'_1 \in \text{CV}$ , and  $R_n = T_n$ , and  $y_1 = \gamma(t_0)$  for some sufficiently large  $t_0$ . Namely, we choose  $t_0$  large enough so that  $\pi(\gamma(t_0))$  is further than  $10M$  to the right of  $\pi(S'_1)$  along  $\pi(\gamma)$ .

Recall that, by assumption, in the factor graph  $\mathcal{FF}$ , we have that  $R_n$  projects farther than  $B_0$  to the right of  $S'_1$  along  $\pi(\gamma)$ . By Equation 8 and Lemma 6.2 this implies that

$$(\pi(R_n) \mid \pi(\gamma(t_0)))_{\pi(S'_1)} \geq B_0 - C - \mathfrak{c}.$$

Lemma 6.3 now implies that

$$(\pi(R) \mid \pi \circ \gamma(t_0))_{\pi(S'_1)} \geq B_0 - M - C - \mathfrak{c} \geq 9M.$$

This contradicts the fact that  $\pi(R) = \pi(S) = \pi(S'_1) = \langle a \rangle$ .

Hence,  $S' \in \text{CV}$ , as desired.  $\square$

Notice that, for sufficiently large  $n$ , we have that  $\widetilde{S}_n$ ,  $S'_n$ , and  $S'$  are all in the same simplex, so  $\pi(\widetilde{S}_n) = \pi(S'_n) = \pi(S')$ . Thus, Item 1 at the start of the proof, together with [BF14, Lemma 3.1], implies  $d_{\mathcal{FF}}(S, S') \leq A_0 + 4$ .

After passing to a subsequence, we see that  $R_n \rightarrow T$  in  $\overline{\text{CV}}$  for some  $T \in \partial\text{CV}$ . Applying [BR15, Lemma 7.3], we have that after passing to a further subsequence certain initial segments of the  $\gamma'_n$  converge uniformly on compact sets to a folding ray  $\gamma'$ . The folding ray  $\gamma'$  has a limit  $S \in \partial\text{CV}$  such that (1) if  $T$  is arational, then  $\pi(S) = \pi(T) \in \partial\mathcal{FF}$  and (2) if  $T$  is not arational, then  $\pi(T)$  and  $\pi(S)$  are uniformly close in  $\mathcal{FF}$ , since  $T$  and  $S$  share a reducing factor. In particular,  $d_{\mathcal{FF}}(\pi(T), \pi(S)) \leq \mathfrak{d}$ , for  $\mathfrak{d}$  as defined in Equation 7.

**Claim 7.7.** *There exists a point  $z$  on  $\gamma'$  such that  $d_{\mathcal{FF}}(S, \text{Pr}_\gamma(z)) \geq C_0 + 10L_{\text{CV}}$ , where  $L_{\text{CV}}$  is the Lipschitz constant for  $\pi: \text{CV} \rightarrow \mathcal{FF}$ .*

*Proof of Claim.* We start by assuming that  $B_0 > C_0 + 10K + M + 3C + 2\mathfrak{c} + \mathfrak{d}$ , where still  $M$  is the constant provided by Lemma 6.3, and  $\delta$  is the hyperbolicity constant of  $\mathcal{FF}$ , and  $C$  adjusts  $2\delta$  for our using quasi-geodesics,  $\mathfrak{c}$  is the constant of Lemma 6.2, and the definition of  $\mathfrak{d}$  is as in Equation 7. Let  $y_0 = \pi(S) = \pi(S'_1) = \pi(S'_n)$ .

By Item 2 at the start of the proof we know  $d_{\mathcal{FF}}(\widetilde{S}_n, \text{Pr}_\gamma(R_n)) \geq B_0$  and thus, by Lemma 6.2,

$$(\pi(\widetilde{S}_n) \mid \mathbf{n}_{\pi \circ \gamma}(\pi(R_n)))_{y_0} \geq B_0 - \mathfrak{c} - C.$$

Therefore, Lemma 6.3 implies that, for  $t_0$  sufficiently large,

$$(9) \quad (\pi(T) \mid \pi(\gamma(t_0)))_{y_0} \geq B_0 - M - C - \mathfrak{c}.$$

We consider separately two cases:

- (1)  $T$  is arational and (2)  $T$  is not arational.

Suppose first that we are in Case 1, i.e. that  $T$  is arational, so that  $\pi(T) = \pi(S)$ . Then, by [BR15], we know that  $\pi(\gamma'(t))$  converges to the point  $\pi(T)$  in  $\partial\mathcal{FF}$  as  $t \rightarrow +\infty$ . Since for ideal triangles in  $\delta$ -hyperbolic spaces, the Gromov product converges in a lim sup sense, we have that, for any point  $\pi(z)$  on  $\pi(\gamma')$  sufficiently far away from the point  $y_0$ :

$$(\pi(z) \mid \pi(\gamma(t_0)))_{y_0} \geq B_0 - M - 2C - \mathfrak{c}.$$

Equation 8 then says

$$d_{\mathcal{FF}}(y_0, \mathbf{n}_{\pi \circ \gamma}(\pi(z))) \geq B_0 - M - 3C - \mathfrak{c}.$$

This in turn implies that

$$d_{\mathcal{FF}}(y_0, \text{Pr}_\gamma(z)) \geq B_0 - M - 3C - 2\mathfrak{c} \geq C_0 + 10L_{\text{CV}}.$$

Thus the claim is satisfied using any such  $z$ .

We now suppose that we are in Case 2, i.e. that  $T$  is not arational, so that  $d_{\mathcal{FF}}(\pi(T), \pi(S)) \leq \mathfrak{d}$ . By the second bullet point of Definition 6.1, we have for all  $t$  sufficiently large:

$$d_{\mathcal{FF}}(\pi(S), \pi(\gamma'(t))) \leq \eta,$$

for some uniform constant  $\eta > 0$  independent of  $\gamma, \gamma'$ , or  $T$  (see Definition 6.1). Hence, for large  $t$

$$d_{\mathcal{FF}}(\pi(T), \pi(\gamma'(t))) \leq \eta + \mathfrak{d}.$$

By taking  $t$  large enough and setting  $z = \gamma'(t)$ , by Equation 9, we have

$$(\pi(z) \mid \pi(\gamma(t_0)))_{y_0} \geq B_0 - M - 2C - \mathfrak{c} - \mathfrak{d}.$$

Using a similar argument as in Case 1, we now also have that the claim is satisfied in this case with our  $z = \gamma'(t)$ , where  $t$  is sufficiently large.  $\square$

We now complete the proof as in [DH17, Lemma 5.21]. Recall that certain initial subsegments of the  $\gamma'_n$  converge uniformly on compact sets to  $\gamma$ . So we can choose a sequence  $\{z_n\}$  in  $CV$  converging to the point  $z$  of Claim 7.7 and satisfying that  $z_n \in \gamma'_n$  for each  $n$ . Now take  $n$  sufficiently large so that both  $d_{\text{sym}}(S'_n, S') < \frac{1}{2}$  and  $d_{\text{sym}}(z_n, z) < \frac{1}{2}$ . Claim 7.7 implies that  $d_{\mathcal{FF}}(S, \text{Pr}_\gamma(z_n)) \geq C_0 + 3L_{CV}$ . Observe that, since  $d_{\text{sym}}(z_n, z) < \frac{1}{2}$  and  $S'$  is fixed,  $d_{\text{sym}}(z_n, S')$  is bounded also, and hence  $d_{\text{sym}}(z_n, S'_n)$  is bounded. This implies that eventually the  $z_n$  lie on the segments  $\sigma_n$ . However,  $\text{diam}_{\mathcal{FF}}(\sigma_n) \leq C_0$  for all  $n$  and  $y_0 \subset \pi(\sigma_n)$  for all  $n$ , so that  $d_{\mathcal{FF}}(S, \text{Pr}_\gamma(z_n)) \leq C_0$ . This contradicts that  $d_{\mathcal{FF}}(S, \text{Pr}_\gamma(z_n)) \geq C_0 + 3L_{CV}$ .  $\square$

**Theorem 7.8.** *Let  $\gamma$  be a greedy folding axis in  $CV$  of a fully irreducible outer automorphism  $\varphi \in \text{Out}(F_r)$ . Then  $\gamma$  has the bounded geodesic image property.*

*Proof.* By combining Lemma 7.5 with Proposition 7.4, we see that for each  $S$  in the image of  $\gamma$  there exist constants  $B_1, D_1 \geq 0$  so that  $\gamma$  will be  $(B_1, D_1)$ -contracting at  $S$ . It follows that  $\gamma$  is  $(B_1, D_1)$ -contracting at each translate  $\varphi^i(S)$ . If necessary, replace  $\varphi$  with a positive power so that  $d_{\mathcal{FF}}(S, \varphi(S)) \geq B_1$ .

First, we refer the reader back to §6 for properties of  $\mathcal{FF}$  and add the following: In a  $\delta$ -hyperbolic metric space  $X$ , every quasigeodesic has the bounded geodesic image property. More precisely, for any  $\delta \geq 0$  and  $Q \geq 1$  there exist constants  $E, E_1 > 0$  with the following property. For any  $x, y$  in a  $\delta$ -hyperbolic metric space  $X$  and a  $Q$ -quasigeodesic  $\alpha$  in  $X$ , if  $\mathbf{n}_\alpha(y)$  occurs at least distance  $E$  to the right of  $\mathbf{n}_\alpha(x)$  along  $\alpha$ , then any  $Q$ -quasigeodesic  $[x, y]$  contains a subsegment  $[x', y']$  such that  $d(x', \mathbf{n}_\alpha(x)) \leq E_1$ , and  $d(y', \mathbf{n}_\alpha(y)) \leq E_1$ , and  $[x', y']$  has Hausdorff distance at most  $E_1$  from  $[\mathbf{n}_\alpha(x), \mathbf{n}_\alpha(y)]$ .

We set  $l = \max\{d_{\text{sym}}(S, \varphi(S)), d_{\mathcal{FF}}(S, \varphi(S))\}$  and recall that  $\mathbf{c}$  is the constant of Lemma 6.2. Note that  $l \geq B_1$ .

Now define  $C_1 = K(10Kl + 2\mathbf{c} + E + K)$  and suppose that we have  $X, Y \in CV$  such that  $d_{\text{sym}}(\text{Pr}_\gamma(X), \text{Pr}_\gamma(Y)) \geq C_1$  with the projection of  $Y$  to  $\gamma$  occurring to the right of the projection of  $X$  along the orientation of  $\gamma$ . Let  $[X, Y]$  be any geodesic from  $X$  to  $Y$  in  $CV$ .

We know  $\pi(\gamma)$  is a  $K$ -parameterized quasigeodesic in  $\mathcal{FF}$ , and  $\pi(\text{Pr}_\gamma(X))$  and  $\pi(\text{Pr}_\gamma(Y))$  are within  $\mathbf{c}$  of the nearest point projections of  $\pi(X)$  and  $\pi(Y)$  to  $\pi(\gamma)$ . Thus, by the choice of  $C_1$ ,

$$d_{\mathcal{FF}}(\pi(\text{Pr}_\gamma(X)), \pi(\text{Pr}_\gamma(Y))) \geq 10Kl + E + 2\mathbf{c}$$

and so

$$d_{\mathcal{FF}}(\mathbf{n}_{\pi(\gamma)}(\pi(X)), \mathbf{n}_{\pi(\gamma)}(\pi(Y))) \geq 10Kl + E.$$

By the bounded geodesic image property for  $\pi(\gamma)$ , there exists a subsegment  $[x', y']$  of the (unparameterized)  $Q$ -quasigeodesic  $\pi([X, Y])$  in  $\mathcal{FF}$  such that

$$d(x', \mathbf{n}_{\pi(\gamma)}(\pi(X))) \leq E_1 \text{ and } d(y', \mathbf{n}_{\pi(\gamma)}(\pi(Y))) \leq E_1.$$

We take  $\pi(S_x)$  to be the translate  $\varphi^i(\pi(S))$  occurring immediately to the right of  $\mathbf{n}_{\pi(\gamma)}(\pi(X))$  along  $\pi(\gamma)$ . Note that  $d_{\mathcal{FF}}(\pi(S_x), \mathbf{n}_{\pi(\gamma)}(\pi(X))) \leq lK$ . Similarly, we let  $\pi(S_y)$  be the translate  $\varphi^j(\pi(S))$  which is the *second to last* translate of  $\pi(S)$  before  $\mathbf{n}_{\pi(\gamma)}(\pi(Y))$  along  $\pi(\gamma)$ . Then  $d_{\mathcal{FF}}(\pi(S_y), \mathbf{n}_{\pi(\gamma)}(\pi(Y))) \leq 2lK$ . Our selection of  $S_x$  and  $S_y$  guarantees that we may apply the bounded geodesic image property at both of these points.

We first apply the  $(B_1, D_1)$ -contraction property for  $S_x$ , using the segment of  $[x', y']$  that crosses a subsegment of  $\pi(\gamma)$  starting at  $\pi(S_x)$  with length  $B_1$ , and find a point  $Z_1$  on the corresponding segment  $J_1$  of  $[X, Y]$  such that  $d_{\text{sym}}(Z_1, S_x) \leq D_1$ . Then we apply the  $(B_1, D_1)$ -contraction property for  $S_y$ , using the segment of  $[x', y']$  that crosses a subsegment of  $\pi(\gamma)$  starting at  $\pi(S_y)$  with length

$B_1$ , and find a point  $Z_2$  on the corresponding segment  $J_2$  of  $[X, Y]$  such that  $d_{\text{sym}}(Z_2, S_y) \leq D_1$ . By construction,  $J_1$  occurs before  $J_2$  in  $[X, Y]$  and therefore  $Z_1$  comes before  $Z_2$  along  $[X, Y]$ . Since  $\gamma$  is thick, there is a  $C_2$  such that  $d_{\text{sym}}(Z_1, S_x)$  and  $d_{\text{sym}}(Z_2, S_y)$  are both less than  $C_2$ .  $\square$

**Corollary 7.9.** *Let  $\gamma$  be a greedy folding axis in CV of some fully irreducible  $\varphi \in \text{Out}(F_r)$ . Then there exist constants  $\rho \geq 0$  and  $c_0 \geq 0$  with the following property. Let  $C_1, C_2 > 0$  be the bounded geodesic image property constants for  $\gamma$  (where we know that this property holds for  $\gamma$  by Theorem 7.8). Let  $R \geq 1$  be an arbitrary sufficiently large number.*

*Let  $X, Y \in \text{CV}$  be such that  $d_{\text{sym}}(\text{Pr}_\gamma(X), \text{Pr}_\gamma(Y)) \geq C_1$  and such that  $\text{Pr}_\gamma(X) = \gamma(t_1)$ ,  $\text{Pr}_\gamma(Y) = \gamma(t_2)$  where  $t_2 - t_1 \geq R$ . Then any geodesic segment  $[X, Y]$  contains a subsegment of length  $\geq R - c_0$  that  $\rho$ -fellow travels a subsegment of the same length in  $\gamma$ .*

To prove the corollary, we need the following lemma.

**Lemma 7.10.** *Let  $K \geq 0$  and  $C \geq 0$ . Then there exists a constant  $A' = A'(K, C) > 0$  with the following property. Let  $\gamma$  be a geodesic in CV such that  $\pi \circ \gamma$  is a  $K$ -quasigeodesic in  $\mathcal{FF}$ . Let  $P, Q \in \gamma$  and  $Z_1, Z_2 \in \text{CV}$  be such that  $d_{\text{sym}}(Z_1, P) \leq C$  and  $d_{\text{sym}}(Z_2, P) \leq C$ . Let  $[P, Q]$  denote the segment of  $\gamma$  from  $P$  to  $Q$ . Then for any geodesic segment  $[Z_1, Z_2]$  in CV, we have that the segments  $[P, Q]$  and  $[Z_1, Z_2]$  are  $A'$ -Hausdorff close with respect to  $d_{\text{sym}}$ .*

*Proof.* Consider the geodesic segments  $[P, Z_1]$ ,  $[Z_2, Q]$  and the path  $\alpha = [P, Z_1] \cup [Z_1, Z_2] \cup [Z_2, Q]$ . Since  $d_{\text{sym}}(Z_1, P) \leq C$  and  $d_{\text{sym}}(Z_2, P) \leq C$ , the path  $\alpha$  is a  $C'$ -quasigeodesic in CV for some constant  $C' \geq 0$  depending only on  $C$ . Therefore, by [DT17, Theorem 4.1], for some constant  $A = A(C', K) \geq 0$ , the paths  $\alpha$  and  $[P, Q]$  are  $A$ -Hausdorff close with respect to  $d_{\text{sym}}$ . It follows that  $[P, Q]$  and  $[Z_1, Z_2]$  are  $(A + C)$ -close, and the lemma holds with  $A' = A + C$ .  $\square$

We can now prove Corollary 7.9.

*Proof of Corollary 7.9.* By the bounded geodesic image property for  $\gamma$ , there exists a subsegment  $[Z_1, Z_2]$  in  $[X, Y]$  such that  $d_{\text{sym}}(\text{Pr}_\gamma(X), Z_1) \leq C_2$  and  $d_{\text{sym}}(\text{Pr}_\gamma(Y), Z_2) \leq C_2$ . Since  $\gamma$  is the axis of a fully irreducible,  $\pi \circ \gamma$  is a  $K$ -quasigeodesic in  $\mathcal{FF}$  for some  $K \geq 0$ . Thus, by Lemma 7.10, the segments  $[Z_1, Z_2]$  and  $[\text{Pr}_\gamma(X), \text{Pr}_\gamma(Y)] = [\gamma(t_1), \gamma(t_2)]$  are  $A'$ -Hausdorff close with respect to  $d_{\text{sym}}$ .

By the triangle inequality we have  $d_{\text{CV}}(Z_1, Z_2) \geq R - 2C_2$ . Then by Lemma 5.8 the segment  $[Z_1, Z_2]$  has a subsegment of length  $\geq R - 2C_2 - cC_2$  that  $cC_2$  fellow travels a segment of  $\gamma$  of the same length (where  $c$  is the constant depending on thickness of  $\gamma$  provided by Lemma 5.8). Therefore the conclusion of the corollary holds with  $\rho = cC_2$  and  $c_0 = 2C_2 + cC_2$ .  $\square$

## 8. RANDOM OUTER AUTOMORPHISMS ARE TRIANGULAR

Let  $\varphi$  be a principal fully irreducible outer automorphism (which exists by Proposition 5.11) and let  $\gamma$  be its lone folding axis (see §4.2). Evidently,  $\gamma$  is both a greedy and a simple folding path.

By Theorem 7.8, there are constants  $C_1, C_2 \geq 0$  such that  $\gamma$  is  $(C_1, C_2)$ -contracting. Let  $\rho \geq 0$  be the constant given by Corollary 7.9. With this set-up we have the following statement:

**Proposition 8.1.** *There exists a constant  $R_1 \geq 0$  satisfying the following: Let  $\psi$  be a fully irreducible outer automorphism with simple folding axis  $\gamma'$ . Suppose that the  $\text{Pr}_\gamma$ -projection of  $\gamma'$  to  $\gamma$  has diameter  $\geq R_1$  and that, using the orientations on these paths, the left end of  $\gamma'$  projects to the left of the right end of  $\gamma'$ . Then  $\psi$  is triangular.*

*Proof.* The conclusion follows directly from Corollary 5.14 and Corollary 7.9.  $\square$

Proposition 8.1 is a key technical result that combines the full strength of the bounded geodesic image property (Corollary 7.9) and of the “stability” result for triangular and principal outer automorphisms (Corollary 5.14).



We now turn to the proof of our main results stated in the introduction. To do so, we freely use the constants collected in §6. To simplify notation, we denote the  $\pi$ -image of a point  $x$  in CV by  $\bar{x} \subset \mathcal{FF}$ .

**Theorem A.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ . Let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose  $\langle \text{supp}(\mu) \rangle_+$  contains  $\varphi^{-1}$  for some principal fully irreducible  $\varphi \in \text{Out}(F_r)$ . Then  $w_n$  is an ageometric triangular fully irreducible outer automorphism with probability tending to 1 as  $n \rightarrow \infty$ .*

*Proof.* We work with the constants established at the beginning of this section. As before, we apply Proposition 2.2 to the action  $\text{Out } F_r \curvearrowright \mathcal{FF}$  with  $Q$  equal to the (unparameterized) quasigeodesic constants for the image of a geodesic in CV. We set  $L = L_{\text{CV}} \cdot (R_1 + 2\kappa + 2\mathfrak{c} + 1)$ , for  $R_1$  as in Proposition 8.1 and  $\mathfrak{c}$  as in Proposition 6.2.

Let  $\gamma(t)$  be the (lone) axis in CV of our principal fully irreducible  $\varphi$ . By Remark 3.1, for the left action of  $\text{Out}(F_r)$  on CV, iteration of  $\varphi^{-1}$  on  $\gamma$  translates in the direction of  $t \rightarrow \infty$  along  $\gamma$ . Then  $\bar{\gamma}(t)$  is a directed quasigeodesic axis for  $\varphi^{-1}$  in  $\mathcal{FF}$  (again with respect to the left action of  $\text{Out}(F_r)$ ). Thus we can apply Proposition 2.2 to  $\bar{\gamma}$  since, by assumption,  $\varphi^{-1} \in \langle \text{supp}(\mu) \rangle_+$ .

Letting  $\gamma_{w_n}$  denote a simple folding axis for  $w_n$  (as  $w_n$  is fully irreducible with probability tending to 1 as  $n \rightarrow \infty$ ), it follows by Proposition 2.2 that  $\bar{\gamma}_{w_n}$  has an  $(L, \kappa)$ -oriented match with  $\bar{\gamma}$ . By the hyperbolic geometry of  $\mathcal{FF}$ , this means that there is a  $g_n \in \text{Out}(F_r)$  such that the nearest point projection in  $\mathcal{FF}$  of  $g_n \cdot \bar{\gamma}_{w_n}$  to  $\bar{\gamma}$  has diameter at least  $L_{\text{CV}} \cdot (R_1 + 2\mathfrak{c} + 1)$ . If we instead use the projection  $\pi \circ \text{Pr}_\gamma$ , we find that the diameter of the image of the projection is at least  $L_{\text{CV}} \cdot (R_1 + 1)$  (see Proposition 6.2). Further, since the match is oriented, we have that the left end of  $\bar{\gamma}_{w_n}$  projects to the left of its right end, along  $\gamma$ . Finally, we use the fact that  $\pi: \text{CV} \rightarrow \mathcal{FF}$  is  $L_{\text{CV}}$ -Lipschitz to conclude that the  $\text{Pr}_\gamma$ -projection of  $\gamma_{w_n}$  to  $\gamma$  has diameter at least  $R_1$ . Hence, from Proposition 8.1, we conclude that  $w_n$  is triangular.  $\square$

In the following corollary  $i(\varphi_n)$  is the rotationless index of  $\varphi_n$  (see §4.6) and  $\text{ind}_{\text{geom}}(T_+^{\varphi_n})$  is the geometric index of  $\varphi_n$  (see [CH12]).

**Corollary B.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ , and let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose that  $\langle \text{supp}(\mu) \rangle_+$  contains  $\varphi^{-1}$  for some principal fully irreducible  $\varphi \in \text{Out}(F_r)$ .*

*Then, with probability going to 1 as  $n \rightarrow \infty$ ,  $w_n$  is an ageometric fully irreducible outer automorphism  $\varphi_n$  satisfying:*

- (1) *each component of the ideal Whitehead graph  $IW(\varphi)$  is a triangle;*
- (2) *each branchpoint of  $T_+^{\varphi_n}$  is trivalent;*
- (3)  *$i(\varphi_n) > 1 - r$  and  $\text{ind}_{\text{geom}}(T_+^{\varphi_n}) < 2r - 2$ ;*
- (4) *the tree  $T_+^{\varphi_n}$  is nongeometric.*

*Proof.* Theorem A implies that, with probability going to 1 as  $n \rightarrow \infty$ , the outer automorphism  $\varphi_n$  is triangular fully irreducible. Thus, in particular,  $\varphi_n$  is ageometric, so that the tree  $T_+^{\varphi_n}$  is nongeometric and  $i(\varphi_n) > 1 - r$ . Also, for nongeometric trees we have  $\text{ind}_{\text{geom}}(T_+^{\varphi_n}) < 2r - 2$  (see [CH12, p. 300] for more details). Proposition 5.12 now implies that  $T_+^{\varphi_n}$  is trivalent.  $\square$

**Corollary C.** *Let  $r \geq 3$  and let  $\mu$  be a probability distribution on  $\text{Out}(F_r)$  which is nonelementary and has bounded support for the action on  $\mathcal{FF}$ , and let  $(w_n)$  be the random walk determined by  $\mu$ . Suppose that  $\langle \text{supp}(\mu) \rangle_+$  contains a pair of elements  $\varphi$  and  $\psi$ , such that both  $\varphi$  and  $\psi^{-1}$  are principal fully irreducible outer automorphisms.*

*Then, with probability approaching 1 as  $n \rightarrow \infty$ , both  $w_n$  and  $w_n^{-1}$  are ageometric triangular fully irreducible outer automorphisms. In particular, both fixed trees of  $w_n$  are trivalent.*

*Proof.* Let  $\mathcal{T} \subset \text{Out}(F_r)$  be the subset consisting of the triangular fully irreducible elements. By Theorem A,  $\mu^n(\mathcal{T}) \rightarrow 1$ . Recall that  $\check{\mu}$  denotes the reflected measure,  $\check{\mu}(g) = \mu(g^{-1})$ , so the same theorem implies that  $\check{\mu}^n(\mathcal{T}) \rightarrow 1$ . But the distribution of the random element  $w_n^{-1}$  is  $\check{\mu}^n$ , and Theorem A applies to  $\check{\mu}$  as well. Therefore the probability that both  $w_n$  and  $w_n^{-1}$  lie in  $\mathcal{T}$  tends to 1 as  $n \rightarrow \infty$ .  $\square$

## REFERENCES

- [ABH<sup>+</sup>06] P. Arnoux, V. Berthé, A. Hilion, A. Siegel, et al. Fractal representation of the attractive lamination of an automorphism of the free group. In *Annales de l'institut Fourier*, volume 56, pages 2161–2212. Chartres: L'Institut, 1950-, 2006.
- [ACT15] G. N. Arzhantseva, C. H. Cashen, and J. Tao. Growth tight actions. *Pacific J. Math*, 278(1):1–49, 2015.
- [AK11] Y. Algom-Kfir. Strongly contracting geodesics in outer space. *Geom. Topol*, 15(4):2181–2233, 2011.
- [AKB12] Y. Algom-Kfir and M. Bestvina. Asymmetry of outer space. *Geom. Dedicata*, 156:81–92, 2012.
- [AKKP18] Y. Algom-Kfir, I. Kapovich, and C. Pfaff. Stable Strata of Geodesics in Outer Space. *International Mathematics Research Notices*, 2018(00):pp. 1–30, 2018.
- [AKP17] Y. Algom-Kfir and C. Pfaff. Normalizers and centralizers of cyclic subgroups generated by lone axis fully irreducible outer automorphisms. *New York J. Math*, 23:365–381, 2017.
- [Bes11] M. Bestvina. A Bers-like proof of the existence of train tracks for free group automorphisms. *Fund. Math*, 214:89–100, 2011.
- [Bes14] M. Bestvina. Geometry of outer space. In *Geometric group theory*, volume 21 of *IAS/Park City Math. Ser.*, pages 173–206. Amer. Math. Soc., Providence, RI, 2014.
- [BF94] M. Bestvina and M. Feighn. Outer limits. *preprint*, pages 1–19, 1994.
- [BF14] M. Bestvina and M. Feighn. Hyperbolicity of the complex of free factors. *Advances in Mathematics*, 256:104–155, 2014.
- [BFH97] M. Bestvina, M. Feighn, and M. Handel. Laminations, trees, and irreducible automorphisms of free groups. *Geometric and Functional Analysis*, 7(2):215–244, 1997.
- [BH92] M. Bestvina and M. Handel. Train tracks and automorphisms of free groups. *The Annals of Mathematics*, 135(1):1–51, 1992.
- [Bog08] O. Bogopolski. *Introduction to group theory*. EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008. Translated, revised and expanded from the 2002 Russian original.
- [BR15] M. Bestvina and P. Reynolds. The boundary of the complex of free factors. *Duke Mathematical Journal*, 164(11):2213–2251, 2015.
- [CH12] T. Coulbois and A. Hilion. Botany of irreducible automorphisms of free groups. *Pacific Journal of Mathematics*, 256(2), 2012.
- [CHL08] T. Coulbois, A. Hilion, and M. Lustig. R-trees and laminations for free groups I: algebraic laminations. *Journal of the London Mathematical Society*, 78(3):723–736, 2008.
- [CL95] M. Cohen and M. Lustig. Very small group actions on r-trees and dehn twist automorphisms. *Topology*, 34(3):575–617, 1995.
- [CM15] D. Calegari and J. Maher. Statistics and compression of scl. *Ergodic Theory and Dynamical Systems*, 35(1):64–110, 2015.
- [Cou14] T. Coulbois. Free group automorphisms and train-track representative in python/sage. <https://github.com/coulbois/sage-train-track>, 2012–2014.
- [CV86] M. Culler and K. Vogtmann. Moduli of graphs and automorphisms of free groups. *Inventiones mathematicae*, 84(1):91–119, 1986.
- [DH17] F. Dahmani and C. Horbez. Spectral Theorems for Random Walks on Mapping Class Groups and  $\text{Out}(F_n)$ . *International Mathematics Research Notices*, page rnnw306, 2017.
- [DKL15] S. Dowdall, I. Kapovich, and C. J. Leininger. Dynamics on free-by-cyclic groups. *Geom. Topol.*, 19(5):2801–2899, 2015.
- [DT17] S. Dowdall and S. J. Taylor. Hyperbolic extensions of free groups. *Geom. Topol.*, 22(1):517–570, 2017.
- [FH11] M. Feighn and M. Handel. The recognition theorem for  $\text{Out}(F_n)$ . *Groups Geom. Dyn.*, 5(1):39–106, 2011.
- [FM11] S. Francaviglia and A. Martino. Metric properties of outer space. *Publicacions Matemàtiques*, 55(2):433–473, 2011.
- [GJLL98] D. Gaboriau, A. Jaeger, G. Levitt, and M. Lustig. An index for counting fixed points of automorphisms of free groups. *Duke Math. J.*, 93(3):425–452, 1998.
- [GL95] D. Gaboriau and G. Levitt. The rank of actions on R-trees. In *Annales scientifiques de l'Ecole normale supérieure*, volume 28, pages 549–570. Société mathématique de France, 1995.

- [GM17] V. Gadre and J. Maher. The stratum of random mapping classes. *Ergodic Theory and Dynamical Systems*, pages 1–17, 2017.
- [Ham12] U. Hamenstädt. The boundary of the free factor graph. *arXiv preprint arXiv:1211.1630*, 2012.
- [HM07] M. Handel and L. Mosher. Parageometric outer automorphisms of free groups. *Transactions of the American Mathematical Society*, 359(7):3153–3184, 2007.
- [HM11] M. Handel and L. Mosher. *Axes in outer space*. Number 1004. Amer Mathematical Society, 2011.
- [HM13] M. Handel and L. Mosher. The free splitting complex of a free group, I: hyperbolicity. *Geom. Topol.*, 17(3):1581–1672, 2013.
- [Hor14] C. Horbez. The horoboundary of outer space, and growth under random automorphisms. *arXiv preprint arXiv:1407.3608*, 2014.
- [KL10] I. Kapovich and M. Lustig. Intersection form, laminations and currents on free groups. *Geometric And Functional Analysis*, 19(5):1426–1467, 2010.
- [KM98] V. A. Kaimanovich and H. Masur. The Poisson boundary of Teichmüller space. *J. Funct. Anal.*, 156(2):301–332, 1998.
- [KP15] I. Kapovich and C. Pfaff. A train track directed random walk on  $\text{Out}(F_r)$ . *International Journal of Algebra and Computation*, 25(5):745–798, August 2015.
- [KR14] I. Kapovich and K. Rafi. On hyperbolicity of free splitting and free factor complexes. *Groups Geom. Dyn.*, 8(2):391–414, 2014.
- [LL03] G. Levitt and M. Lustig. Irreducible automorphisms of  $F_n$  have north–south dynamics on compactified outer space. *Journal of the Institute of Mathematics of Jussieu*, 2(01):59–72, 2003.
- [LP97] G. Levitt and F. Paulin. Geometric group actions on trees. *American Journal of Mathematics*, pages 83–102, 1997.
- [Mah11] J. Maher. Random walks on the mapping class group. *Duke Mathematical Journal*, 156(3):429–468, 2011.
- [Mah12] J. Maher. Exponential decay in the mapping class group. *J. Lond. Math. Soc. (2)*, 86(2):366–386, 2012.
- [MP16] L. Mosher and C. Pfaff. Lone axes in outer space. *Algebr. Geom. Topol.*, 16(6):3385–3418, 2016.
- [MS17] J. Maher and A. Sisto. Random subgroups of acylindrically hyperbolic groups and hyperbolic embeddings. *International Mathematics Research Notices*, 2017.
- [MT14] J. Maher and G. Tiozzo. Random walks on weakly hyperbolic groups. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2014.
- [Pau89] F. Paulin. The Gromov topology on  $\mathbf{R}$ -trees. *Topology Appl.*, 32(3):197–221, 1989.
- [Pfa12] C. Pfaff. *Constructing and Classifying Fully Irreducible Outer Automorphisms of Free Groups*. PhD thesis, Rutgers University, 2012. PhD Thesis, <http://www.math.rutgers.edu/~cpfaff/Thesis.pdf>.
- [Rey12] P. Reynolds. Reducing systems for very small trees. *arXiv preprint arXiv:1211.3378*, 2012.
- [Riv08] I. Rivin. Walks on groups, counting reducible matrices, polynomials, and surface and free group automorphisms. *Duke Math. J.*, 142(2):353–379, 2008.
- [Sis11] A. Sisto. Contracting elements and random walks. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, to appear, 2011.
- [Sko89] R. Skora. Deformations of length functions in groups, preprint. *Columbia University*, 1989.
- [Sta83] J.R. Stallings. Topology of finite graphs. *Inventiones Mathematicae*, 71(3):551–565, 1983.
- [Vog15] K. Vogtmann. On the geometry of outer space. *Bull. Amer. Math. Soc. (N.S.)*, 52(1):27–46, 2015.

DEPARTMENT OF MATHEMATICS AND STATISTICS, HUNTER COLLEGE OF CUNY, 695 PARK AVE, NEW YORK, NY 10065

[HTTP://MATH.HUNTER.CUNY.EDU/ILYAKAPO/](http://math.hunter.cuny.edu/ilyakapo/),  
 Email address: [ik535@hunter.cuny.edu](mailto:ik535@hunter.cuny.edu)

CUNY COLLEGE OF STATEN ISLAND AND CUNY GRADUATE CENTER, VICTORY BOULEVARD, STATEN ISLAND, NY 10314

[HTTP://WWW.MATH.CSI.CUNY.EDU/~MAHER/](http://www.math.csi.cuny.edu/~maher/),  
 Email address: [joseph.maher@csi.cuny.edu](mailto:joseph.maher@csi.cuny.edu)

DEPARTMENT OF MATH & STATS, QUEEN’S UNIVERSITY, JEFFERY HALL, 48 UNIVERSITY AVE., KINGSTON, ON CANADA, K7L 3N6

[HTTP://MATH.UCSB.EDU/~CPF AFF/](http://math.ucsb.edu/~cpfaff/),  
 Email address: [cpfaff@math.ucsb.edu](mailto:cpfaff@math.ucsb.edu)

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY, 1805 BROAD ST, PHILADELPHIA, PA 19122

[HTTPS://MATH.TEMPLE.EDU/~SAMUEL.TAYLOR/](https://math.temple.edu/~samuel.taylor/),  
 Email address: [samuel.taylor@temple.edu](mailto:samuel.taylor@temple.edu)