

A DENSE GEODESIC RAY IN THE $\text{Out}(F_r)$ -QUOTIENT OF REDUCED OUTER SPACE.

Yael Algom-Kfir, Catherine Pfaff

ABSTRACT. In [Mas81] Masur proved the existence of a dense geodesic in the moduli space for a Teichmüller space. We prove an analogue theorem for reduced Outer Space endowed with the Lipschitz metric. We also prove two results possibly of independent interest: we show Brun’s unordered algorithm weakly converges and from this prove that the set of Perron-Frobenius eigenvectors of positive integer $m \times m$ matrices is dense in the positive cone \mathbb{R}_+^m (these matrices will in fact be the transition matrices of positive automorphisms). We give a proof in the appendix that not every point in the boundary of Outer Space is the limit of a flow line.

1. INTRODUCTION

1.1. Geodesics in Outer Space. One of the richest and most expansive methods for studying surfaces has been through the ergodic geodesic flow on Teichmüller space. As an example, it was used by Eskin and Mirzakhani [EM11] to count pseudo-Anosov conjugacy classes of a bounded length. For this reason, the papers of Masur [Mas82] and Veech [Vee82] independently proving the ergodicity of the Teichmüller flow were seminal in the field. The existence of an $\text{Out}(F_r)$ -invariant ergodic geodesic flow on Outer Space may similarly expand the tools for studying $\text{Out}(F_r)$. Before giving the proof of the ergodicity theorem in Teichmüller space, Masur performed the following a “litmus test” for its plausibility.

Theorem 1.1 ([Mas81]). *Given a closed surface S_g of genus g , there exists a Teichmüller geodesic in the Teichmüller space T_g whose projection $p : T_g \rightarrow T_g/MCG(S_g)$ to moduli space is dense in both directions.*

Our main theorem is an $\text{Out}(F_r)$ analogue of the above theorem. Some of the terms in the theorem are defined directly below its statement.

Theorem A. For each $r \geq 2$, there exists a geodesic fold ray in the reduced Outer Space $\mathcal{R}\mathcal{X}_r$ whose projection to $\mathcal{R}\mathcal{X}_r/\text{Out}(F_r)$ is dense.

This research was supported by THE ISRAEL SCIENCE FOUNDATION (grant no. 1941/14). The second author was supported first by the ARCHIMEDE Labex (ANR-11-LABX- 0033) and the A*MIDEX project (ANR-11-IDEX-0001-02) funded by the “Investissements d’Avenir,” managed by the ANR. She was secondly supported by the CRC701 grant of the DFG, supporting the projects B1 and C13 in Bielefeld.

- Remark 1.2.** (1) The reduced Outer Space $\mathcal{R}\mathcal{X}_r$ is a subcomplex of the Outer Space \mathcal{X}_r , which consists of those graphs without separating edges (see Definition 2.12). It is an equivariant deformation retract of \mathcal{X}_r .
- (2) The metric on $\mathcal{R}\mathcal{X}_r$ with respect to which the ray in Theorem A is a geodesic is the Lipschitz metric (see Definition 2.25). It is an asymmetric metric (analogous to the Thurston metric on Teichmüller space [Thu]) that has proven to be very useful in the $\text{Out}(F_r)$ context, e.g. [Bes11].
- (3) Because of the asymmetry of the metric, our geodesics will always be “directed geodesics,” i.e. maps $\Gamma: [0, \infty) \rightarrow \mathcal{X}_r$ such that $d(\Gamma(t), \Gamma(t')) = t' - t$ for $t' > t \geq 0$, but not necessarily for $t > t'$.
- (4) A fold line is a special kind of geodesic in \mathcal{X}_r (explicitly described in Definition 3.13) that is analogous to a Teichmüller geodesic.
- (5) Comparing between Theorem 1.1 and Theorem A, one may notice that our theorem declares the existence of a ray in contrast with Masur’s theorem which declares the existence of a geodesic. We could easily extend our ray to a bi-infinite geodesic. However, the density of the image of the ray will follow from techniques that we cannot extend to the backwards direction.

Returning to Remark 1.2(1), we note that for proving algebraic properties of $\text{Out}(F_r)$, one may usually replace Outer Space with $\mathcal{R}\mathcal{X}_r$. However, on the geometric side, it is not known whether or not $\mathcal{R}\mathcal{X}_r$ is convex in any coarse sense.

We hence pose two questions:

Question 1.3. *For each $r \geq 2$, does there exist a geodesic fold line in \mathcal{X}_r that is dense in both directions in $\mathcal{X}_r/\text{Out}(F_r)$?*

Question 1.4. *For each $r \geq 2$, is the reduced Outer Space $\mathcal{R}\mathcal{X}_r$ coarsely convex? For example, given points $x, y \in \mathcal{R}\mathcal{X}_r$ does there always exist a geodesic from x to y which is contained in reduced Outer Space?*

1.2. The unit tangent bundle. Masur obtained Theorem 1.1 as a corollary of his study of the geodesic flow on Teichmüller Space. The unit tangent bundle of Teichmüller space is isomorphic to its unit cotangent bundle Q_0 , which may be described explicitly as the space of unit area quadratic differentials on a closed surface S_g of genus g . The geodesic flow on Teichmüller space is a $\text{MCG}(S_g)$ invariant action of \mathbb{R} on Q_0 . For $t \in \mathbb{R}$ we denote the flow by $T_t: Q_0 \rightarrow Q_0$. Given a quadratic differential $q \in Q_0$, the set of points $\{q_t\}_{t \in \mathbb{R}} = \{T_t q\}_{t \in \mathbb{R}}$ defines a geodesic in the Teichmüller space T_g .

Theorem 1.5 ([Mas81]). *For a closed surface S_g of genus $g > 1$, there exists a quadratic differential $q \in Q_0$ so that the projection of $\{q_t\}$ for either $t > 0$ or $t < 0$ is dense in $Q_0/\text{MCG}(S_g)$.*

The analogue of Theorem 1.5 is not obvious, as it is unclear what the unit tangent bundle should be. In Outer Space the analogues of the following facts are false:

- (1) If Γ_1, Γ_2 are two bi-infinite Teichmüller geodesics satisfying that there exist non-degenerate intervals $I, I' \subset \mathbb{R}$ with $\Gamma_1(I) \subset \Gamma_2(I')$ then, up to reparametrization, $\Gamma_1 = \Gamma_2$.
- (2) There exists a compactification $\overline{T_g}$ of T_g such that for any two points $x, y \in \overline{T_g}$ there exists a geodesic from x to y .

The failure of (1) is an impediment to a “local” description of the tangent bundle. This failure is similar to its failure in a simplicial tree. Two geodesics in Outer Space can meet for a period of time and then diverge from each other (or even alternately meet and diverge). On the other hand, the failure of (2) is an impediment to a “global” description in terms of the boundary of Outer Space. One may want to use equivalence classes of rays emanating from a common base-point. However, we show in §9 that there are points on the boundary i.e. $\overline{\mathcal{X}_r} - \mathcal{X}_r$ that are not ends of geodesic fold rays. (While this result may be known to the experts, to our knowledge this is the first time it appears in print). Thus one would first have to identify the visual boundary as a subset of $\overline{\mathcal{X}_r} - \mathcal{X}_r$ and this has yet to be done.

For the purposes of this paper, we propose the following analogue of the unit tangent bundle. Given a point $x \in \mathcal{R}\mathcal{X}_r$ there are finitely many germs $[\alpha]$ of fold lines α in $\mathcal{R}\mathcal{X}_r$ initiating at x . Define

$$\mathcal{UR}\mathcal{X}_r = \{(x, [\alpha]) \mid x \in \mathcal{R}\mathcal{X}_r, \alpha \text{ is a fold line with } \alpha(0) = x \text{ and } \text{Im}(\alpha) \subset \mathcal{R}\mathcal{X}_r\}.$$

Given a geodesic ray $\gamma: [0, \infty) \rightarrow \mathcal{R}\mathcal{X}_r$, for $t \in \mathbb{R}$ we denote by γ_t the path $\gamma_t(s) = \gamma(t+s)$. We may associate to γ a path in the unit tangent bundle $\tilde{\gamma}(t) = (\gamma(t), [\gamma_t])$. We prove:

Theorem B. For each $r \geq 2$, there exists a Lipschitz geodesic fold ray $\tilde{\gamma}: [0, \infty) \rightarrow \mathcal{UR}\mathcal{X}_r$ so that the projection of $\tilde{\gamma}$ to $\mathcal{UR}\mathcal{X}_r/\text{Out}(F_r)$ is dense.

1.3. Geodesics in other subcomplexes. For each $r \geq 2$, we define the *theta subcomplex* \mathcal{T}_r to be the subspace of $\mathcal{R}\mathcal{X}_r$ consisting of all points in \mathcal{X}_r whose underlying graph is either a rose or a theta graph, see Figure 1. This subcomplex carries the significance of being the minimal subcomplex containing the image of the Cayley graph under the natural map. Both as a warm-up, and for its intrinsic significance, we initially prove Theorem A in \mathcal{T}_r .

Theorem C. For each $r \geq 2$, there exists a fold line in \mathcal{T}_r that projects to a Lipschitz geodesic fold ray in $\mathcal{T}_r/\text{Out}(F_r)$.

1.4. Outline. We begin by outlining the proof of Theorem C. After proving Theorem C, we develop the topological machinery necessary to extend the more combinatorial




FIGURE 1. The underlying graphs of the simplices of \mathcal{T}_r . The graph on the left is called a *rose* and that on the right is called a *theta graph*.

arguments of the proof of Theorem C to the proof of Theorem B (and Theorem A as a corollary).

Recall that points in Outer Space are marked metric graphs equivalent up to homotopy, see Definition 2.2. As described in Definition 2.11, $\text{Out}(F_r)$ acts on the right by changing the marking. To a point $x \in \mathcal{T}_r/\text{Out}(F_r)$, one may associate a positive vector, the “length vector” recording the graph’s edge lengths. The folding operation may be translated to a matrix recording the change in edge lengths from its initial point to its terminal point. In this dictionary, a fold ray in $\mathcal{T}_r/\text{Out}(F_r)$ should correspond to an initial vector and a sequence of fold matrices. However, not every such sequence comes from a fold ray: a particular fold may or may not be allowed for a specific vector depending on whether its image is again a positive vector. Our challenge is to construct a sequence of fold matrices $\{T_k\}_{k=1}^\infty$ satisfying that, for some positive vector w_0 , if we write $w_i = T_i \cdots T_1(w_0)$ for each $i \in \mathbb{N}$ then

- I. the fold T_{i+1} is allowed in $w_i = T_i \cdots T_1(w_0)$ for each integer $1 \leq i < \infty$,
- II. the set of vectors w_i is projectively dense in a simplex and,
- III. the corresponding fold ray is a geodesic ray in the Lipschitz metric.

In order to prove item (II) of the list, we prove the following fact:

Theorem D. For each $r \geq 2$, let $S_{l_1}^r$ be the set of unit vectors according to the l_1 metric in \mathbb{R}_+^r . The set of Perron-Frobenius eigenvectors of matrices arising as the transition matrices of a positive automorphisms in $\text{Aut}(F_r)$ is dense in the $S_{l_1}^r$.

The proof of Theorem D, in §4, uses Brun’s algorithm [Bru57]. We also prove in §4 that Brun’s (unordered) algorithm converges in angle, a result to our knowledge previously absent from the literature in dimensions higher than four. (Brun proved it in [Bru57] for dimensions three and four.)

To construct the sequence $\{T_k\}$ of fold matrices we enumerate the powers of “Brun matrices” (see §4): P_1, P_2, P_3, \dots . To each P_i , we can attach the following data:

- a positive Perron-Frobenius eigenvector v_i ,
- a positive automorphism $g_i \in \text{Aut}(F_r)$, also denoted g_{v_i} , so that the transition matrix of g_i is P_i ,
- and a decomposition of P_i into fold matrices, arising from the decomposition of g_i into Nielsen generators which correspond to moves in Brun’s algorithm.

We remark that this method is reminiscent of Masur’s paper, where he proved the existence of a dense geodesic in $R_g = Q_0/\text{MCG}(S_g)$ using the fact that closed loops in R_g are dense. The resemblance stems from the facts that the decomposition in the third item defines a loop in $\mathcal{R}\mathcal{X}_r/\text{Out}(F_r)$ based at v_i and that the density, established by Theorem D, of the set of Perron-Frobenius eigenvectors $\{v_i\}$ ’s.

We concatenate the fold sequences associated to the matrices P_i together to form the sequence $\{T_k\}_{k=1}^\infty$. We address item (I) on the list, i.e. the allowability of the sequence, in Lemma 5.1. We now have a fold ray through the points $\{w_j\}_{j=1}^\infty$.

For density of the geodesic ray we use the automorphisms g_i related to the P_i . By ensuring that arbitrarily high powers of these automorphisms (hence matrices) occur in the sequence, we ensure that the ray passes through points with length vectors arbitrarily close to the dense set of eigenvectors.

Finally, property (III) on the list, that the fold line is a Lipschitz geodesic, follows from the fact that every g_i is a positive automorphism (see Corollary 3.18).

To extend our proof of Theorem C to Theorems A and B, we prove that for a generic point y in reduced Outer Space, there exist roses x, z and a “positive” fold line $[x, z]$ remaining in reduced Outer Space and so that $y \in [x, z]$. Here by “positive” we mean that the change of marking from x to z is a positive automorphism. Moreover, if G is the underlying graph of y and E, E' are two adjacent edges in G , then one may choose $[x, z]$ so that it contains the fold of the turn $\{E, E'\}$ immediately after the point y . This construction is carried out in §6 and elevates the geodesic’s density in $\mathcal{R}\mathcal{X}_r/\text{Out}(F_r)$ to density in \mathcal{U}_r . Additionally, the geodesic $[x, z]$ varies continuously as a function of y , as we prove in §7. Thus, one may adjust the previous argument to prove Theorems A and B, which we do in §8.

Acknowledgements. We wish to thank Pierre Arnoux for bringing our awareness to Brun’s algorithm and its possible use in proving that the Perron-Frobenius eigenvectors are dense, as well as other helpful discussions. On this subject we are also heavily indebted to Jon Chaika who pointed out to us that Brun’s algorithm is ergodic and that we could use this to elevate our proof of the Perron-Frobenius eigenvectors being dense in a simplex to additionally prove that Brun’s algorithm converges in angle (also somewhat simplifying our proof that the Perron-Frobenius eigenvectors are dense in a simplex). We would further like to thank Jayadev Athreya for posing the question and helpful discussions. Inspirational ideas given by Pascal Hubert were also particularly valuable. We are indebted to Lee Mosher for pointing out that Keane has a paper on a complication we were facing with regard to Brun’s algorithm. And we are indebted to Fritz Schweiger for his generosity in helping us understand arguments of his books. Finally, we are indebted to Valérie Berthé, Mladen Bestvina, Kai-Uwe Bux, Albert Fisher, Vincent Guirardel, Ilya Kapovich, Amos Nevo, and Stefan Witzel for helpful discussions.

CONTENTS

1. Introduction	1
2. Definitions and background	6
3. Fold paths and geodesics	11
4. Brun's algorithm and density of Perron-Frobenius eigenvectors	15
5. Dense Geodesics in theta complexes	21
6. Finding a rose-to-graph fold path terminating at a given point	23
7. Existence and continuity of rose-to-rose fold lines	32
8. Constructing the fold ray	35
9. Appendix: Limits of fold geodesics.	37
References	38

2. DEFINITIONS AND BACKGROUND

2.1. Outer Space and the action of $\text{Out}(F_r)$. Culler and Vogtmann introduced Outer Space in [CV86]. Points of Outer Space are “marked metric graphs.”

Definition 2.1 (Graph, positive edges). A *graph* will mean a connected 1-dimensional cell complex. $V(G)$ will denote the vertex set and $E(G)$ the set of unoriented edges. The degree of a vertex $v \in V(G)$ will be denoted $\deg_G(v)$, or $\deg(v)$ when G is clear.

For each edge $e \in E(G)$, one may choose an orientation. Once the orientation is fixed that oriented edge E will be called *positive* and the edge with the reverse orientation \bar{E} will be called *negative*. Given an oriented edge E , $i(E)$ will denote its initial vertex and $ter(E)$ its terminal vertex. A *directed* graph is a graph G with a choice of orientation on each edge $e \in E(G)$, we call this choice an *orientation* on G .

Given a free group F_r of rank $r \geq 2$, we choose once and for all a free basis $A = \{X_1, \dots, X_r\}$. Let $R_r = \vee_{i=1}^r \mathbb{S}^1$ denote the graph with one vertex and r edges, we call this graph an *r -petaled rose*. We choose once and for all an orientation on R_r and identify each positive edge of R_r with an element of the chosen free basis. Thus, a cyclically reduced word in the basis corresponds to an immersed loop in R_r .

Definition 2.2 (Marked metric F_r -graph). For each integer $r \geq 2$ we define a *marked F_r -graph* to be a pair $x = (G, m)$ where:

- G is a graph such that $\deg(v) \geq 3$ for each vertex $v \in V(G)$.
- $m: R_r \rightarrow G$ is a homotopy equivalence, called a *marking*.

A *marked metric graph* is a triple (G, m, ℓ) so that (G, m) is a marked graph and:

- The map $\ell: E(G) \rightarrow \mathbb{R}_+$ is an assignment of lengths to the edges. We require that $\sum_{e \in E(G)} \ell(e) = 1$. The quantity $\text{vol}(G) = \sum_{e \in E(G)} \ell(e)$ is called the *volume* of G .

Remark 2.3 (A metric graph as a metric space). The assignment of lengths to the edges does not quite determine a metric on G , but a homeomorphism class of metrics. This choice is inconsequential in this paper.

Define an equivalence relation on marked metric F_r -graphs by $(G, m, \ell) \sim (G', m', \ell')$ when there exists an isometry $\varphi: (G, \ell) \rightarrow (G', \ell')$ so that m' is homotopic to $\varphi \circ m$.

Definition 2.4 (Underlying set of Outer space). For each $r \geq 2$, as a set, the (*rank- r*) *Outer Space* \mathcal{X}_r is the set of equivalence classes of marked metric F_r -graphs.

Remark 2.5. On occasion we may think of graphs with valence-2 vertices as living in Outer Space by considering them equivalent to the graphs obtained by unsubdividing at their valence-2 vertices.

Definition 2.6. The *simplex* σ in \mathcal{X}_r corresponding to the marked graph (G, m) is

$$\sigma_{(G,m)} = \{(G, m, \ell) \in \mathcal{X}_r \mid \text{vol}(G) = 1\}.$$

By enumerating the edges of G , we can identify $\sigma_{(G,m)}$ with the open simplex

$$\mathcal{S}_{|E|} = \left\{ \vec{v} \in \mathbb{R}_+^{|E|} \mid \sum_{i=1}^{|E|} v_i = 1 \right\}.$$

Here $E = E(G)$. We denote this identification by $n: \sigma_{(G,m)} \rightarrow \mathcal{S}_{|E|}$. We call the open simplex corresponding to (R_r, id) the *base simplex* and denote it by σ_0 .

Outer Space has the structure of a simplicial complex built from open simplices (see [Vog02]), faces of $\sigma_{(G,m)}$ arise by letting the edges of a tree in G have length 0.

Definition 2.7 (Simplicial metric). Given a simplex $\sigma_{(G,m)}$ in \mathcal{X}_r , the *simplicial metric* on $\sigma_{(G,m)}$ is defined by $d_s(\ell, \ell') = \sqrt{\sum_{e \in E(G)} (\ell(e) - \ell'(e))^2}$, for $\ell, \ell' \in \sigma_{(G,m)}$. We also denote by d_s the extension of this metric to a path metric on \mathcal{X}_r .

Remark 2.8. In §2.4 we define the Lipschitz metric on \mathcal{X}_r . The simplicial metric and Lipschitz metric on Outer Space differ in important ways. However, open balls with respect to the Lipschitz metric (in either direction, see Remark 2.26) contain open balls of the simplicial metric). Therefore, a set dense with respect to the simplicial topology will also be dense with respect to the Lipschitz topology.

Definition 2.9 (Unprojectivized Outer Space). [CV86] The (*rank- r*) *unprojectivized Outer Space* $\hat{\mathcal{X}}_r$ is the space of metric marked F_r -graphs where $\text{vol}(G)$ is not necessarily 1.

There is a map from $\hat{\mathcal{X}}_r$ to \mathcal{X}_r normalizing the graph volume, i.e.

$$(1) \quad \begin{aligned} q: \mathbb{R}_+^m &\rightarrow \mathcal{S}_m \\ q(x_1, \dots, x_m) &= \left(\frac{x_1}{\sum_{i=1}^m x_i}, \dots, \frac{x_m}{\sum_{i=1}^m x_i} \right) \end{aligned}$$

and taking the point (G, μ, ℓ) to the point $(G, \mu, q(\ell))$.

We call the full preimage under q of a simplex in \mathcal{X}_r an *unprojectivized simplex*.

Definition 2.10 (Topological Outer Space). [CV86] The topological space consisting of the set of equivalence classes of marked metric F_r -graphs, endowed with the simplicial topology, is called the (*rank- r*) *Outer Space* and is also denoted by \mathcal{X}_r .

Definition 2.11 ($\text{Out}(F_r)$ action). If $\Phi \in \text{Aut}(F_r)$ is an automorphism, let $f_\Phi: R_r \rightarrow R_r$ be a homotopy equivalence corresponding to Φ via the identification of $E(R_r)$ with the chosen free basis A of F_r . We define a *right action of $\text{Out}(F_r)$* on \mathcal{X}_r . An outer automorphism $[\Phi] \in \text{Out}(F_r)$ acts by $[G, m, \ell] \cdot [\Phi] = [G, m \circ f_\Phi, \ell]$.

Definition 2.12 (Reduced Outer Space $\mathcal{R}\mathcal{X}_r$ and \mathcal{M}_r). For each integer $r \geq 2$, the (*rank- r*) *reduced Outer Space* $\mathcal{R}\mathcal{X}_r$ is the subcomplex of \mathcal{X}_r consisting of precisely those simplices $\sigma_{(G,m)}$ such that G contains no separating edges. This space is connected and an $\text{Out}(F_r)$ -equivariant deformation retract of \mathcal{X}_r .

Let \mathcal{M}_r denote the quotient space of $\mathcal{R}\mathcal{X}_r$ by the $\text{Out}(F_r)$ action. Hence, \mathcal{M}_r contains a quotient of a simplex for each graph (no longer marked). Note that, as a result of graph symmetries, simplices in \mathcal{X}_r do not necessarily project to simplices in \mathcal{M}_r . Thus, \mathcal{M}_r is no longer a simplicial complex but a union of cells which are quotients of simplices in \mathcal{X}_r .

2.2. Train track structures. Much of the following definitions and theory can be found in [BH92] or [Bes12], for example. However, it should be noted that some of our definitions, including that of an illegal turn, are somewhat nonstandard.

Definition 2.13 (Regular maps). We call a continuous map $f: G \rightarrow H$ of graphs *regular* if for each edge $e \in E(G)$, we have that $f|_{\text{int}(e)}$ is locally injective and that f maps vertices to vertices.

Definition 2.14 (Paths and loops). Depending on the context an *edge-path* in a graph G will either mean a continuous map $[0, n] \rightarrow G$ that, for each $1 \leq i \leq n$, maps $(i-1, i)$ homeomorphically to the interior of an edge, or if the graph G is directed, a sequence of oriented edges e_1, \dots, e_n such that $\text{ter}(e_i) = i(e_{i+1})$ for each $1 \leq i \leq n-1$. We may on occasion also allow for e_1 and e_n to be partial edges. Given any path $\gamma = e_1 \cdots e_n$, we will denote its initial vertex, i.e. $i(e_1)$, by $i(\gamma)$ and its terminal vertex, i.e. $\text{ter}(e_n)$ by $\text{ter}(\gamma)$.

A *loop* α in G is the image of an immersion $\alpha: \mathbb{S}^1 \rightarrow G$. We will associate to each loop an edge-path unique up to cyclic ordering.

In a directed graph G , we will call a path *directed* that either crosses all edges in a positive direction (a *positive path*) or crosses all edges in a negative direction (a *negative path*). The operation of path concatenation will be denoted by $*$.

Definition 2.15 (Illegal turns and gates). Let $f: G \rightarrow H$ be a regular map. Let $e, e' \in E(G)$ be oriented edges with the same initial vertex. We call $\{e, e'\}$ a *turn*. We say a turn $\{e, e'\}$ is an *illegal turn* for f if the first edge of the edge-path $f(e)$ equals the first edge of the path $f(e')$. The property of forming an illegal turn is an equivalence relation and the equivalence classes are called *gates*.

Definition 2.16 (Train track structures). Let $f: G \rightarrow H$ be a regular map. If every vertex of G has ≥ 2 gates, then we call the partition of the turns of G into gates a *train track structure* and say that f induces a *train track structure* on G . An immersed path $\alpha: I \rightarrow G$ will be considered *legal* with respect to a given train track structure if it does not contain a subpath $e_i e_j$ where $\{\bar{e}_i, e_j\}$ is an illegal turn.

Remark 2.17. The image of a legal path will be locally embedded.

Definition 2.18 (Transition matrix). The *transition matrix* of a regular self-map $f: G \rightarrow G$ is the square $|E(G)| \times |E(G)|$ matrix (a_{ij}) such that a_{ij} , for each i and j , is the number of times $g(e_i)$ passes over e_j in either direction.¹

We define the *transition matrix* for an element $\Phi \in \text{Aut}(F_r)$ to be the transition matrix of f_Φ (see Definition 2.11).

2.3. Perron-Frobenius theory. We are particularly interested in positive matrices (defined below) because of their known properties (due to Perron-Frobenius theory) of contracting, the *positive cone* $\mathbb{R}_+^d = \{v \in \mathbb{R}^d \mid v_i > 0, i = 1, \dots, d\}$.

Definition 2.19 (Positive matrices, Perron-Frobenius eigenvalues and eigenvectors). We call a matrix $A = [a_{ij}]$ *positive* if each entry of A is strictly positive. By Perron-Frobenius theory, we know that each such matrix has a unique eigenvalue of maximal modulus and that this eigenvalue is real. This eigenvalue is called the *Perron-Frobenius (PF) eigenvalue* of A . It has an associated eigenvector whose entries are each strictly positive. We call the eigenvector with strictly positive entries and such that all entries sum to one the *Perron-Frobenius (PF) eigenvector*.

Definition 2.20 (Weak convergence). A sequence $\{A_k\}_{k=1}^\infty$ of $d \times d$ matrices, restricted to vectors in \mathbb{R}_+^d , *converges weakly* if the sequence $\{A_k(\mathbb{R}_+^d)\}_{k=1}^\infty$ converges projectively to a point.

Remark 2.21. Perron-Frobenius theory also tells us that, for a positive matrix M , the sequence $\{M^k\}_{k=1}^\infty$ weakly converges to the line spanned by the PF eigenvector.

¹This matrix is the transpose of the transition matrix as Bestvina and Handel define it in [BH92], but this definition will have a stronger relationship with the change-of-metric matrix we define later.

2.4. The Lipschitz metric.

Definition 2.22 (Difference in markings). Let $x = (G, m, \ell)$ and $y = (G', m', \ell')$ be two points in \mathcal{X}_r . Denote by $h: G \rightarrow R_r$ a homotopy inverse of m . A *difference in markings* is a linear map $f: G \rightarrow G'$ homotopic to $m' \circ h$.

Definition 2.23 (Stretch). Let α be a conjugacy class in F_r , equipped with a free basis A . By abuse of notation we may think of α as an immersed loop $\alpha: \mathbb{S}^1 \rightarrow R_r$ in R_r via the identification of the edges of R_r with A . For $x = (G, m, \ell) \in \mathcal{X}_r$, let α_x denote the unique immersed simplicial loop in G homotopic to $m(\alpha)$.

Given a conjugacy class α in F_r and $x \in \mathcal{X}_r$, we define $l(\alpha, x)$ as the length of α_x . (Notice that since α_x is a simplicial loop in x , its length does not depend on the particular metric structure chosen for x , see Remark 2.3). For $x, y \in \mathcal{X}_r$ define the *stretch* of α from x to y as $\text{st}_\alpha(x, y) := \frac{l(\alpha, y)}{l(\alpha, x)}$.

The following theorem is attributed to either White or Thurston. It can be found in [AK11].

Theorem 2.24. *Given a continuous map f of metric spaces, let $\text{Lip}(f)$ denote the Lipschitz constant for f . Then for each pair of points $x, y \in \mathcal{X}_r$, we have*

$$(2) \quad \inf\{\text{Lip}(f) \mid f: x \rightarrow y \text{ is a difference in marking}\} = \sup\{\text{st}_\alpha(x, y) \mid \alpha \in F_r\}.$$

Moreover, both the infimum and supremum are realized.

Definition 2.25 (Lipschitz metric). The *Lipschitz metric* $d(x, y)$ is defined as the log of either of the quantities in Equation 2. This function is not symmetric but satisfies the other axioms of a metric [Bes12].

A difference in marking that achieves the minimum Lipschitz constant of (2) is called an *optimal map*. A loop that achieves the maximum stretch is called a *witness*. For each $x, y \in \mathcal{X}_r$ there exist optimal maps and witnesses.

Remark 2.26. An open ball based at x with radius R is either

$$B_{\rightarrow}(x, R) = \{y \in \mathcal{X}_r \mid d(x, y) < R\} \text{ or } B_{\leftarrow}(x, R) = \{y \in \mathcal{X}_r \mid d(y, x) < R\}.$$

In either case, the simplicial topology is finer (or equal) to the Lipschitz topology.

For a given difference of marking f , the subgraph of G where the Lipschitz constant is achieved is called the *tension graph*, usually denoted Δ_f . Notice that f induces a train track structure on Δ_f . Proposition 2.27 gives one way to identify witnesses.

Proposition 2.27. [BF14] *Let x and y be two points in \mathcal{X}_r , let $f: x \rightarrow y$ be a map, and let Δ_f be the tension graph of f . If Δ_f contains a legal loop, then f is an optimal map and any legal loop in Δ_f is a witness. Conversely, if $\alpha \subset x$ is a witness, then it is a legal loop in Δ_f .*

Proposition 2.28. *Let $f: x \rightarrow y$ and $g: y \rightarrow z$ be difference in markings maps. Let α be a conjugacy class in F_r satisfying that α_x is f -legal and contained in Δ_f and that α_y is g -legal and contained in Δ_g . Then $d(x, z) = d(x, y) + d(y, z)$.*

Proof. Since α_x is contained in Δ_f , the map f stretches each edge of α_x by $\lambda_f = \text{Lip}(f)$. Moreover, since α_x is f -legal, $l(\alpha, y) = \lambda_f \cdot l(\alpha, x)$. Similarly, if $\lambda_g = \text{Lip}(g)$, then $l(\alpha, z) = \lambda_g \cdot l(\alpha, y) = \lambda_f \lambda_g \cdot l(\alpha, x)$. Thus, $\text{st}_\alpha(x, z) = \lambda_f \lambda_g$ and therefore $d(x, z) \geq \log(\lambda_f \lambda_g)$. By Proposition 2.27, α_x and α_y are both witnesses, hence $d(x, y) = \text{st}_\alpha(x, y) = \log \lambda_f$ and $d(y, z) = \text{st}_\alpha(y, z) = \log \lambda_g$. Thus, we have $d(x, z) \geq \log \lambda_f + \log \lambda_g = d(x, y) + d(y, z)$. The triangle inequality gives us an equality. \square

3. FOLD PATHS AND GEODESICS

In this section we introduce fold lines and prove results that will allow us to construct Lipschitz geodesics from certain infinite sequences of nonnegative matrices (unfolding matrices).

Definition 3.1 (Unparametrized geodesic). Let $I \subset \mathbb{R}$ be a generalized interval in \mathbb{R} . An *unparametrized geodesic* in \mathcal{X}_r is a map $\Gamma: I \rightarrow \mathcal{X}_r$ satisfying that:

- (1) for each $s < r < t$, we have $d(\Gamma(r), \Gamma(t)) = d(\Gamma(r), \Gamma(s)) + d(\Gamma(s), \Gamma(t))$ and
- (2) there exists no nontrivial subinterval $I' \subset I$ and point $x_0 \in \mathcal{X}_r$ such that $\Gamma(t) = x_0$ for each $t \in I'$.

Remark 3.2. If Γ is an unparametrized geodesic then there exists a generalized interval I' and a homeomorphism $h: I' \rightarrow I$ so that $\Gamma \circ h$ is an honest directed geodesic, i.e. for all $s < t$, we have that $d(\Gamma \circ h(s), \Gamma \circ h(t)) = t - s$.

Again $A = \{X_1, \dots, X_r\}$ will denote a fixed free basis of F_r .

Definition 3.3 (Fold automorphism). By a *fold automorphism* we will mean a “left Nielsen generator,” i.e. an automorphism of the following form ($i \neq j$):

$$(3) \quad \Phi_{ij}(X_k) = \begin{cases} X_j X_k & \text{for } k = i \\ X_k & \text{for } k \neq i. \end{cases}$$

By notation abuse Φ_{ij} will also denote the map $R_r \rightarrow R_r$ that corresponds to the above automorphism after identifying its positively oriented edges with the basis A .

To a fold automorphism one can associate a matrix.

Definition 3.4 ((Un)folding matrix). Let $i \neq j \in \{1, \dots, r\}$. Then the (i, j) *folding matrix* T_{ij} has entries t_{kl} , where

$$(4) \quad t_{kl} = \begin{cases} 1 & \text{if } k = l \\ -1 & \text{if } (k, l) = (i, j) \\ 0 & \text{otherwise} \end{cases}$$

Notice that T_{ij} is not the transition matrix of f_{ij} , though it will relate to the change-of-metric matrix coming from a folding operation. The matrix T_{ij} is invertible and we call $M_{ij} := T_{ij}^{-1}$ the (i, j) *unfolding matrix*. Notice that the entries of M_{ij} are m_{kl} , where

$$(5) \quad m_{kl} = \begin{cases} 1 & \text{if } k = l \text{ or } (k, l) = (i, j) \\ 0 & \text{otherwise.} \end{cases}$$

Notice also that the nonnegative *matrix* M_{ij} is the transition matrix of Φ_{ij} . We hence sometimes write $M(\Phi_{ij})$ for this matrix.

Definition 3.5 (A combinatorial fold). Let G be a graph whose oriented edges are numbered. Let (e_k, e_j) be a pair of distinct oriented edges with $i(e_k) = i(e_j)$. A combinatorial fold is a tuple $(G, (e_k, e_j), G', f)$ where G' is a graph and $f: G \rightarrow G'$ is a quotient map that identifies an initial segment of e_k with an initial segment of e_j .

- (1) when f identifies part of e_k with all of e_j we will call it a *proper full fold*. We will call this tuple “folding e_k over e_j .”
- (2) when f identifies all of e_k with all of e_j we will call it a *full fold* and say that “ e_k and e_j are fully folded.”
- (3) when f identifies a proper segments of e_k and e_j we will call it a *partial fold* and say that “ e_k and e_j are partially folded.”

Notation We sometimes repress some of the data depending on the context. We will denote the combinatorial fold by f , or $G \rightarrow G'$, or $(G, (e_i, e_j))$ depending on which data we want to emphasize.

Definition 3.6 (Direction matching folds). Let G be an oriented graph. A combinatorial fold $(G, (e_i, e_j))$ is *direction matching* if e_1 and e_2 are either both negative edges (we then call f *negative*) or both positive edges (we then call f *positive*).

Observation 3.7. Let G be an oriented graph and $f: G \rightarrow H$ a direction matching combinatorial fold, then f induces an orientation on the edges of H . Moreover, f maps each positive edge of G to a positive edge-path in H (of simplicial length ≤ 2).

Definition 3.8 (Allowable folds). Let $x_0 = (G, m, \ell)$ be a point in Outer Space. The full fold $(G, (e, e'))$ is *allowable* in x_0 if the following two conditions hold:

- (1) $\ell(e) \geq \ell(e')$.
- (2) If $\text{ter}(e) = \text{ter}(e')$, then $\ell(e) > \ell(e')$. In this case this is a *proper full fold*.

Suppose G is a rose with its edges enumerated. Let $n_\tau: \sigma_{(G,m)} \rightarrow \mathcal{S}_{|E|}$ be the induced homeomorphism. The set where Φ_{ij} is allowable will be denoted by $\sigma_{(G,m)}^{(i,j)}$. The target graph G' is also a rose and Φ_{ij} induces an enumeration of the edges of G' by declaring the edge $\Phi_{ij}(e_k)$ to be the k^{th} edge for each $k \neq i$ and calling the remaining edge of G' the i^{th} edge.

Lemma 3.9. *Let $\hat{\sigma}_0$ be the unprojectivized base simplex, and let Φ be a fold automorphism. Assume that the edges of R_r have been enumerated and let $n_0: \hat{\sigma}_0 \rightarrow \mathcal{S}_r$ be the induced identification. $\Phi: \hat{\sigma}_0 \rightarrow \hat{\sigma}_1$ induces a new enumeration of the edges of the target rose and we get a new identification $n_1: \hat{\sigma}_1 \rightarrow \mathcal{S}_r$. Then for each $x \in \hat{\sigma}_1$ we have $n_1(x) = n_0(x \cdot \Phi^{-1})$ (as defined in Definition 2.11).*

If a fold $(G, (e, e'))$ is allowable at a point x_0 in Outer Space, one can construct a path $\{\hat{x}_t\}_{t \in [0,1]}$ in unprojectivized outer space, called a “fold path.” This is done by identifying initial segments of e and e' of length $t\ell_0(e')$, for $0 \leq t \leq 1$. The quotient map $f_{t,0}: x_0 \rightarrow x_t$ is a homotopy equivalence, as are the quotient maps $f_{t,s}$ for $0 \leq s \leq t \leq 1$. By projectivizing we get a path $\{x_t\}$ in Outer Space.

Definition 3.10 (Basic fold paths). Given an allowable fold as above, the path $\mathcal{F}: [0, 1] \rightarrow \mathcal{X}_r$ defined by $t \mapsto x_t$ is the *fold path in \mathcal{X}_r starting at x_0 and defined by folding e over e'* . The path $t \rightarrow \hat{x}_t$ will be called the *(unprojectivized) basic fold path*. We will not always use the hat notation when discussing unprojectivized paths but will mention whether the image lies in \mathcal{X}_r or $\hat{\mathcal{X}}_r$, if it is otherwise unclear.

Definition 3.11 (Change-of-metric matrix). Let G, G' be graphs and assume we have numbered their oriented edges. Let Ψ be a linear map from a subset of the unprojectivized simplex $\hat{\sigma}_{(G,\mu)}$ to the unprojectivized simplex $\hat{\sigma}_{(G',\mu')}$. Then Ψ may be represented by an $|E(G')| \times |E(G)|$ matrix. This matrix will be called the *change-of-metric matrix*.

Lemma 3.12. *Let Φ be an allowable fold automorphism on the point $x_0 = (R_r, m, \ell)$ and suppose the edges of R_r have been numbered so that $\Phi = \Phi_{ij}$. Let x_1 be the folded graph and suppose $\hat{\sigma}_0$ and $\hat{\sigma}_1$ are unprojectivized open simplices containing respectively x_0 and x_1 . The change-of-metric matrix for the fold operation from $n_0(\hat{\sigma}_{(R,m)}^{(i,j)})$ to $n_1(\hat{\sigma}_{(R,\Phi \circ m)})$ is the matrix T_{ij} of Definition 3.4.*

Definition 3.13 (Fold paths). A *fold path* $\mathcal{F}: [0, k] \rightarrow \mathcal{X}_r$ is a path in \mathcal{X}_r that may be divided into a sequence of basic fold paths $\{\mathcal{F}_i\}_{i=1}^k$ as in Definition 3.10, so that $\mathcal{F}_i(1) = \mathcal{F}_{i+1}(0)$ for each i . As above, we will denote by $x_t = (G_t, m_t, \ell_t)$ the points of the path in \mathcal{X}_r . The maps $f_{t,s}$ for $s, t \in [0, k]$ will be defined similarly as above.

If \mathcal{F} is a fold path from $\mathcal{F}(0) = x$ to $\mathcal{F}(k) = y$ we sometimes denote it by $\mathcal{F}: x \rightarrow y$. The following Lemma follows from Lemma 3.12 and its proof is left to the reader.

Lemma 3.14. *Let f_1, \dots, f_k be a sequence of combinatorial proper full folds of the r -rose, with associated change-of-metric matrices T_1, \dots, T_k having respective inverse matrices M_1, \dots, M_k . Suppose that $v \in M_1 \cdots M_k(\mathbb{R}_+^r)$. Then the combinatorial fold f_l is allowable in the metric graph $n_l^{-1}(T_l \cdots T_1(v))$, for each $2 \leq l \leq k$ (and any marking). Furthermore, applying f_1, \dots, f_k to $x_0 \in \sigma_0$ will result in the point $n_k^{-1}(T_k \cdots T_1(n_0(x_0)))$ of the simplex $\sigma_{(R_r, f_k \circ \dots \circ f_1 \circ m)}$.*

Lemma 3.15. *Let $\{D_i\}_{i=1}^\infty$ denote a sequence of nonnegative invertible matrices such that, for some $N \in \mathbb{N}$, we have that $D_1 \cdots D_i$ is strictly positive for all $i \geq N$. Then there exists a vector $w_0 \in \mathbb{R}_+^r$ so that, if we define $w_{l+1} := D_{l+1}^{-1}w_l$, for each l , then each w_l is a positive vector.*

Proof. Let $\overline{\mathbb{R}_+^r}$ denote the set of vectors with nonnegative entries, and recall the map q from Equation 1. Let $M = D_1 \cdots D_N$, by the assumption it is a nonnegative matrix. Consider $\mathcal{I} = \bigcap_{i=1}^\infty D_1 \cdots D_i(\mathbb{R}_+^r)$. Note that

$$\mathcal{I} \supset \bigcap_{i=1}^\infty D_1 \cdots D_i(M(\overline{\mathbb{R}_+^r})) \supset \bigcap_{i=1}^\infty q(D_1 \cdots D_i(M(\overline{\mathbb{R}_+^r}))$$

and the latter is nonempty since it is an intersection of nested compact sets. Moreover, $\mathcal{I} \subset M(\mathbb{R}_+^r) \subset \mathbb{R}_+^r$. Choose $w_0 \in \mathcal{I}$. Then $w_0 \in \mathcal{I} \subset D_1 \cdots D_l(\mathbb{R}_+^r)$ implies that $w_l = D_l^{-1} \cdots D_1^{-1}w_0 \in \mathbb{R}_+^r$ for each $1 \leq l < \infty$. \square

Corollary 3.16. *Let $\{f_i\}_{i=0}^\infty$ be a sequence of combinatorial proper full folds of the r -rose, with associated change-of-metric matrices $\{T_i\}_{i=0}^\infty$ having respective inverse matrices $\{M_i\}_{i=0}^\infty$. If there exists some $N \in \mathbb{N}$, so that for all $i > N$ the matrix $D_1 \cdots D_i$ is strictly positive, then there exists a vector $w_0 \in \mathbb{R}_+^r$ so that the infinite fold sequence $\{f_i\}_{i=0}^\infty$ is allowable in the rose $x_0 = n_0^{-1}(w_0)$.*

Proposition 3.17. *Let $\{\mathcal{F}_i: x_i \rightarrow x_{i+1}\}_{i=0}^k$ be a sequence of fold paths with fold maps $\{f_{s,t}\}_{s \geq t \geq 0}$. Suppose there is a conjugacy class α in F_r satisfying that, for each i , the realization α_{x_i} of α in x_i is legal with respect to the train track structure induced by $f_{i+1,i}$. Then the corresponding fold path $\text{Im}(\mathcal{F}) = \{x_t\}_{t \in [0,k]}$ is an unparametrized geodesic, i.e. for each $r \leq s \leq t$ in $[0, k]$, we have $d(x_r, x_t) = d(x_r, x_s) + d(x_s, x_t)$.*

Proof. The proof uses Propositions 2.27 and 2.28 and is left to the reader. \square

Suppose y_0 is a graph and $f_0: y_0 \rightarrow y_1$ is direction matching, then by Observation 3.7, we have that y_1 inherits an orientation such that the image of each edge is a positive edge-path (see Definition 2.14).

Corollary 3.18. *Let y_0 be a directed metric r -rose graph with length vector v and let $\{f_i: y_i \rightarrow y_{i+1}\}_{i=0}^{i=k}$ be an allowable sequence of proper full folds. Suppose that for each $i = 0, \dots, k-1$ the fold f_{i+1} is direction matching with respect to the orientation*

of y_{i+1} inherited by f_i . For each $0 \leq i \leq k$, let $\{\mathcal{F}_i: y_i \rightarrow y_{i+1}\}$ denote the fold path determined by $\{f_i\}$. Then the corresponding fold path $\text{Im}(\mathcal{F}) = \{y_t\}_{t \in [0, k]}$ is an unparametrized geodesic.

Proof. Without generality loss we can assume that the marking on R_r is the identity. Then, by Proposition 3.17, it suffices to show that there exists a conjugacy class α in F_r satisfying that, for each i , the realization α_{y_i} of α in y_i is legal with respect to the train track structure induced by the map f_i . We claim that this holds for the conjugacy class α of a positive generator X_1 . By induction suppose that α_{y_i} is a positive loop. Since f_i maps positive edges to positive edge-paths, $\alpha_{y_{i+1}}$ is also a positive loop. Since f_{i+1} is direction matching, $\alpha_{y_{i+1}}$ is f_{i+1} -legal and positive. \square

4. BRUN'S ALGORITHM AND DENSITY OF PERRON-FROBENIUS EIGENVECTORS

We introduce fibered systems so that we can use an algorithm of Brun to prove, in Theorem 4.14, that the sequence of Brun matrices converge weakly for a full measure set of points. These matrices are unfolding matrices, which will be significant for proving Theorems A, B, and C in the next sections.

4.1. Fibered Systems. The following definitions are taken from [Sch00].

Definition 4.1 (Fibered systems). A pair (B, T) is called a *fibered system* if B is a set, $T: B \rightarrow B$ is a map, and there exists a partition $\{B(i) \mid i \in I\}$ of B such that I is countable and $T|_{B(i)}$ is injective. The sets $B(i)$ are called *time-1 cylinders*.

Definition 4.2 (Time- s cylinder). For each $x \in B$, one can define a sequence $\Phi(x) = (i_1(x), i_2(x), \dots) \in I^{\mathbb{N}}$ by letting $i_s(x) = i \iff T^{s-1}x \in B(i)$. In other words, $i_s(x)$ tells us which cylinder $T^{s-1}x$ lands in. Then a *time- s cylinder* is a set of the form

$$B(i_1, \dots, i_s) = \{x \in B \mid i_1(x) = i_1, \dots, i_s(x) = i_s\}.$$

From the definitions we have $B(i_1, \dots, i_{s+1}) = B(i_1, \dots, i_s) \cap T^{-s}B(i_{s+1})$.

4.2. The unordered Brun algorithm in the positive cone. The following algorithm (commonly referred to as *Brun's algorithm*) was introduced by Brun [Bru57] as an analogue of the continued fractions expansion of a real number in dimensions 3 and 4. It was later extended to all dimensions by Schweiger in [Sch82].

Definition 4.3 (Brun's (unordered) algorithm). *Brun's unordered algorithm* is the fibered system (C^n, T) defined on

$$C^n = \mathbb{R}_+^n = \{(x_1, \dots, x_n) \in \mathbb{R}_+^n \mid x_i > 0 \text{ for all } 1 \leq i \leq n\}$$

by

$$\begin{aligned} T: C^n &\rightarrow C^n \\ (x_1, \dots, x_n) &\mapsto (x_1, \dots, x_{m-1}, x_m - x_s, x_{m+1}, \dots, x_n), \end{aligned}$$

where

$$m(x) := \min \left\{ i \mid x_i = \max_{1 \leq j \leq n} x_j \right\}, \quad s(x) := \min \left\{ i \neq m(x) \mid x_i = \max_{1 \leq j \neq m(x) \leq n} x_j \right\}.$$

In other words, for each $x = (x_1, \dots, x_n) \in C^n$, we have that $m(x)$ is the first index that achieves the maximum of the coordinates and $s(x)$ is the first index that achieves the maximum of all of the coordinates except for $m(x)$.

Notice that, letting $(i, j) = (m(x), s(x))$, the transformation T is just left multiplication by the matrix $T_{i,j}$ from Definition 3.4. Then, given a vector $v_0 = (x_1, \dots, x_n) \in C^n$ with rationally independent coordinates, one obtains an infinite sequence

$$\{v_k = (x_1^k, \dots, x_n^k)\}_{k=1}^\infty \subset C^n,$$

where v_{k+1} is recursively defined by $v_{k+1} = T_{m(v_k), s(v_k)} v_k$.

Definition 4.4 (Brun sequence). In light of the above, the sequence for Brun's unordered algorithm (which we call the *Brun sequence*) will consist of the ordered pairs $\vec{k}_s = (i_s, j_s)$, where $(i_s, j_s) = (m(v_s), s(v_s))$. Further, the sequence $\Theta(x) = \{\vec{k}_s\}_{s=1}^\infty$ will determine a sequence of folding matrices, which we denote by $\{T_s^x\}_{s=1}^\infty$, where $T_s^x = T_{\vec{k}_s(x)}$. We let $\{M_s^x\}_{s=1}^\infty$ denote the corresponding inverses, i.e. the unfolding matrices. We denote their finite products by

$$(6) \quad A_s^x = M_1^x M_2^x \cdots M_s^x \text{ for each } s \in \mathbb{N}.$$

Then, as above, for each $v_0 \in C^n$, we have a sequence of vectors $\{v_s\}_{s=1}^\infty \subset C^n$, where $v_{s+1} = T v_s = T_{\vec{k}_s(v_0)} v_s$. Thus, $v_s = M_{s+1}^{v_0} v_{s+1}$, and hence $v_0 = A_s^{v_0} v_s$. So

$$(7) \quad v_0 \in \bigcap_{s=1}^\infty A_s^{v_0}(C^n).$$

This fact will become particularly important in the proof of Theorem 4.5.

Definition 4.5 (Brun matrix). For each vector $v_0 \in C^r$, we call each matrix

$$(8) \quad A_n = M_{i_1, j_1}^{v_0} \cdots M_{i_n, j_n}^{v_0}$$

a *Brun matrix*. We let \mathcal{B}_r denote the set of $r \times r$ Brun matrices.

Remark 4.6. When it is clear from the context, we may leave out reference to the starting vector v_0 and simply write A_s, T_s, M_s , etc.

Proposition 4.7. *Let Y_r be the set of rationally independent vectors in C^r , then for each $x \in Y_r$ there exists an $N \in \mathbb{N}$ so that A_n^x is a positive matrix for each $n > N$.*

Proof. We fix v_0 and omit the index v_0 from the notation below. We first prove that, for each i , and for each $h \in \mathbb{N}$, there exists some $m > h$ and some j such that $\vec{k}_m = (i, j)$. Starting with v_h and until $(v_m)_i$ becomes the largest coordinate, at each step one subtracts a number $\geq (v_m)_i = (v_h)_i$ from some coordinate $\geq (v_m)_i$. This can only happen a finite number of times before each coordinate apart from $(v_m)_i$ becomes less than $(v_m)_i = (v_h)_i$.

Consider A_n as in (6). To prove the proposition, it suffices to show that, for each (i, j) , there exists a large enough N so that for all $n > N$ the (i, j) -th entry of A_n is positive. In fact, it is enough to show that this entry is positive for some A_n . Indeed A_{n+1} is obtained from A_n by adding one of its columns to another one of its columns, so that an entry positive in A_n , will be positive in A_{n+1} .

Fix i, j . Let

$$a = \min\{t \mid \vec{k}_t = (i, c) \text{ for some } c\}, \quad b = \min\{t > a \mid \vec{k}_t = (j, d) \text{ for some } d\}.$$

Let c_1 be such that $\vec{k}_a = (i, c_1)$. Observe that, since c_1 is the second largest coordinate in v_a , in the next vector v_{a+1} , either i is still the largest coordinate or c_1 becomes the largest coordinate. There is some $N_1 \geq 1$ and some index c_2 so that

$$A_{a+N_1} = A_{a-1} M_{(i, c_1)}^{N_1} M_{(c_1, c_2)}.$$

We continue in this way, $A_n = A_{a-1} M_{(i, c_1)}^{N_1} M_{(c_1, c_2)}^{N_2} M_{(c_2, c_3)}^{N_3} \cdots M_{(c_t, c_{t+1})}^{N_t}$. When $n = b$, we have that $c_t = j$. Thus, for $n = b - 1$, we have

$$A_{b-1} = A_{a-1} M_{(i, c_1)}^{N_1} M_{(c_1, c_2)}^{N_2} M_{(c_2, c_3)}^{N_3} \cdots M_{(c_t, j)}^{N_t}.$$

It is elementary to see that the (i, j) -th entry of $M_{(i, c_1)}^{N_1} M_{(c_1, c_2)}^{N_2} M_{(c_2, c_3)}^{N_3} \cdots M_{(c_t, j)}^{N_t}$ is positive. This implies that the (i, j) -th entry of A_{b-1} is positive. \square

4.3. Other versions of Brun's algorithm. To use the results of Schweiger's books, we must give two other different, but related, versions of Brun's algorithm.

Definition 4.8 (Brun's ordered algorithm). *Brun's ordered algorithm* is the fibered system (Δ^n, T') defined on $\Delta^n := \{x \in C^n \mid x_1 \geq \cdots \geq x_n\}$ by

$$\begin{aligned} T' : \Delta^n &\rightarrow \Delta^n \\ (x_1, \dots, x_n) &\mapsto (x_2, \dots, x_{i-1}, x_1 - x_2, x_i, \dots, x_n), \end{aligned}$$

where $i = i(x) \geq 2$ is the first index so that $x_1 - x_2 \geq x_i$ and, if there is no such index, we let $i(x) = n$. The time-1 cylinders are $\Delta^n(i) = \{x \mid x_{i-1} > x_1 - x_2 \geq x_i\}$.

Notice that the transformation T' is just left multiplication by the matrix $T_{1,2}$, followed by a permutation matrix that we denote P_i , determined by the cylinder $\Delta^n(i)$. (We denote $P_i T_{1,2}$ by T'_i and its inverse by M'_i .) Hence, given a sequence of indices $\omega(x) = (i_1, i_2, \dots)$ with $1 \leq i_j \leq n$ for each i_j , one obtains a sequence

of matrices $\{T'_{i_j}\}_{j=1}^\infty$. Given $v_0 \in \Delta^n(i_1, \dots, i_m)$, this gives a sequence of points $v_0, \dots, v_m \in \Delta^n$ such that $v_{k+1} = T'_{i_j} v_k$ for each $1 \leq k \leq m-1$. The fibered system sequence here will be $\omega(x) = (i_1, i_2, \dots)$ when

$$x \in \bigcap_{m=1}^{\infty} \Delta^n(i_1, i_2, \dots, i_m).$$

If $\omega(x) = (i_1, i_2, \dots)$, we define

$$(9) \quad A_k^x = M'_{i_1} \cdots M'_{i_k}$$

for each $k \in \mathbb{N}$.

Definition 4.9 (Brun's homogeneous algorithm). *Brun's homogeneous algorithm* is the fibered system $(B_n, \overline{T'})$ defined on

$$B_n = \{x \in \mathbb{R}^n \mid 1 \geq x_1 \geq x_2 \geq \cdots \geq x_n \geq 0\}$$

and where $\overline{T'}: B_n \rightarrow B_n$ is such that the following diagram commutes:

$$\begin{array}{ccc} \Delta^{n+1} & \xrightarrow{T'} & \Delta^{n+1} \\ \downarrow p & & \downarrow p \\ B_n & \xrightarrow{\overline{T'}} & B_n \end{array}$$

where $p: \Delta^{n+1} - \{0\} \rightarrow B_n$ is defined by

$$(10) \quad p(x_1, \dots, x_{n+1}) = \left(\frac{x_2}{x_1}, \dots, \frac{x_{n+1}}{x_1} \right).$$

We denote the time-1 cylinders by $B_n(i)$.

4.4. Relating the algorithms.

Definition 4.10. Let $\mathcal{O}: C^n \rightarrow \Delta^n$ be defined by

$$(x_1, \dots, x_n) \mapsto (x_{i_1}, \dots, x_{i_n})$$

where $x_{i_1} \geq x_{i_2} \geq \cdots \geq x_{i_n}$. Note that for a particular x , $\mathcal{O}(x)$ is a permutation.

Lemma 4.11. *For each $x \in C^n$ and for each $m \in \mathbb{N}$, there exist permutation matrices P_{i_1}, P_{i_2} so that*

$$A_m^x = P_{i_1} (A_m^{\mathcal{O}(x)})' P_{i_2}.$$

Proof. This follows from the following commutative diagram:

$$\begin{array}{ccc} C^r & \xrightarrow{T} & C^r \\ \downarrow \mathcal{O} & & \downarrow \mathcal{O} \\ \Delta^r & \xrightarrow{T'} & \Delta^r \end{array} .$$

□

Corollary 4.12. *For each $x \in C^n$ and for each $m \in \mathbb{N}$, we have that A_m^x is a positive matrix if and only if $(A_m^{\mathcal{O}(x)})'$ is a positive matrix.*

Corollary 4.13. *For each irrational $x \in \Delta^n$ there exists an N so that $(A_n^x)'$ is positive for all $n > N$.*

Proof. This follows from Proposition 4.7 and Corollary 4.12. □

4.5. Weak convergence and consequences. Recall the definitions of the $(r - 1)$ -dimensional simplex \mathcal{S}_r in Definition 2.6 and the projection map $q: C^r \rightarrow \mathcal{S}_r$ of Equation 1. We will show that for each $r \geq 2$, the set of Perron-Frobenius eigenvectors for the transition matrices of positive automorphisms in $\text{Aut}(F_r)$ is dense in the simplex \mathcal{S}_r .

It is proved in [Sch00, Theorem 21, pg. 5] that Brun’s ordered algorithm is ergodic, conservative, and admits an absolutely continuous invariant measure. The proof uses Rényi’s condition, which further says that the measure is equivalent to Lebesgue measure. We are very much indebted to Jon Chaika for pointing out to us that we could use the ergodicity of Brun’s algorithm to prove the following theorem.

Theorem 4.14. *There exists a set $K \subset \mathcal{S}_r$ of full Lebesgue measure such that for each $x \in K$*

$$(11) \quad \bigcap_{j=1}^{\infty} A_j^x(\mathbb{R}_+^r) = \text{span}_{\mathbb{R}_+} \{x\}.$$

Remark 4.15. Before proceeding with the proof, we explain what we saw as an impediment to proving that the PF eigenvectors are dense in a simplex. It is possible to have a sequence of invertible positive integer $d \times d$ matrices $\{M_i\}_{i=1}^{\infty}$ so that

$$\bigcap_{k=1}^{\infty} M_1 \cdots M_k(\mathbb{R}_+^d)$$

is more than just a single ray. The existence of such sequences of matrices was proved in the context of non-uniquely ergodic interval exchange transformations. There are several papers on the subject (including [KN76], [Kea77], [Vee82], [Mas82]). Because it may not be straightforward to the reader outside of the field, we briefly explain how [Kea77] implies the existence of such a sequence.

We consider a sequence of pairs of positive integers $\{(m_k, n_k)\}_{k=1}^{\infty}$ satisfying the conditions of [Kea77, Theorem 5]. We look at

$$\bigcap_{k=1}^{\infty} A_{m_1, n_1} A_{m_2, n_2} \cdots A_{m_k, n_k}(\mathbb{R}_+^d),$$

as defined on pg. 191. Keane explains on pg. 191 that the product of any two successive A_{m_i, n_i} is strictly positive and that always $\det(A_{m_k, n_k}) = 1$. We let $B_k = A_{m_1, n_1} A_{m_2, n_2} \cdots A_{m_k, n_k}$. [Kea77] projectivizes B_k to \tilde{B}_k , so that \tilde{B}_k is a map of the 3-dimensional simplex \mathcal{S}_4 . By Lemma 4, $\{\tilde{B}_k(0, 1, 0, 0)\}_{k=1}^\infty$ is a sequence of vectors converging to a vector whose 2^{nd} entry is at least $\frac{1}{3}$. By Lemma 3 (when Theorem 5(ii) holds), $\{\tilde{B}_k(0, 0, 1, 0)\}_{k=1}^\infty$ is a sequence of vectors converging to a vector whose 3^{rd} entry is at least $\frac{7}{10}$. But $\frac{1}{3} + \frac{7}{10} > 1$ and we have assumed that we are in \mathcal{S}_4 . So these limits must be distinct vectors.

Proof of Theorem 4.14. Choose any N -cylinder $\Omega := \Delta^r(i_1, \dots, i_N)$ such that the corresponding matrix $Z := A'_N$ is positive (see Corollary 4.13). Let $\bar{\Omega} := p(\Omega)$. Then $\mu(\bar{\Omega}) > 0$ where μ is the Lebesgue measure on B_r . Since $\bar{T}' : B_r \rightarrow B_r$ is ergodic with respect to the Lebesgue measure, by Birkhoff's Theorem, there exists a set $\bar{K} \subset B_r$ such that $\mu(\bar{K}) = 1$ and so that for each $\bar{x} \in \bar{K}$ the following set is infinite:

$$J(\bar{x}) := \{n \in \mathbb{N} \mid \bar{T}'^n(\bar{x}) \in \bar{\Omega}\}.$$

We let $K' := p^{-1}(\bar{K})$. Then for each $x' \in K'$ the set $I(x') := \{n \in \mathbb{N} \mid (T')^n(x') \in \Omega\}$ is infinite, as $n \in I(x')$ if and only if $n \in J(p(x'))$.

Let $K'' := \mathcal{O}^{-1}(K') \subset \mathbb{R}_+^r$. If $x \in K''$, then $\mathcal{O}(x) \in K'$, and hence for each $n \in I(\mathcal{O}(x))$ we have

$$(T')^n(\mathcal{O}(x)) \in \Omega.$$

Let $s \in \mathbb{N}$ be arbitrary. Consider the first s integers $\{j_1, \dots, j_s\}$ in $I(\mathcal{O}(x))$ satisfying that any difference between two of these numbers is $> N$ (where N came from the original N -cylinder we started with). Let $N_1 \geq j_s + N + 1$. Then, for each $m > N_1$,

$$(A_m^{\mathcal{O}(x)})' = D_1 \cdots D_{j_1-1} Z D_{j_1+N} \cdots D_{j_2-1} \cdots D_{j_s-1} Z D_{j_s+N} \cdots D_m,$$

where Z is the positive matrix that we started with and the D_i are the M_i 's of Brun's ordered algorithm. The matrix Z appears in this product s times. By Lemma 4.11, for each $x \in K''$ and each $m \in \mathbb{N}$, there exist permutation matrices P_{i_1}, P_{i_2} so that

$$A_m^x = P_{i_1} (A_m^{\mathcal{O}(x)})' P_{i_2}.$$

Hence, for this arbitrary s we have found an $N_1(s) \in \mathbb{N}$ so that for all $m > N_1$,

$$A_m^x = P_1 D_1 \cdots D_{j_1-1} Z D_{j_1+N} \cdots D_{j_2-1} \cdots D_{j_s-1} Z D_{j_s+N} \cdots D_m P_2,$$

where Z appears in this product at least s times and the other matrices in this product are all invertible nonnegative integer matrices. Then, by [Fis09, Corollary 7.9], Equation 11 holds true for each $x \in K''$.

Let $q : \mathbb{R}_+^r \rightarrow \mathcal{S}_r$ be the projection to the simplex in the positive cone (see (1)) and let μ' be the Lebesgue measure on \mathcal{S}_r . Denote $K = q(K'')$. Then, since $\mu(\bar{K}) = 1$, we arrive at $\mu'(K) = 1$, as desired. \square

Definition 4.16. We let \mathcal{P}_r be defined as:

$$\mathcal{P}_r := \{v_{PF} \in \mathcal{S}_r \mid v_{PF} \text{ is the PF eigenvector for some positive Brun matrix } M \in \mathcal{B}_r\}.$$

Definition 4.17. Suppose

$$(12) \quad A = M_{i_1, j_1} \cdots M_{i_n, j_n} \in \mathcal{B}_r.$$

Then each M_{i_k, j_k} is an unfolding matrix as in (5) and we can associate to it the fold-automorphism f_{i_k, j_k} (see Definition 3.3). Notice that M_{i_k, j_k} is in fact the transition matrix for f_{i_k, j_k} . To each $A \in \mathcal{B}_r$ as in (12) we associate the automorphism

$$(13) \quad g_A = f_{i_n, j_n} \circ \cdots \circ f_{i_1, j_1},$$

whose transition matrix is A . We also call this automorphism g_v , where v is the PF eigenvector of A .

Theorem D. For each $r \geq 2$, let $S_{l_1}^r$ be the set of unit vectors according to the l_1 metric in \mathbb{R}_+^r . The set of Perron-Frobenius eigenvectors of matrices arising as the transition matrices of a positive automorphisms in $\text{Aut}(F_r)$ is dense in the $S_{l_1}^r$.

Proof. It will suffice to show that \mathcal{P}_r is dense. By Theorem 4.14, we know that Brun's algorithm converges in angle on a dense set of points $K \subset \mathcal{S}_r$. Thus, given any $x \in \mathcal{S}_r$ and $\varepsilon > 0$, there exists some $x' \in B(x, \frac{\varepsilon}{2}) \cap K$ on which Brun's algorithm weakly converges. Hence, there exists some N such that, for each $n \geq N$, we have that $q(A_n^{x'} C^r) \subset B(x', \frac{\varepsilon}{2})$. By possibly replacing N with a larger integer, we can further assume that the $A_n^{x'}$ are positive (see Proposition 4.7), so have PF eigenvectors. And the PF eigenvector v_n for each $A_n^{x'}$ is contained in $A_n^{x'} C^r$ and hence is in $B(x', \frac{\varepsilon}{2})$. Hence, there exists a $v_i \in \mathcal{P}_r$ such that $d(v_i, x) < \varepsilon$. \square

5. DENSE GEODESICS IN THETA COMPLEXES

In this section we construct a geodesic ray dense in the theta subcomplex whose top-dimensional simplex has underlying graph as in the right-hand graph of Figure 1. One could consider this a warm-up to the proof of Theorem B or interesting in its own right, as this is the minimal subcomplex containing the projection of the Cayley graph for $\text{Out}(F_r)$.

5.1. Construction of the fold ray. We enumerate the vectors in \mathcal{P}_r from Definition 4.16 as $\{v_i\}_{i=1}^\infty$. For each i there exists a positive matrix A_{v_i} in \mathcal{B}_r so that v_i is the PF eigenvector of A_{v_i} . Further, there exists an automorphism g_{v_i} in \mathcal{A}_r corresponding to A_{v_i} (Equation 13). We also enumerate all possible fold automorphisms, as in (3), by $h_1, \dots, h_{r(r-1)}$.

We construct a sequence that contains each $g_{v_i}^k \circ h_j$ with $i, k, j \in \mathbb{N}$:

$$(14) \quad g_{v_1} \circ h_1, g_{v_2} \circ h_1, g_{v_1}^2 \circ h_2, g_{v_1}^3 \circ h_3, g_{v_2}^2 \circ h_2, g_{v_3} \circ h_1, g_{v_4} \circ h_1 \dots$$

Decompose each g_{v_i} in (14) according to (13), to obtain an infinite sequence of fold automorphisms $\{\Phi_k\}_{k=1}^\infty$ (15). Orienting R_r and identifying its positive edges with the basis, Φ_1 can be represented by the combinatorial fold f_1 . The new graph R_r inherits an orientation and an enumeration of edges and thus we may continue inductively to define the combinatorial fold $f_k: G_{k-1} \rightarrow G_k$. The enumeration of edges induces a homeomorphism $n_k: \hat{\sigma}_{(G_k, f_k \circ \dots \circ f_1)} \rightarrow \mathcal{S}_r$.

Lemma 5.1. *Let σ_0 be the base simplex, then there exists a point $x_0 \in \sigma_0$ so that, for each $k \geq 0$, the fold f_{k+1} is allowable in the folded rose after performing the sequence of folds f_1, \dots, f_k . Moreover, this folded rose is $x_k = n_k^{-1}(T_k \circ \dots \circ T_1(n_0(x_0)))$ in the simplex $\sigma_{(G, f_k \circ \dots \circ f_1 \circ m)}$.*

Proof. This follows from Corollary 3.16. Note that the positivity condition follows from the positivity of the matrices A_{v_i} . \square

Remark 5.2. [Fis09, Corollary 7.9] implies the metric on x_0 is unique, as the same positive matrix occurs in infinitely many of the products $M_{i_1} \dots M_{i_k}$.

Definition 5.3 (\mathcal{R}). We let \mathcal{R} denote the infinite fold ray (Definition 3.13) initiating at x_0 and defined by the sequence of folds $\{f_i\}$ as constructed above.

Theorem C. For each $r \geq 2$, there exists a fold line in \mathcal{T}_r that projects to a Lipschitz geodesic fold ray in $\mathcal{T}_r/\text{Out}(F_r)$.

Proof. We recall \mathcal{R} from Definition 5.3. It is clear that \mathcal{R} is contained in \mathcal{T}_r . \mathcal{R} is a geodesic ray by Corollary 3.18.

First recall that the simplicial and Lipschitz metrics on \mathcal{X}_r induce the same topology on \mathcal{X}_r . Hence, it suffices to prove density in the simplicial metric.

Let $\bar{a} \in \mathcal{M}_r$ and let $\varepsilon > 0$ be arbitrary. Lift \bar{a} to a point $a \in \mathcal{X}_r$ in the interior of a top dimensional simplex τ . Let $y \in \tau$ be a point such that $d_s(a, y) < \varepsilon$, and so that its coordinates are rationally independent. The point y lies on a fold line $\mathcal{F}_{i,j}$ from a point x on one face of τ to a point z in another face. Without generality loss assume $z \in \sigma_0$, the base simplex. Moreover, without generality loss assume the combinatorial fold is $f_{1,2}$. Let e'_1, \dots, e'_r denote the edges of R_r as numbered in σ_0 . Enumerate the edges of G_τ , the underlying graph of τ , as e_1, \dots, e_{r+1} , so that if $c: G \rightarrow R$ is the map collapsing e_2 in G , then $c(e_1) = e'_1$ and $c(e_i) = e'_{i-1}$ for each $i \geq 3$. We parameterize the path $[x, z]$ as an unfolding path $\hat{\gamma}_{1,2}(z, t)$ in unprojectivized Outer Space as follows. Let $\zeta = n_0(z) \in \mathcal{S}_r$, then

$$n_\tau(\hat{\gamma}_{1,2}(z, t)) = (\zeta_1 + t, t, \zeta_2 - t, \zeta_3, \dots, \zeta_r).$$

We note that $z = \hat{\gamma}_{1,2}(z, 0)$, $x = \frac{1}{1-\zeta_2}\hat{\gamma}_{1,2}(z, \zeta_2)$, and for some $0 < t_0 \leq \zeta_2$ we have $y = \frac{1}{1-t_0}\hat{\gamma}_{1,2}(z, t_0)$. Since the function $\gamma_{1,2}(z, t) = \frac{1}{1-t}\hat{\gamma}_{1,2}(z, t)$ is continuous when t is

bounded away from 1, there exists an $\varepsilon' > 0$ so that for each $w \in B(z, \varepsilon')$, the path $\gamma_{1,2}(w, t)$ passes through $B(a, \varepsilon)$.

Now, by the density of PF eigenvectors (Theorem *D*), there exists a vector $v_i \in \mathcal{P}_r$ such that $v_i \in n_0(B(z, \varepsilon'))$. Let $\varepsilon'' > 0$ be such that $n_0(B(z, \varepsilon'))$ contains $B(v_i, \varepsilon'')$. Let K be large enough so that $q(A_{v_i}^k(\mathbb{R}_+^r)) \subset B(v_i, \varepsilon'')$ for all $k > K$.

Let Ψ_k be the composition of Φ_1, Φ_2, \dots from (15) up to the first fold automorphism in the decomposition of $g_{v_i}^k$. Choose $k > K$ so that the last fold automorphism in Ψ_k is $\Phi_{1,2}$. Let z_k be the rose-point in \mathcal{R} directly after performing the fold sequence of Ψ_k . Let x_k be the rose-point directly before z_k .

We claim that $\mathcal{R} \cdot \Psi_k^{-1}$ is ε -close to a . To see this recall that directly after z_k in \mathcal{R} we perform the folds corresponding to $g_{v_i}^k$. Let w_k be the rose-point in \mathcal{R} directly after these folds and let n_s, n_m be the appropriate identifications of the simplices containing z_k and w_k with \mathcal{S}_r . Then $n_s(z_k) = A_{v_i}^k(n_m(w_k))$. Hence $n_s(z_k)$ is ε'' -close to v_i . Thus $n_s(z_k)$ is ε' -close to $n_0(z)$.

The point z_k is in $\sigma_0 \cdot \Psi_k$. Hence, by Lemma 3.9, $n_0(z_k \cdot \Psi_k^{-1}) = n_s(z_k)$. Thus the fold path $[x_k, z_k] \cdot \Psi_k^{-1}$, which is a path induced by the fold $\Phi_{1,2}$, satisfies that its endpoint, $z_k \cdot \Psi_k^{-1}$, is ε' -close to z . Thus, this fold line intersects $B(a, \varepsilon)$, as desired. \square

6. FINDING A ROSE-TO-GRAPH FOLD PATH TERMINATING AT A GIVEN POINT

This section is the first step in our expansion of our methods of Section 5 to obtain a dense geodesic ray in the full quotient of reduced Outer Space.

In this section and in the next one, we find, for a dense set of points y in reduced Outer Space, roses x, z so that $y \in [x, z]$ and the difference in marking map $x \rightarrow z$ is positive. The path $[x, z]$ will be called a positive rose-to-rose fold line. It will replace our basic fold lines $\mathcal{F}_{i,j}$ in the proof of Theorem C.

A rose-to-rose fold path will have two parts: a rose-to-graph part $[x, y]$ and a graph-to-rose part $[y, z]$. We begin in (Subsection 6.1) with decomposing the graph of y into a union of positive loops (Lemma 6.6). This allows us to find the rose x .

6.1. Decomposing a top graph into a union of directed loops.

Definition 6.1 (Paths and distance in trees). Let T be a tree. Then for each pair of points p, q in T there is a unique (up to parametrization) path from p to q . We denote its image by $[p, q]_T$ and, when there is no chance for confusion, we drop the subscript T . Given a tree T , let $d_T(\cdot, \cdot)$ denote the distance in T .

Definition 6.2 (Rooted trees). A *rooted tree* is a tree T with a preferred vertex v_0 called a *root*. A rooted tree can be thought of as a finite set with a partial order that has a minimal element - which is the root. We will refer to the partial ordering induced by the pair (T, v_0) as \leq_T , i.e. $w \leq_T w' \implies w \in [v_0, w']_T$.

Remark 6.3. In the figures to follow the root will always appear at the bottom.

We use special spanning trees to guide us in finding the loop decomposition of G :

Definition 6.4 (Good tree). Let T be a rooted tree in G and $e = (v, w)$ an edge. We call e *bad* if $v \not\prec_T w$ and $w \not\prec_T v$. Let $B(T)$ be the number of bad edges in G with respect to T . When $B(T) = 0$ we call T *good* (sometimes elsewhere called *normal*).

We prove a somewhat stronger version of [Die05, Proposition 1.5.6].

Proposition 6.5. *For each $G \in \mathcal{X}_r$, and for each $E \in E(G)$ that is not a loop, there exists a rooted spanning tree (T, v_0) so that $B(T) = 0$ and $E \in E(T)$ and $v_0 = \text{ter}(E)$. Moreover, when G is trivalent with no separating edges, T can be chosen so that $\text{deg}_T(v_0) = 1$.*

Proof. For an edge $e = (v, w)$ in G , the union $[v_0, v]_T \cup [v_0, w]_T$ forms a tripod. Denote the middle vertex of this tripod by q_e , i.e. q_e satisfies $[v_0, q_e]_T = [v_0, v]_T \cap [v_0, w]_T$. Let $d(e) = d(v_0, q_e)$. We define the complexity $C(T) = (n(T), m(T))$ of T by defining $n(T)$ and $m(T)$ as follows:

$$n(T) = \begin{cases} -\min\{d(e) \mid e \text{ is bad}\} & \text{if } B(T) \neq 0 \\ -\infty & \text{otherwise} \end{cases}$$

and let

$$m(T) = \#\{e \in G \mid e \text{ is bad and } d(e) = -n(T)\}.$$

In particular, $m(T) = 0$ if there are no bad edges, so that, if there are no bad edges, $C(T) = (-\infty, 0)$.

The complexity is ordered by the lexicographical ordering of the pairs. Note that for T bad, the complexity is bounded from below. Indeed, when $G \in \mathcal{X}_r$, we have $|V(G)| \leq 2r - 2$, so $n(T) \geq 2 - 2r$, hence $C(T) \geq (2 - 2r, 1)$. We will show that we can always decrease the complexity, so that T can no longer have a bad edge.

Given an edge E in G , we let $v_0 = \text{ter}(E)$. Let G_1, \dots, G_N denote the components of $G - \{v_0\}$ with $\{v_0\}$ added back to each component separately. Thus $G = \cup_{i=1}^N G_i / \{v_0\}$. We will construct a good tree T_i (rooted at v_0) in each G_i . Then $T = \cup_{i=1}^N T_i$ will also be a good tree rooted at v_0 , since all edges $e \in E(G - T)$ have endpoints inside some G_i .

Let G_1 be the component containing the edge E . For $i \neq 1$ we choose a spanning tree T_i in G_i arbitrarily. For $i = 1$ we construct a spanning tree T_1 in G_1 such that $\text{deg}(v_0)_{T_1} = 1$: Denote by v_1 the endpoint of E distinct from v_0 . We define T_1 to be the union of E with a spanning tree in $G_1 - E$.

We will modify each T_i separately to make it a good tree in G_i . Suppose $e = (v, w)$ is a bad edge in G_i realizing the minimal distance $-n(T_i)$. Let e_1 be the first edge of $[v, q_e]_{T_i}$, and let e_2 be the last edge of $[v, q_e]_{T_i}$. Let $T'_i = T_i \cup \{e\} \setminus \{e_2\}$. We

claim that $e_2 \neq E$. If $G_i \neq G_1$, then this is obvious. Otherwise, $\deg_{T_1}(v_0) = 1$ and $\deg(q_e) \geq 2$, so that $v_0 \neq q_e$ and hence $e_2 \neq E$. Therefore, $E \in E(T'_1)$ after the move and still $\deg_{T'_1}(v_0) = 1$. Next, notice that some bad edges of T_i have become good in T'_i , for example e is no longer bad, as is any edge from a vertex in $[v, q_e]_{T_i}$ to a vertex in $[q_e, w]_{T_i}$. Some bad edges remain bad. But the only edges that were good and became bad are edges with one endpoint in $[v, q_e]_{T_i}$ and one endpoint in a component of $T_i \setminus \{v\}$ that does not contain e_1 or w . For such an edge f , we have that $q_f = v$, so $d(f) = d(v, v_0)_{T'_i} > d(q_e, v_0)_{T'_i} = d(e)$ and the complexity has decreased.

If G is trivalent with no separating edges, then no edge is a loop. This implies that there are three edges E, E', E'' incident at v_0 . If the tripod $E \cup E' \cup E''$, was separating then each of its edges would be separating - a contradiction. Therefore, $G = G_1$ in this case and the proof is complete. \square

Lemma 6.6. *If G is a trivalent graph with no separating edges and $\{E, E'\}$ is a turn at the vertex v_0 , then there exists an orientation on the edges of G so that,*

- (1) $G = \cup_{i=1}^r \alpha_i$ where each α_i is a positive embedded loop,
- (2) $\alpha_i \cap (\cup_{j=1}^{i-1} \alpha_j)$ is a connected arc containing v_0 for each i , and
- (3) v_0 is the terminal point of both E and E' .

To prove Lemma 6.6, we use Proposition 6.5 to find a good tree (T, v_0) containing E and so that v_0 has valence 1 in T . Denote by e_1 the third edge at v_0 distinct from E, E' . Since $E \in E(T)$, we have $E', e_1 \notin E(T)$. Lemma 6.6 now follows from:

Lemma 6.7. *Let G be a trivalent graph with no separating edges and let (T, v_0) be a good spanning tree in G . Let $e_1 \in E(G - T)$ so that $i(e_1) = v_0$. Then one can enumerate $E(G - T) - \{e_1\}$ as e_2, \dots, e_r and orient all of the edges of G so that*

- (1) there exist positive embedded loops $\alpha_1, \dots, \alpha_r \subset G$,
- (2) for each i , we have that i is the smallest index such that α_i contains e_i , and
- (3) for each i , we have that $\alpha_i \cap (\cup_{j=1}^{i-1} \alpha_j)$ is a connected arc containing e_1 .

We will need the following definitions in our proof of Lemma 6.7.

Definition 6.8 ($v_-(e)$ and $v_+(e)$). Let T be a good tree in G . Given an edge $e \in E(G)$, one of its endpoints is closer (in T) to v_0 than the other. We denote by $v_-(e)$ the vertex closer to v_0 and by $v_+(e)$ the vertex further from v_0 .

Definition 6.9. Let α be an embedded oriented path in the graph G and let x, y be two points in the image of α , i.e $x = \alpha(s)$ and $y = \alpha(t)$, for some s, t . If $s < t$, then we denote by $[x, y]_\alpha$ the image of the subpath of α initiating at x and terminating at y , i.e. $\alpha([s, t])$.

Definition 6.10 (Left-right splitting). Let v_0 be a vertex of a graph G and let α be an embedded directed loop based at v_0 . Let e be an edge of α , and m the midpoint

of e . Then the *left-right* splitting of α at e is

$$L_\alpha^e = [v_0, m]_\alpha, \quad R_\alpha^e = [m, v_0]_\alpha.$$

Definition 6.11 (Aligned edges). Let G be a graph and (T, v_0) a spanning good tree in G . Suppose $f_1, f_2 \in E(T)$ satisfy that the vertices $\{v_+(f_1), v_+(f_2), v_0\}$ span a line in T . Then we call f_1, f_2 *aligned*. If $v_+(f_1) < v_+(f_2)$ we say f_1 *lies below* f_2 .

Definition 6.12 (Highlighted subpaths). Let G be a graph and (T, v_0) a spanning good tree in G . Let f_1, f_2 be aligned edges in T such that f_1 lies below f_2 . For each i , let α_{f_i} be an embedded loop containing f_i and v_0 . We define $H_{f_1}^{f_1, f_2}$ and $H_{f_2}^{f_1, f_2}$, the *highlighted* subpaths of α_{f_1} and α_{f_2} , respectively, as follows:

$$(16) \quad H_{f_1}^{f_1, f_2} = L_{\alpha_{f_1}}^{f_1} \quad \text{and} \quad H_{f_2}^{f_1, f_2} = R_{\alpha_{f_2}}^{f_2} \quad \text{when } i(f_1) <_T \text{ter}(f_1)$$

$$(17) \quad H_{f_1}^{f_1, f_2} = R_{\alpha_{f_1}}^{f_1} \quad \text{and} \quad H_{f_2}^{f_1, f_2} = L_{\alpha_{f_2}}^{f_2} \quad \text{when } i(f_1) >_T \text{ter}(f_1)$$

Proof of Lemma 6.7. Enumerate $E(G-T) - \{e_1\}$ so that $k < j$ implies that $v_-(e_k) \not\leq v_-(e_j)$. We prove this lemma by induction on i for $1 \leq i \leq r$. We will define a loop α_i and orient its previously unoriented edges so that items (1), (2), and (3) of the lemma hold and moreover the following items (4) and (5) hold. Denote by α_e the first loop that contains e (for example when $e = e_j \notin E(T)$ then $\alpha_e = \alpha_j$).

- (4) If $f \in E(T)$ and $f' \in E(G)$ are such that $v_+(f) \leq v_+(f')$ and f' is oriented, then f is oriented.
- (5) Let $f_1, f_2 \in E(T)$ be aligned and $H_{f_1}^{f_1, f_2}, H_{f_2}^{f_1, f_2}$ the corresponding highlighted paths with respect to $\alpha_{f_1}, \alpha_{f_2}$, then $H_{f_1}^{f_1, f_2} \cap H_{f_2}^{f_1, f_2}$ contains no half-edges.

We include (5) to ensure that the loop in the induction step is embedded.

The base case: We begin the base of the induction with the edge $e_1 \in E(G-T)$, which is adjacent with v_0 . Define the directed circle $\alpha_{e_1} = e_1 * [t(e_1), v_0]_T$, this is clearly an embedded loop. We orient the edges of α_{e_1} accordingly, i.e. e_1 is directed away from v_0 , and the edges $f \in [t(e_1), v_0]_T$ are directed toward v_0 . Clearly items (1)-(4) hold at this stage, i.e. in the subgraph consisting of precisely α_1 .

The induction hypothesis: We now assume that we have oriented some subset of G so that: for each $j \leq i-1$ we have that e_j is oriented. We call $G' = \cup_{j=1}^{i-1} \alpha_j$ the oriented subgraph. We let $e := e_i$ and note that $e \notin G'$.

The induction step: Let $I = [v_-(e), v_+(e)]_T$. Let t_1 be the edge of I adjacent at $v_-(e)$. We claim as follows that t_1 is oriented, see Figure 2. Indeed, if $v_-(t_1) = v_0$ this follows from the base case. Otherwise, there is an edge $t_3 \in E(T)$ adjacent to $v_-(t_1)$. Since t_3 is nonseparating, there is an edge $e' \in E(G-T)$ so that $v_-(e') \leq v_-(t_3)$ and $v_+(e') \geq v_+(t_3)$. But since the graph is trivalent, $v_+(e') \geq v_+(t_1)$. Now, since $v_-(e') < v_-(e)$, we have that e' is oriented. Hence, by item (4) in the induction hypothesis, t_1 is oriented.




FIGURE 2. The green line indicates the path α_e takes near e . The edge t_1 is the first edge in I and t_2 is the last oriented edge in I .

Let t_2 be the last oriented edge of I . Denote $J = [v_+(t_2), v_+(e)]_T$. Note that we allow $t_1 = t_2$ and leave the adjustments of this case, from cases A and C of Figure 2, to the reader. The loop α_e is constructed as in Figure 2 from the following segments by adding or removing a half-edge of t_2 or removing a half-edge of t_1 or t_2

$$(18) \quad H_{t_2}^{t_1, t_2}, J, e, H_{t_1}^{t_2, t_1}.$$

The orientation of α_e is chosen according to the direction of t_1 . When $i(t_1) >_T t(t_1)$, cases (A) and (B) of Figure 2, we orient α_e so that $H_{t_2}^{t_1, t_2}$ is the left (first) segment and when $i(t_1) <_T t(t_1)$, cases (C) and (D), we orient so that $H_{t_1}^{t_2, t_1}$ is the first segment.

We observe that α_e is indeed embedded as follows. The paths $H_{t_2}^{t_1, t_2}, H_{t_1}^{t_2, t_1} \subset G'$ while $J, e \subset G - G'$, hence they are edge-disjoint. Moreover, by item (5) for G' , the paths $H_{t_2}^{t_1, t_2}, H_{t_1}^{t_2, t_1}$ are half-edge disjoint so that even if we add t_1 or t_2 they remain edge disjoint. Therefore the path α_e does not self-intersect in an edge. It cannot self-intersect at a vertex since the graph is trivalent. This proves item (1).

Note that α_e is contained in $G' \cup T \cup \{e\}$, hence it does not contain e_j for $j > i$. This proves item (2). Item (3) is also clear. Item (4) is satisfied since for $f' = e$ and for all $f \in E(T)$ so that $v_+(f) \leq v_+(e)$ we have that f is oriented.

We are left with proving item (5) for each pair of aligned edges $f_1, f_2 \in E(T)$ such that at least one of them is oriented in the i^{th} step, i.e. $\alpha_f = \alpha_i$. It is less difficult to check that the claim holds when both edges are newly oriented. We leave this case to the reader and check the case where $f_1 \subset G' \cap T$ and $f_2 \subset T$ is newly oriented, i.e. $f_2 \subset J$. We illustrate the different cases in Figure 3.

Suppose $f_1 \notin I$ (Case I in Figure 3), then for each $f \in \{f_2, t_1, t_2\}$, we have $f_1 <_T f$. Thus, $H_{f_1}^{f_1, f} = H_{f_1}^{f_1, f'}$ for $f, f' \in \{f_2, t_1, t_2\}$.




FIGURE 3. Checking Claim 5 for f_1, f_2 depending on their location. In Case I, f_1 lies below t_1 , and in the other cases t_1 lies below f_1 . In Cases IIa, IIb, f_1 is pointing down and in Case III, f_1 is pointing up.

Moreover, by our construction of $\alpha_{f_2} = \alpha_e$, we have $H_{f_2}^{f_1, f_2} \subset H_{t_k}^{f_1, t_k} \cup J \cup \{t_2\}$ for either $k = 1$ or $k = 2$. Thus, for the same k we have (see Figure 2) :

$$H_{f_1}^{f_1, f_2} = H_{f_1}^{f_1, t_k} \quad \text{and} \quad H_{f_2}^{f_1, f_2} \subset H_{t_k}^{f_1, t_k} \cup \{e\} \cup J \cup \{t_2\}.$$

Since J is newly oriented, $H_{f_1}^{f_1, t_k}$ and J are edge-disjoint. By the induction hypothesis, $H_{f_1}^{f_1, t_k} \cap H_{t_k}^{f_1, t_k}$ contains no edges or half-edges. Moreover, if $H_{f_2}^{f_1, f_2} \not\subset H_{t_k}^{f_1, t_k} \cup J \cup \{e\}$ then $k = 2$ and a half of t_2 must be in $H_{t_2}^{f_1, t_2}$. Since $H_{f_1}^{f_1, t_k} \cap H_{t_k}^{f_1, t_k}$ contains no half-edges, then t_2 is not contained in $H_{f_1}^{f_1, t_k}$. This implies that $H_{f_1}^{f_1, f_2} \cap H_{f_2}^{f_1, f_2}$.

If $f_1 \in I$, then there are two classes of cases: $i(f_1) \geq_T \text{ter}(f_1)$ (Cases IIa, IIb of Figure 3) and $i(f_1) \leq_T \text{ter}(f_1)$ (Case III). We will prove Case IIa and leave the others to the reader. If f_1 is pointing down, then $H_{f_1}^{f_1, f_2} = R_{\alpha_{f_1}}^{f_1}$, $H_{f_2}^{f_1, f_2} = L_{\alpha_{f_2}}^{f_2}$. There are two subcases: f_2 is pointing down (Case IIa) or up (Case IIb). If f_2 is pointing down, then t_1 is pointing up (see Figure 2), thus $L_{\alpha_e}^{f_2} \subset L_{\alpha_{t_1}}^{t_1} \cup e \cup J$. In this configuration,

$$\begin{aligned} H_{f_1}^{f_1, t_1} &= R_{\alpha_{f_1}}^{f_1} & H_{t_1}^{f_1, t_1} &= L_{\alpha_{t_1}}^{t_1} \\ H_{f_1}^{f_1, f_2} &= R_{\alpha_{f_1}}^{f_1} & H_{f_2}^{f_1, f_2} &= L_{\alpha_e}^{f_2} \subset L_{\alpha_{t_1}}^{t_1} \cup \{e\} \cup J \end{aligned}$$

Hence, the fact that $H_{f_1}^{f_1, t_1} \cap H_{t_1}^{f_1, t_1}$ contains no half-edges implies that $H_{f_1}^{f_1, f_2} \cap H_{f_2}^{f_1, f_2}$ contains no half-edges. The other cases are similar. \square

6.2. Rose-to-graph fold line. Given a point x whose underlying graph is trivalent with no separating edges, we wish to find a rose-point x_0 and a line in \mathcal{X}_r from x_0 to x . This is done by simultaneous folding as defined below.

Definition 6.13 (Rose-to-graph fold line $\mathcal{F}(x_0, \{s_{ij}\})$). Let $x_0 = (R, \mu, \ell_0)$ be a point in $\hat{\mathcal{X}}_r$ whose underlying graph is a rose with r petals. There are $K = r(r-1)$ turns and we enumerate them in any way $\{\tau_i\}_{i=1}^K$. Let $\vec{s} \in \mathbb{R}^K$ be a nonnegative vector so that s_i is no greater than the length of each edge in the turn τ_i . Given the data (x_0, \vec{s}) we construct a continuous family of graphs $\{x_t\}$ for $0 \leq t \leq T = \sum s_i$, and maps $f_{t,0}: x_0 \rightarrow x_t$ as follows. In the i^{th} step let $\tau_i = \{e_j, e_m\}$ and fold initial segments of length s_i in $f_{t,0}(e_j)$ and $f_{t,0}(e_m)$. We caution that these $f_{t,0}$ are not always homotopy equivalences. However, if $f_{T,0}$ is a homotopy equivalence, then for each $t < T$ the map $f_{t,0}$ is a homotopy equivalence. In this case we get a path

$$\begin{aligned} \mathcal{F}(x_0, \vec{s}): [0, T] &\rightarrow \hat{\mathcal{X}}_r \\ \mathcal{F}(x_0, \vec{s})(t) &= x_t. \end{aligned}$$

We denote its projectivization by $\bar{\mathcal{F}} = q(\mathcal{F})$.

Lemma 6.14. *Let G be a trivalent graph such that $\pi_1(G) \cong F_r$. Then for each $x \in \hat{\mathcal{X}}_r$ with underlying graph G , there exists a point x_0 whose underlying graph is a rose and there exists a nonnegative vector \vec{s} in \mathbb{R}^K , where $K = r(r-1)$, so that*

$$x = \mathcal{F}(x_0, \vec{s})(T).$$

Additionally, x_0, \vec{s} are linear functions of the lengths of $E(G)$ as they vary throughout the unprojectivized simplex. (As above, $T = \sum s_i$.)

Proof. Let E_1, \dots, E_{3r-3} be the edges in G . Lemma 6.6 provides a decomposition of G as $G = \cup_{i=1}^r \alpha_i$. Let $R = \sqcup_{i=1}^r \alpha_i / \{v_0\}$, then R is an r -petaled rose. Let e_1, \dots, e_r be the edges of R . Let $\ell_R(e_i) = \ell_G(\alpha_i)$. We write α_i as a sequence of edges $\alpha_i = E_{m(i,1)} \cdots E_{m(i,k_i)}$. Then

$$(19) \quad \ell_R(e_i) = \ell_G(\alpha_i) = \sum_{j=1}^{k_i} \ell_G(E_{m(i,j)}).$$

Let x_0 denote R with these edge lengths. There is a natural map $\theta: x_0 \rightarrow x$ defined by the inclusion of α_i in G . This is a quotient map and, moreover, $\theta|_{e_i}$ is an isometry.

Recall that the intersection of α_i and α_j is an arc containing v_0 . Let a, b be the endpoints of the arc. For $\tau_k = (e_i, e_j)$ and $\tau_m = (\bar{e}_i, \bar{e}_j)$ we define

$$(20) \quad s_k = l_G([v_0, a]_{\alpha_i}) \quad \text{and} \quad s_m = l_G([b, v_0]_{\alpha_i}).$$

Consider the folding line $\mathcal{F}(x_0, \vec{s})$. By the definitions, θ is precisely the map $f_{T,0}: x_0 \rightarrow x_T$, since the points identified by θ are precisely those that are identified in the folds. Therefore, x_T equals the point x that we started with.

Moreover, θ is a homotopy equivalence by Lemma 6.7(2). Therefore, $\mathcal{F}(x_0, \vec{s})$ is a path in unprojectivized Outer Space. Equations 19, 20 show that the dependence of l_i and s_k on edge lengths in x is linear. \square

Lemma 6.15. *Suppose that $x = \mathcal{F}(x_0, \vec{s})(T)$. Then there exists a neighborhood of (x_0, \vec{s}) so that for each (y_0, \vec{u}) in this neighborhood, the endpoint $y := \mathcal{F}(y_0, \vec{u})(T')$ of the fold line $\mathcal{F}(y_0, \vec{u})$ lies in the same unprojectivized open simplex in $\hat{\mathcal{X}}_r$ as x .*

Additionally, the edge lengths of y are linear combinations of the edge lengths of y_0 and \vec{u} .

Proof. Consider the positive edges E_1, \dots, E_m in G and let $G = \cup_{i=1}^r \alpha_i$ be the decomposition guaranteed by Lemma 6.6. The edge E_i is contained in a loop $\alpha_{j(i)}$. Since G is a trivalent graph, at each endpoint $\{v, w\}$ of E_i there is an edge, E_k, E_d resp., not contained in α_i . The edges E_k, E_d are contained in $\alpha_{j(k)}, \alpha_{j(d)}$ resp. Now $v_0, v \in \alpha_{j(i)} \cap \alpha_{j(k)}$. Thus, by Lemma 6.6(3), either $[v_0, v]_{\alpha_{j(i)}} \subseteq \alpha_{j(i)} \cap \alpha_{j(k)}$ or $[v, v_0]_{\alpha_{j(i)}} \subseteq \alpha_{j(i)} \cap \alpha_{j(k)}$. This situation is similar for v_0, w . Therefore,

$$(21) \quad \ell_G(E_i) = \begin{cases} s_m - s_n & \text{when } \tau_m = (e_{j(d)}, e_{j(i)}) \text{ and } \tau_n = (e_{j(k)}, e_{j(i)}) \\ s_n - s_m & \text{when } \tau_m = (\overline{e_{j(d)}}, \overline{e_{j(i)}}) \text{ and } \tau_n = (\overline{e_{j(k)}}, \overline{e_{j(i)}}) \\ |\ell(\alpha_{j(i)}) - (s_n + s_m)| & \text{otherwise} \end{cases}$$

Note this dependence of the $\ell_G(E_i)$ on the variables s_m will be the same for all points in the same unprojectivized simplex since they only depend on the loop decomposition. We also get that the dependence of $\ell_G(E_i)$ on s_{ij} and $\ell(e_i)$ is linear. Let U be the open subset of the unprojectivized rose simplex cross $\mathbb{R}_+^{r(r-1)}$ so that each expression in the right-hand side of (21) is positive for each i . This is an open set containing (x_0, \vec{s}) . For any point y in the unprojectivized simplex of x one can use (19) and (20) to get y_0 and \vec{u} so that $y = \mathcal{F}(y_0, \vec{u})$. Thus (y_0, \vec{u}) solves the equations in (21) for the edge lengths of y , hence $(y_0, \vec{u}) \in U$. Thus, there exists a solution of (21) in U for each choice of edge lengths of G . We see that the dependence of the edge lengths $l_G(E_i)$ on \vec{s} and $\ell(e_i)$ is linear. \square

6.3. Folding a transitive graph to a rose.

Definition 6.16. A *transitive* graph G is a directed graph G with the following property: for any two vertices w, w' there exists a directed path from w to w' .

Note that it is enough to check that for any choice of preferred vertex v , there exists a directed path to and from each other vertex v' .

The proof of the following is left to the reader.

Observation 6.17. Let G be a directed graph and let $f: G \rightarrow G'$ be a direction matching fold of two oriented edges e_1, e_2 in G , starting at a common vertex v . Then:

- (1) If G is transitive, then G' is transitive.
- (2) If the lengths of the edges of G are rationally independent, then the lengths of the edges of G' are rationally independent.

Lemma 6.18. *Let G be any transitive graph with rationally independent edge lengths and let $\{E, E'\}$ be either a positive or negative turn. Then there exists a fold sequence f_1, \dots, f_k containing only direction matching folds and satisfying that $f_k \circ \dots \circ f_1(G)$ is a rose. Further, assuming G is trivalent, we may choose f_1 so that it folds the turn $\{E, E'\}$.*

Proof. We perform the following steps:

Step 1: Let $c(G)$ be the number of directed *embedded* paths in G between (distinct) vertices. For each pair of vertices w, w' there exists a directed path from w to w' , thus it follows that there exists an embedded directed path from w to w' . We decrease $c(G)$ by folding two directed embedded paths α, β so that $i(\alpha) = i(\beta)$ and $ter(\alpha) = ter(\beta)$ and $\alpha \cap \beta = \{i(\alpha), ter(\alpha)\}$. Note that the edges in α and β are distinct and thus we are consecutively performing regular Stallings folds. Also, if G is trivalent and we choose a decomposition as in Lemma 6.6, then we can choose the first α to contain E and the first β to contain E' and fold α, β so that the first combinatorial fold folds the turn $\{E, E'\}$. Denote the new graph by G' . Then, by Observation 6.17, G' is transitive and has rationally independent edge lengths. The complexity has decreased, i.e. $c(G') < c(G)$.

At the end of this step we may assume that we have a connected graph G so that $G = \coprod_{i=1}^n \gamma_i / \sim$, where γ_i is a circle and for each $i \neq j$: $\gamma_i \cap \gamma_j$ is either empty or a single point (see Figure 4). We call such a graph a *gear graph*. Notice that given a gear graph, such a decomposition into circles is unique up to reindexing




FIGURE 4. This graph is an example of a gear graph.

Step 2: For a gear graph, we define a new complexity. Let V be the set of vertices (of valence > 2) and let $m = \max\{\text{val}(w) \mid v_0 \neq w \in V\}$. Define the complexity:

$$c(G) = (|V|, m).$$

Let w be a vertex that realizes the valence m and let γ_1, γ_2 be circles so that $w = \gamma_1 \cap \gamma_2$ and suppose that $v \neq w$ is a vertex on γ_2 (see Figure 4). Let $\gamma_2 = \alpha\beta$ where $i(\alpha) = v = ter(\beta)$ and $ter(\alpha) = w = i(\beta)$. By folding, wrap β over γ_1 until v is on the image of γ_1 (we may have to wrap β multiple times over γ_1). Now there

are two paths from v to w : α and γ'_1 , the remaining part of γ_1 . Fold $\bar{\alpha}$ over $\overline{\gamma'_1}$. G' is a gear graph. Moreover, the valence of the image of w decreases (w may now even have valence 2). Thus $c(G') < c(G)$. When all vertices other than v_0 have valence 2 we can no longer decrease the complexity, and G will be a rose. \square




FIGURE 5. Step 2 (β' is the remaining portion of β , after it is wrapped around γ_1)

7. EXISTENCE AND CONTINUITY OF ROSE-TO-ROSE FOLD LINES

In Subsection 6.2 we defined a rose-to-graph fold line $[x_0, x] = \mathcal{F}(x_0, \vec{s})$, given a point x and a loop decomposition of its underlying graph. We also had two continuity statements: (1) x_0, \vec{s} vary continuously as a function of x by Lemma 6.14 and (2) $\mathcal{F}(x_0, \vec{s})$ vary continuously as a function of x_0, \vec{s} by Lemma 6.15. We need similar existence and continuity statements for the full rose-to-rose fold line.

Proposition 7.1. *Let x be any point of $\hat{\mathcal{X}}_r$ satisfying that $q(x)$ is in a top-dimensional simplex of reduced Outer Space with rationally independent edge lengths. Let $\{E, E'\}$ be a pair of adjacent edges in the underlying graph of x . Then there exists a positive rose-to-rose fold line, which we denote by $\mathcal{R}(x_0, \vec{s}): [0, L] \rightarrow \mathcal{X}_r$, containing x and containing the fold of $\{E, E'\}$.*

Proof. By Lemma 6.14 there exists a rose point x_0 and a vector \vec{s} so that the rose-to-graph fold segment $\mathcal{F}(x_0, \vec{s})$ terminates at the point x . We consider the orientation on x given by the loop decomposition in Lemma 6.6. We may apply Lemma 6.18 to obtain a fold sequence f_1, \dots, f_k which terminates in a rose z' with some valence-2 vertices and so that f_1 folds the turn $\{E, E'\}$. Removing the valence-2 vertices gives a rose, which we denote by z . The line just described will be denoted $\mathcal{R}(x_0, \vec{s})$. It satisfies the statement in the theorem. \square

Notation: Let x_0 be a rose, let $\vec{s} \in \mathbb{R}^{r(r-1)}$, and let $\mathcal{R}(x_0, \vec{s})$ be the fold line defined by these parameters. (We call such a fold line a *rose-to-rose* fold line.) We will denote by x the trivalent metric graph at the end of the rose-to-graph fold segment, i.e. $x = \mathcal{R}(x_0, \vec{s})(\sum s_i)$, by z the end-point of the rose-to-rose fold segment, and by z' the point in the graph-to-rose fold segment from which z is obtained by removing the valence-2 vertices.

Definition 7.2 (Proper fold line). Let $\mathcal{R}(x_0, \vec{s})$ be a rose-to-rose fold line and let f_1, \dots, f_k be the sequence of combinatorial folds from x to z' . If for each l the fold f_l is a proper full fold, then we will say that $\mathcal{R}(x_0, \vec{s})$ is a *proper fold line*.

For example, for each x as in Proposition 7.1, the fold line constructed in the proposition is a proper fold line since the edge lengths in x are rationally independent.

Proposition 7.3. *For each proper rose-to-rose fold line $\mathcal{R}(x_0, \vec{s})$ and $\varepsilon > 0$, there exists a neighborhood U of (x_0, \vec{s}) so that for any point $(y_0, \vec{u}) \in U$:*

- (1) *The endpoints of the rose-to-graph fold segments*

$$y := \mathcal{F}(y_0, \vec{u})(T'), \quad x := \mathcal{F}(x_0, \vec{s})(T)$$

lie in the same unprojectivized open simplex and are ε -close.

- (2) *The sequence of combinatorial folds from x to z' appearing in the graph-to-rose fold segment is allowable in y .*
- (3) *Let $\mathcal{R}(y_0, \vec{u})$ be the fold line defined by concatenating $\mathcal{F}(y_0, \vec{u})$ with the fold segments from (2), then the terminal points $w := \mathcal{R}(y_0, \vec{u})(L')$ and $z := \mathcal{R}(x_0, \vec{s})(L)$ are ε -close.*

Proof. By Lemma 6.15 there exists a neighborhood U of (x_0, \vec{s}) so that for each $(y_0, \vec{u}) \in U$ the fold line $\mathcal{F}(y_0, \vec{u})$ terminates at some point y lying in the same unprojectivized top-dimensional simplex as x . Since the edge lengths of y vary continuously with the edge lengths of y_0 and \vec{s} , we can make U smaller, if necessary, to ensure that y and x are ε -close. This proves (1).

Let f_1, \dots, f_k be the combinatorial fold sequence from x to z' , as described in Lemma 6.18. For each combinatorial proper full fold folding e_i over e_j , there corresponds a square $m \times m$ folding matrix (here $m > r$), which we denote $T'_{ij} = (a_{kl})$, so that $a_{kl} = 1$ for $k = l$ and $a_{ij} = -1$, but otherwise $a_{kl} = 0$ when $k \neq l$ (compare with Definition 3.4). Let T'_1, \dots, T'_k be the fold matrices corresponding to f_1, \dots, f_k . Then just as in Lemma 3.14, the combinatorial fold sequence f_1, \dots, f_k is allowable in the point y if and only if for each l the vector $T'_1 \cdots T'_k(\ell(y))$ is positive for each $1 \leq l \leq k$. This defines an open neighborhood of x where item (2) holds. Item (3) follows from the fact that matrix multiplication is continuous. \square

Lemma 7.4. *Let $h: x_0 \rightarrow z$ be a homotopy equivalence representing the map from the initial rose x_0 to the terminal rose z in a rose-to-rose fold line. Let H be a matrix representing the change-of-metric from the rose z to the rose x_0 . Then H is a nonnegative invertible integer matrix and it equals the transition matrix of h .*

Proof. Let x_0 be the initial rose with edges e_1, \dots, e_r and let z be the terminal rose with edges e'_1, \dots, e'_r . Let z' be the graph on the fold line just before z , i.e. z is obtained from z' by unsubdividing at all of the valence-2 vertices. Let $h: x_0 \rightarrow z$

and $g: x_0 \rightarrow z'$ be the relevant homotopy equivalences. Since g is a subdivision followed by folding maps, each of which is a positive map, we have for each i that $g|_{e_i}$ is a local isometry. Thus, $h|_{e_i}$ is a local isometry. Moreover, h maps the unique vertex of x_0 to the unique vertex of z . Suppose $h(e_i)$ contains a part of an edge e'_i , then $h(e_i)$ contains a full appearance of e'_i (since there is no backtracking and the vertex maps to the vertex). Therefore, $h|_{e_i}$ is an edge-path in z . Thus, we may write

$$(22) \quad l(e_i) = \sum_j m(i, j) l(e'_j),$$

where the $m(i, j)$ are the nonnegative integer entries of the transition matrix of h . By Equation 22, the change-of-metric matrix from z to x_0 coincides with the transition matrix of the homotopy equivalence h . Therefore, H is nonnegative, and integer. Moreover, since all the folds in a rose-to-rose fold line are direction matching, h is a positive map. Thus, H is equal to $Ab(h)$, the map induced by h (viewed as an automorphism) by abelianization. Therefore, H is invertible. \square

Lemma 7.5. *For each $\varepsilon > 0$ and proper fold line $\mathcal{R}(x_0, \vec{s})$, there exists a neighborhood U of the terminal rose z such that, for each $w \in U$, there exists a proper rose-to-rose fold line $\mathcal{R}(y_0, \vec{u})$ terminating at w satisfying that*

- (1) *the top graphs x, y are ε -close,*
- (2) *the combinatorial fold sequence corresponding to the graph-to-rose segments are the same in both lines, and*
- (3) *the change-of-metric matrix for both fold lines is the same.*

Proof. We prove (1) and (2). By Lemma 7.4, the change-of-metric matrix H from z to x_0 is nonnegative. Thus, for any w in the same unprojectivized simplex as z , we have that $Hl(w)$ is also positive. By Proposition 7.3, there exists a neighborhood V of x_0 so that for each $y_0 \in V$ there exists a vector \vec{u} so that if y is the top graph of the fold line $\mathcal{F}(y_0, \vec{u})$, then x and y are ε -close and their combinatorial fold sequences are the same. The neighborhood U can be taken in $H^{-1}(V)$. This proves (1) and (2). Since the combinatorial folds are the same, the transition matrix $y_0 \rightarrow w$ is the same as the transition matrix $x_0 \rightarrow z$. (3) follows from Lemma 7.4. \square

Definition 7.6 (Rational rose-to-rose fold lines). A rose-to-rose fold line $\mathcal{R}(x_0, \vec{s})$ is called *rational* if for each edge e in G and for each loop α_i in the loop decomposition of the underlying graph of x , the quotient $\frac{l(e, x)}{l(\alpha_i, x)}$ is rational.

Proposition 7.7. *For each $\varepsilon > 0$ and for each x in an unprojectivized top-dimensional simplex of reduced Outer Space, with rationally independent edge-lengths, there exists a rational proper rose-to-rose fold line passing through some x' in the same open unprojectivized simplex as x , satisfying that $d(x', x) < \varepsilon$.*

Proof. Let x be a point in an unprojectivized top-dimensional simplex and having rationally independent edges. Let $\mathcal{R}(x_0, \vec{s})$ be a proper rose-to-rose fold line containing x . Let V be a neighborhood of (x_0, \vec{s}) guaranteed by Proposition 7.3. Let U be a neighborhood of x in the same top-dimensional simplex and such that for each $y \in U$ there exists some $(y_0, \vec{u}) \in V$ with y the terminal point of $\mathcal{F}(y_0, \vec{u})$. This is possible by Lemma 6.14. Let x' be a point in U which is ε -close to x and such that the ratios $\frac{l(e', x')}{l(\alpha_i, x')}$ are rational. Then the resulting fold line through x' will have the required properties. \square

8. CONSTRUCTING THE FOLD RAY

Enumerate \mathcal{P}_r by $\{v_i\}_{i=1}^\infty$ (see §4). For each i there are countably many rational proper rose-to-rose fold lines. Thus there are countably many rational proper rose-to-rose fold lines terminating at a rose with length-vector v_i . For each such fold line \mathcal{R}_{ij} , let F_{ij} denote its fold matrix and H_{ij} its inverse - an invertible nonnegative matrix. Let U_{ij} denote the neighborhood of v_i from Lemma 7.5 (for $\varepsilon = 1$ will suffice), i.e. for each $w \in U_{ij}$ there exists a proper fold line terminating at w , passing through the same simplices as \mathcal{R}_{ij} and satisfying that their fold matrices are the same.

Since each $A_i = A_{v_i}$ is a positive matrix, for each i, j there exists an integer $n(i, j)$ satisfying that for each $n > n(i, j)$ we have $A_i^n(\mathbb{R}_+^r) \subset U_{ij}$.

Recall that g_{v_i} has a decomposition into fold automorphisms obtained from Brun's algorithm, this induces a decomposition of A_i to unfolding matrices and A_i^{-1} into folding matrices. We then create a sequence of pairs denoted $\{a_k\}$ which satisfies:

- (1) If $a_k = (i, j)$ for an odd k then there exists some n such that $n > n(i, j)$ so that $a_{k+1} = (i, n)$.
- (2) For each $i, N \in \mathbb{N}$ there exists an $n > N$ and an even k so that $(i, n) = a_k$,
- (3) For each $i, j \in \mathbb{N}$, there exist infinitely many odd k 's so that $a_k = (i, j)$,

To each a_k in this sequence we attach an (unfolding) matrix or sequence of matrices and automorphisms: if k is odd and $a_k = (i, j)$ we attach the matrix H_{ij} related to the rose-to-rose fold line \mathcal{R}_{ij} and define f_k to be its change-of-marking automorphism. And if k is even and $a_k = (i, n)$ we attach a sequence of unfolding matrices according to a Brun's algorithm decomposition $A_i^n = (M_1^i \dots M_k^i)^n$ and a sequence of fold automorphisms. We get a sequence of matrices, which we denote by $\{D_l\}_{l=1}^\infty$. For each l , either $D_l = H_{ij}$ or $D_l = M_j^i$ for some i, j . We emphasize that we have not decomposed H_{ij} . Moreover we get a sequence of automorphisms $\{f_l\}_{l=1}^\infty$. Let $\{Z_l\}_{l=1}^\infty$ denote the sequence of folding matrices, i.e. $Z_l = D_l^{-1}$ for each l .

Definition 8.1 (Ray \mathfrak{R}). We construct as follows the geodesic fold ray \mathfrak{R} that we later prove is dense. Let x_0 be a rose in the unprojectivized base simplex (i.e. having the identity marking) with the length vector w_0 provided by Lemma 3.15 for the

matrix sequence $\{D_l\}$ that we defined in the paragraph above. For each k let x_k be the rose with length vector inductively defined as $w_k = Z_k w_{k-1}$ and with the marking $f_k \circ \cdots \circ f_1$. For each l , if Z_l is a single fold matrix coming from the matrix decomposition of A_i^n , then let the line from x_l to x_{l+1} be the single proper full fold line corresponding to the matrix Z_l . This fold is allowable since $Z_l w_l = w_{l+1}$ is positive. If $Z_l = H_{i,j}$ for some i, j coming from $a_k = (i, j)$, then $a_{k+1} = (i, n)$ for $n > n(i, j)$. Hence there is a number s so that the following s matrices $\{D_d\}_{d=l+1}^{l+s}$ are the matrices of the decomposition of A_i^n . Therefore, $w_{l+1} = A_i^n(w_{l+s})$ for $n > n(i, j)$. Thus $x_{k+1} \in U_{ij}$, so the fold line corresponding to \mathcal{R}_{ij} is allowable in $x_k = H_{ij}(x_{k+1})$. We insert this fold line between such x_k and x_{k+1} . This defines a fold ray connecting the x_k 's.

Theorem B. For each $r \geq 2$, there exists a geodesic ray $\tilde{\gamma}: [0, \infty) \rightarrow \mathcal{R}\mathcal{X}_r$ so that the projection of $\tilde{\gamma}$ to $\mathcal{UR}\mathcal{X}_r/\text{Out}(F_r)$ is dense.

Proof. We will show that, for each $r \geq 2$, the fold ray \mathfrak{R} of Definition 8.1 is contained in $\mathcal{UR}\mathcal{X}_r$ and projects densely into \mathcal{U}_r . The ray is geodesic by Corollary 3.18.

To prove that \mathfrak{R} never leaves $\mathcal{R}\mathcal{X}_r$, i.e. contains no graph with a separating edge, it will suffice to show that at each point $x \in \mathfrak{R}$, the underlying graph can be directed so that it is a transitive graph. This clearly holds for each proper full fold of a rose, hence for the fold sequences coming from the decompositions of the $g_{v_i}^k$. Moreover, for each i, j \mathcal{R}_{ij} consists of transitive graphs since all folds are direction matching, see Observation 6.17(2).

Let x be a point in an unprojectivized top-dimensional simplex with rationally independent edge lengths. Let G be its underlying graph and $\{E, E'\}$ a turn. By Theorem 7.1 we can construct a proper positive rose-to-rose fold line $\mathcal{R} = \mathcal{R}(x_0, \vec{s})$ containing x and the combinatorial fold f_1 of $\{E, E'\}$ directly after x . We may assume that the terminal point of this rose-to-rose fold line z lies in σ_0 . For each $\varepsilon > 0$, by Lemma 7.7, there exists a proper *rational* rose-to-rose fold line \mathcal{R}' containing a point x' in the same unprojectivized open simplex as x , so that x, x' are ε -close, and so that the fold f_1 is the fold following x' . Let H be the unfolding matrix corresponding to \mathcal{R}' .

For each $\varepsilon > 0$, there exists, as in Lemma 7.4, an open neighborhood U of the terminal point of \mathcal{R}' so that for each $w \in U$, there exists a proper rose-to-rose fold line $\mathcal{R}'(y_0, \vec{u})$, terminating at w , such that the top graphs x', y are ε -close, the combinatorial fold sequence in the graph-to-rose segments are the same, and the change-of-metric matrix for $\mathcal{R}'(y_0, \vec{u})$ is H . Since the set of PF eigenvectors is dense, there exists an i so that the PF eigenvector $v_i \in U$. Hence, there exists a rose-to-rose fold line R_{ij} passing through the same unprojectivized simplices as \mathcal{R}' and having the same change-of-metric matrix $H_{ij} = H$. Moreover, there exists a point $x''_{ij} \in R_{ij}$ in the same unprojectivized top-dimensional simplex as x' and ε -close to x' . By

Definition 8.1, there exist infinitely many k 's so that the fold line between x_k and x_{k+1} is the one passing through the same unprojectivized simplices as R_{ij} (hence \mathcal{R}'). In fact, these occur before arbitrarily high powers of g_{v_i} , so that they terminate arbitrarily close to a rose with length vector v_i . Let k be such a number and let Ψ_k be the composition of the automorphisms f_1, f_2, \dots up to x_k . Thus, by Lemma 7.4, there exists a point $\xi \in [x_{k-1}, x_k] \cdot \Psi_k^{-1}$ in the same unprojectivized top-dimensional simplex as x''_{ij} and ε -close to x''_{ij} . Hence, ξ is the point on the ray defined in Definition 8.1 that is 3ε -close to our original point x and the fold immediately after ξ is the one folding the turn $\{E, E'\}$. \square

Theorem A. For each $r \geq 2$, there exists a geodesic fold ray in the reduced Outer Space $\mathcal{R}\mathcal{X}_r$ whose projection to $\mathcal{R}\mathcal{X}_r/\text{Out}(F_r)$ is dense.

Proof. This is an immediate corollary of Theorem B. \square

9. APPENDIX: LIMITS OF FOLD GEODESICS.

In many cases, as in the case of the geodesic that we construct in this paper, a concatenation of fold segments $\{\gamma_i: [i, i+1] \rightarrow \hat{\mathcal{X}}_r\}_{i=1}^\infty$ that glue together to a ray $\gamma: [0, \infty) \rightarrow \hat{\mathcal{X}}_r$, projecting under q to a Lipschitz geodesic, satisfies the properties of a semi-flow line below.

Definition 9.1 (Semi-flow line). ([HM11, pg. 3], definition of a “fold line”) A semi-flow line in unprojectivized Outer Space is a continuous, injective, proper function $\mathbb{R} \rightarrow \hat{\mathcal{X}}_r$ defined by a continuous 1-parameter family of marked graphs $t \rightarrow G_t$ for which there exists a family of homotopy equivalences $h_{ts}: G_s \rightarrow G_t$ defined for $s \leq t \in \mathbb{R}$, each of which preserves marking, such that the following hold:

- (1) Train track property: For all $s \leq t \in \mathbb{R}$, the restriction of h_{ts} to the interior of each edge of G_s is locally an isometric embedding.
- (2) Semiflow property: $h_{ut} \circ h_{ts} = h_{us}$ for all $s \leq t \leq u \in \mathbb{R}$ and $h_{ss}: G_s \rightarrow G_s$ is the identity for all $s \in \mathbb{R}$.

Handel and Mosher ([HM11, §7.3]) prove each flow line converges (in the axes or Gromov-Hausdorff topologies) to a point T_∞ in $\overline{\mathcal{X}}_r$, the direct limit of the system.

Theorem 9.2. *For any flow line, its direct limit T_∞ is an F_r -tree that has trivial arc stabilizers. Hence, in particular, not every point in the boundary is the direct limit of a flow line.*

Proof. We lift the maps to a direct system on trees $\{f_{ts}: T_s \rightarrow T_t\}$ that are F_r -equivariant, restrict to isometric embeddings on edges, and form a direct system. In [HM11, §7.3] it is shown that the maps $f_{\infty s}: T_s \rightarrow T_\infty$ that are given by the direct limit construction are also edge-isometries and F_r -equivariant.

Assume $x, y \in T_\infty$ are such that $\gamma \in \text{Stab}[x, y]$. Without generality loss, assume $d(x, y) = 1$ in T_∞ . We will show this leads to a contradiction. For each $t \in [a, \infty)$, we denote by $A_t(\gamma)$ the axis of γ in T_t . Letting $\varepsilon = \frac{1}{5}$, there exists some $s \geq a$ so that: $f_{\infty s}(x_s) = x$, $f_{\infty s}(y_s) = y$, and $d(x_s, \gamma x_s) < \varepsilon$, $d(y_s, \gamma y_s) < \varepsilon$, and $|d(x_s, y_s) - 1| < \varepsilon$.

Since f_{ts} is distance non-increasing for all t, s , we have for all $t \geq s$ that

$$1 \leq d(f_{ts}(x_s), f_{ts}(y_s)) \leq 1 + \varepsilon.$$

Note that for all $f_{ts}(x_s)$ and $f_{ts}(y_s)$, they are at most a distance of $\frac{\varepsilon}{2}$ from $A_t(\gamma)$. Otherwise, for example, $d(f_{ts}(x_s), \gamma f_{ts}(x_s)) \geq \varepsilon$, which contradicts the fact that the maps are distance non-increasing.

Thus, for each $t \geq s$ there exist z_t, w_t such that $[z_t, w_t] \subset A_t(\gamma) \cap [f_{ts}(x_s), f_{ts}(y_s)]$, and $d(z_t, f_{ts}(x_s)) < \frac{\varepsilon}{2}$ and $d(w_t, f_{ts}(y_s)) < \frac{\varepsilon}{2}$. Hence, $d(z_t, w_t) \geq 1 - \varepsilon$.

Let N be the number of $f_{\infty s}$ -illegal turns in the path $[x_s, y_s]$. Thus, the number of $f_{\infty t}$ -illegal turns in the path $[f_{ts}(x_s), f_{ts}(y_s)]$ is $\leq N$. Hence, the number of $f_{\infty t}$ -illegal turns in the path $[z_t, w_t]$ is $\leq N$. Let us denote the points of $[z_t, w_t]$ where the illegal turns occur by a_t^1, \dots, a_t^N , and we also denote $a_t^0 := z_t$ and $a_t^{N+1} := w_t$.

However, for sufficiently large t , the translation length of γ in T_t is $< \frac{1-\varepsilon}{3(N+1)}$. Note that since $[z_t, w_t]$ is on $A(\gamma)$, we have that $[z_t, w_t] \cap \gamma[z_t, w_t]$ is equal to $[z_t, w_t]$ with (possibly) segments of lengths $\leq \frac{1-\varepsilon}{3(N+1)}$ cut from either end. Since there are $\leq N+1$ segments (a_t^i, a_t^{i+1}) in $[z_t, w_t]$, one of them has length $\geq \frac{1-\varepsilon}{N+1}$. Thus, for some $i = 0, 1, \dots, N$, we have $\gamma(a_t^i) \in (a_t^i, a_t^{i+1})$ or $\gamma^{-1}(a_t^i) \in (a_t^{i-1}, a_t^i)$, which are both legal segments. Without generality loss suppose the former. Thus, the turn taken by $[z_t, w_t]$ at a_t^i is illegal, but the turn taken by $\gamma[z_t, w_t]$ at $\gamma(a_t^i)$ is legal, since it equals the turn taken by $[z_t, w_t]$ at $\gamma(a_t^i)$. This is a contradiction to the equivariance. \square

REFERENCES

- [AK11] Y. Algom-Kfir, *Strongly contracting geodesics in outer space*, *Geom. Topol.* **15** (2011), no. 4, 2181–2233. MR 2862155
- [Bes11] Mladen Bestvina, *A Bers-like proof of the existence of train tracks for free group automorphisms*, *Fund. Math.* **214** (2011), no. 1, 1–12. MR 2845630 (2012m:20046)
- [Bes12] M. Bestvina, *PCMI lectures on the geometry of outer space*, http://www.math.utah.edu/pcmi12/lecture_notes/bestvina.pdf (2012).
- [BF14] M. Bestvina and M. Feighn, *Hyperbolicity of the complex of free factors*, *Adv. Math.* **256** (2014), 104–155. MR 3177291
- [BH92] M. Bestvina and M. Handel, *Train tracks and automorphisms of free groups*, *The Annals of Mathematics* **135** (1992), no. 1, 1–51.
- [Bru57] V. Brun, *Algorithmes euclidiens pour trois et quatre nombres*, *Treizième congrès des mathématiciens scandinaves, tenu à Helsinki (1957)*, 45–64.
- [CV86] M. Culler and K. Vogtmann, *Moduli of graphs and automorphisms of free groups*, *Inventiones mathematicae* **84** (1986), no. 1, 91–119.
- [Die05] Reinhard Diestel, *Graph theory (graduate texts in mathematics)*.

- [EM11] Alex Eskin and Maryam Mirzakhani, *Counting closed geodesics in moduli space*, J. Mod. Dyn. **5** (2011), no. 1, 71–105. MR 2787598 (2012f:37070)
- [Fis09] A. Fisher, *Nonstationary mixing and the unique ergodicity of adic transformations*, Stochastics and Dynamics **9** (2009), no. 03, 335–391.
- [HM11] M. Handel and L. Mosher, *Axes in Outer Space*, no. 1004, Amer Mathematical Society, 2011.
- [Kea77] Michael Keane, *Non-ergodic interval exchange transformations*, Israel Journal of Mathematics **26** (1977), no. 2, 188–196.
- [KN76] Harvey B Keynes and Dan Newton, *A minimal, non-uniquely ergodic interval exchange transformation*, Mathematische Zeitschrift **148** (1976), no. 2, 101–105.
- [Mas81] H. Masur, *Dense geodesics in moduli space*, Riemann Surfaces and Related Topics (Stony Brook Conference), Ann. Math. Studies, vol. 97, 1981, pp. 417–438.
- [Mas82] ———, *Interval exchange transformations and measured foliations*, Ann. of Math **115** (1982), no. 1, 169–200.
- [Sch82] F. Schweiger, *Ergodische Eigenschaften der Algorithmen von Brun und Selmer*, Österreich. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II **191** (1982), no. 8-9, 325–329. MR 716668 (84m:10046)
- [Sch00] ———, *Multidimensional continued fractions*, Oxford University Press, 2000.
- [Thu] William Thurston, *Minimal stretch maps between hyperbolic surfaces*, arXiv:math.GT/9801039 math.GT.
- [Vee82] W. A. Veech, *Gauss measures for transformations on the space of interval exchange maps*, Annals of Mathematics (1982), 201–242.
- [Vog02] K. Vogtmann, *Automorphisms of free groups and outer space*, Geometriae Dedicata **94** (2002), no. 1, 1–31.

Department of Mathematics, University of Haifa
<http://www.math.haifa.ac.il/algomkfir/>,
E-mail address: yalgom@univ.haifa.ac.il

Department of Mathematics, University of California at Santa Barbara
<http://math.ucsb.edu/~cpfaff/>,
E-mail address: cpfaff@math.ucsb.edu