

## THE SPACE OF BINARY THETA SERIES

ERNST KANI

RÉSUMÉ. Le but de cet article est d'étudier l'espace  $\Theta_D$  engendré par les séries thêta binaires  $\vartheta_f$  qui sont attachées aux formes  $f$  primitives positives binaires quadratiques d'un discriminant  $D < 0$  fixé. En particulier, nous expliquons comment une telle série thêta  $\vartheta_f$  peut être exprimée comme une combinaison linéaire des éléments de la base (étendue) d'Atkin-Lehner pour les formes modulaires de poids un. De plus, nous montrons que cet espace a une base naturelle  $\{\vartheta_\chi\}_\chi$  consistant de fonctions propres de Hecke  $\vartheta_\chi$  (attachées aux caractères  $\chi$  du groupe  $\text{Cl}(D)$  des classes des formes de discriminant  $D$ ) et nous calculons les fonctions  $L(s, \vartheta_\chi)$  associées.

ABSTRACT. The purpose of this paper is to study the space  $\Theta_D$  generated by the binary theta series  $\vartheta_f$  attached to the primitive positive binary quadratic forms  $f$  of a fixed discriminant  $D < 0$ . In particular, we explain how each such theta series  $\vartheta_f$  can be expressed as a linear combination of the elements of the (extended) Atkin-Lehner basis for modular forms of weight one. Furthermore, we show that this space has a natural basis  $\{\vartheta_\chi\}_\chi$  consisting of Hecke eigenforms  $\vartheta_\chi$  (attached to characters  $\chi$  of the class group  $\text{Cl}(D)$  of forms of discriminant  $D$ ) and work out the associated  $L$ -functions  $L(s, \vartheta_\chi)$ .

### 1. Introduction

Let  $\Theta_D$  denote the space generated by the binary theta series  $\vartheta_f$  attached to the primitive, positive-definite binary quadratic forms  $f(x, y) = ax^2 + bxy + cy^2$  of discriminant  $D = b^2 - 4ac < 0$ . It is curious that while much has been written about the space generated by theta series attached to quadratic forms in  $2k \geq 4$  variables (cf. [15] and the references therein), the binary case does not seem to have been treated in detail in the literature. It is the purpose of this paper to fill this gap by presenting a variety of known and unknown results about this space.

By the work of Weber [33], Hecke [12] and Schoeneberg [26], it is known that  $\Theta_D$  is a subspace of the space  $M_1(|D|, \psi_D)$  of modular forms of weight 1, level  $|D|$  and Nebentypus  $\psi_D$ , where  $\psi_D = \left(\frac{D}{\cdot}\right)$  is the Kronecker-Legendre character. Contrary to the case of higher weight, the space of binary theta series is often a proper subspace of  $M_1(|D|, \psi_D)$  (cf. Remark 16) and so it is of interest to be able to identify it inside the space of modular forms.

As a first step towards this, we explain in this paper how each theta series  $\vartheta_f$  can be expressed explicitly as a linear combination of the *canonical* (extended) Atkin-Lehner

basis of  $M_1(|D|, \psi_D)$ ; cf. Remark 30(b). This (and other results) are used in the paper [17] to give an *intrinsic* description of the space of all theta series.

To this end we first observe that  $\Theta_D$  has a natural basis  $\{\vartheta_\chi\}$  indexed by the characters  $\chi \in \text{Cl}(D)^*$  of the class group  $\text{Cl}(D)$  of forms of discriminant  $D$ . It turns out that each  $\vartheta_\chi$  is a normalized eigenform with respect to Hecke algebra  $\mathbb{T}(D)$ ; the latter is the algebra generated by all Hecke operators  $T_n$  with  $(n, D) = 1$ .

**Theorem 1.** *The space  $\Theta_D$  is a  $\mathbb{T}(D)$ -submodule of  $M_1(|D|, \psi_D)$  of multiplicity one, and has a canonical basis  $\{\vartheta_\chi\}$  consisting of normalized  $\mathbb{T}(D)$ -eigenforms. Furthermore,  $\vartheta_\chi$  is a cusp form if and only if  $\chi$  is not a quadratic character.*

This theorem implies the following interesting result.

**Theorem 2.** *We have  $\Theta_D = \Theta_D^E \oplus \Theta_D^S$ , where  $\Theta_D^E = \Theta_D \cap E_1(|D|, \psi_D)$  denotes the Eisenstein space part and  $\Theta_D^S = \Theta_D \cap S_1(|D|, \psi)$  denotes the cusp space part of  $\Theta_D$ . Moreover,*

$$(1) \quad \dim \Theta_D^E = g_D \quad \text{and} \quad \dim \Theta_D^S = \frac{1}{2}(h_D - g_D),$$

where  $h_D = |\text{Cl}(D)|$  denotes the number of classes of forms of discriminant  $D$ , and  $g_D = [\text{Cl}(D) : \text{Cl}(D)^2]$  denotes the number of genera.

Note that it follows from (1) that  $\Theta_D$  has no non-zero cusp forms if and only if  $h_D = g_D$ , i.e., if and only if  $D$  is an *idoneal discriminant*. As is explained in Remark 17(b) below, this can be viewed as an alternate version of Theorem 3 of Kitaoka [18]. Moreover, Theorem 2 implies that a classical result of Siegel [29] (for quadratic forms in  $m \geq 4$  variables) extends to binary forms; cf. Corollary 18.

If  $D$  is a *fundamental discriminant*, i.e., if  $D = d_K$ , where  $d_K$  is the discriminant of  $K := \mathbb{Q}(\sqrt{D})$ , then Theorems 1 and 2 are essentially well-known, although I have not been able to find an explicit reference (except for the comments in Hecke [14], p. 792). Indeed, in this case each  $\vartheta_\chi$  is the modular form  $f(z; \chi)$  associated to  $\chi$  (viewed as a Hecke character) by Hecke’s construction, and so each  $\vartheta_\chi$  is a primitive form (newform); cf. Remark 16. In particular, in this case the  $\vartheta_\chi$ ’s are part of the canonical Atkin-Lehner basis.

However, if  $D$  is not a fundamental discriminant, then this is no longer true:  $\Theta_D$  is *not contained* in the space generated by the modular forms  $f(z; \chi)$  associated to Hecke characters (cf. Theorem 3 and Corollary 33). More precisely,  $\vartheta_\chi$  is equal to the modular form  $f(z; \chi)$  associated to the character  $\chi$  (viewed as Hecke character) if and only if  $\chi \in \text{Cl}(D)^*$  is *primitive*, i.e.,  $\chi$  is *not equal* to the lift  $\chi' \circ \pi$  of any character  $\chi' \in \text{Cl}(D')^*$  of some “lower level”  $D'|D$ ,  $D' \neq D$ , via the canonical map  $\pi = \pi_{D,D'} : \text{Cl}(D) \rightarrow \text{Cl}(D')$ . In the general case the situation is as follows.

**Theorem 3.** *Let  $\chi$  be a character on the class group  $\text{Cl}(D)$ , where  $D = f_D^2 d_K$ .*

(a) *There is a unique divisor  $f_\chi | f_D$  and a unique primitive character  $\chi_{pr}$  on the class group  $\text{Cl}(D_\chi)$ , where  $D_\chi = f_\chi^2 d_K$ , such that  $\chi = \chi_{pr} \circ \bar{\pi}_{D,D_\chi}$ .*

(b) *The form  $\vartheta_{\chi_{pr}} \in \Theta_{D_\chi}$  is a primitive form of level  $|D_\chi|$  whose  $L$ -function is the Hecke  $L$ -function associated to a suitable Hecke character  $\tilde{\chi}_{pr}$ ; in other words,*

$L(s, \vartheta_{\chi_{pr}}) = L(s, \tilde{\chi}_{pr})$ . Moreover, there exist constants  $c_n(\chi) \in \mathbb{R}$  such that

$$(2) \quad \vartheta_\chi(z) = \sum_{n|\bar{f}_\chi^2} c_n(\chi) \vartheta_{\chi_{pr}}(nz),$$

where  $\bar{f}_\chi = f_D/f_\chi$ . Furthermore, the function  $n \mapsto c_n(\chi)$  is multiplicative and has the generating function

$$(3) \quad C(s, \chi) := \sum_{n|\bar{f}_\chi^2} c_n(\chi) n^{-s} = L(s, \vartheta_\chi)/L(s, \vartheta_{\chi_{pr}}) = L(s, \vartheta_\chi)/L(s, \tilde{\chi}_{pr}).$$

Note that while  $L(s, \vartheta_{\chi_{pr}}) = L(s, \tilde{\chi}_{pr})$  is a classical Hecke  $L$ -function and hence is well-understood, the  $L$ -function  $L(s, \vartheta_\chi)$  is more complicated and is, in fact, *unknown*. (It is computed incorrectly in [21] and in [1]; cf. Remark 11.) Thus, (3) does not help in determining the constants  $c_n(\chi)$ . Instead, we calculate  $C(s, \chi)$  directly in Theorem 28 by using facts about ideals in quadratic orders which are presented in the Appendix (see §6). As a consequence, we thus obtain not only the desired explicit formula for  $\vartheta_f$  in terms of the Atkin-Lehner basis (cf. Remark 30(b)) but also an explicit expression for  $L$ -function  $L(s, \chi)$  which seems to be new; cf. Corollary 31. In particular, we obtain

**Corollary 4.** *If  $p^{\bar{e}_p} || \bar{f}_\chi$  denotes the largest power of  $p$  dividing  $\bar{f}_\chi$ , then the  $p$ -Euler factor  $L_p(s, \chi)$  of  $L(s, \chi)$  at  $p|\bar{f}_\chi$  is*

$$L_p(s, \chi) = \frac{1 - p^{(1-2s)\bar{e}_p}}{1 - p^{1-2s}} + \frac{\left(1 - \frac{1}{p}\psi_{D_\chi}(p)\right) p^{(1-2s)\bar{e}_p}}{1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s}}.$$

At the end of §5, we present some special cases and numerical examples of these results; cf. Examples 35 and 37.

After the first version of this paper was completed, Norm Hurt drew my attention to the papers of Sun and Williams [30, 31] which are partially related to some of the topics mentioned here. Indeed, Corollaries 31 and 4 can be viewed (for  $D < 0$ ) as a generalization of many of their results; cf. Remark 32(c) for more details.

## 2. The basis $\{\vartheta_\chi\}$ of $\Theta_D$

As in the introduction, let  $f(x, y) = ax^2 + bxy + cy^2$  be a primitive, positive definite binary quadratic form of discriminant  $D = b^2 - 4ac < 0$ . Thus  $a, b, c \in \mathbb{Z}$  and  $(a, b, c) = 1$ , and  $D = f_D^2 d_K$ , where  $d_K$  is the (fundamental) discriminant of the imaginary quadratic field  $K = \mathbb{Q}(\sqrt{D})$ . For a fixed  $D$ , let  $Q_D$  denote the set of all such forms. The *binary theta series* attached to  $f \in Q_D$  is the function on the upper half-plane  $\mathfrak{H}$  given by

$$\vartheta_f(z) = \sum_{x, y \in \mathbb{Z}} e^{2\pi i f(x, y)z} = \sum_{n=0}^{\infty} r_n(f) e^{2\pi i n z},$$

where  $r_n(f) = \#\{(x, y) \in \mathbb{Z}^2 : f(x, y) = n\}$  denotes the number of representations of  $n$  by  $f$ . The following fact is fundamental for most of this paper.

**Proposition 5.** *If  $f \in Q_D$ , then  $\vartheta_f$  is a modular form of level  $|D|$  and Nebentypus  $\psi_D = \left(\frac{D}{\cdot}\right)$ . Thus  $\Theta_D := \sum_{f \in Q_D} \mathbb{C}\vartheta_f$  is a subspace of  $M_1(|D|, \psi_D)$ .*

**Proof.** See Schoeneberg [27], ch. IX, Theorems 4 and 5, or Miyake [22], Corollary 4.9.5(3).  $\square$

**Remark 6.** (a) This result was proved by Weber [33] in 1893 (see his formula §15 (21)), except that he did not verify that  $\vartheta_f$  is holomorphic at all the cusps. In 1926 Hecke [12] proved a more general result from which above result can be deduced (cf. Remark 11(c)). Siegel [29] proved something weaker but for quadratic forms in an arbitrary number  $m \geq 2$  of variables; cf. [29], Hilfssatz 30 and 31 and also Satz 4, where the case  $m = 2$  is excluded. In 1939 Schoeneberg [26] proved a very general result about theta series which includes the above result as a special case.

(b) The  $L$ -function associated to the modular form  $\vartheta_f$  is

$$Z_f(s) = L(s, \vartheta_f) = \sum_{n \geq 1} \frac{r_n(f)}{n^s} = \sum_{(x,y) \neq (0,0)} f(x,y)^{-s}, \quad \text{where } \operatorname{Re}(s) > 1.$$

This function is often called the *Epstein zeta-function* of  $f$  (cf. e.g. [7]), even though it was introduced by Dirichlet in 1839 (cf. [8], §6.18 (p. 358)) and was studied intensively by him and by Kronecker [20] (and by others) many years before Epstein.

Recall that the group  $\operatorname{GL}_2(\mathbb{Z})$  acts on binary quadratic forms by change of coordinates, and that this action preserves the set  $Q_D$ . It is immediate that  $r_n(fT) = r_n(f)$ , for all  $n \geq 0$  and  $T \in \operatorname{GL}_2(\mathbb{Z})$ , so  $\vartheta_{fT} = \vartheta_f$ . We can thus index the theta series by the quotient set

$$\overline{Q}_D = Q_D / \operatorname{GL}_2(\mathbb{Z}) = Q_D / \approx,$$

where  $\approx$  denotes the equivalence relation induced by  $\operatorname{GL}_2(\mathbb{Z})$ -equivalence.

**Proposition 7.** <sup>1</sup> *The set  $\{\vartheta_f : f \in \overline{Q}_D\}$  is a basis of the space  $\Theta_D$ . Thus*

$$\dim \Theta_D = \overline{h}_D := \#\overline{Q}_D.$$

As we shall see, this follows easily from the following basic facts about binary quadratic forms which are due to Dirichlet [8] and Weber [32].

**Lemma 8.** *Let  $f, f_1, f_2 \in Q_D$  be primitive quadratic forms. Then:*

- (a) *There exist infinitely many prime numbers  $p$  with  $r_p(f) > 0$ .*
- (b) *If there is a prime number  $p \nmid D$  with  $r_p(f_i) > 0$ , for  $i = 1, 2$ , then  $f_1 \approx f_2$ .*

**Proof.** (a) This was first proved by Dirichlet [8] for prime discriminants and by Weber [32] in the general case. A proof using class field theory is given in Cox [6], Theorem 9.12.

(b) Weber [33] states this (elementary) result on p. 259 and points out that it follows easily from his paper [32] (or from a paper of Schering). It is restated and proved (without references) as Satz 1 in Piehler [24].  $\square$

<sup>1</sup>The referee pointed out that this can be viewed as a special case of the lemma on p. 158 of Kitao-ka [19]

**Proof of Proposition 7.** Let  $f_1, \dots, f_N \in Q_D$  be a system of representatives of  $\overline{Q}_D$ ; thus,  $N = \overline{h}_D$ . It is clear from what was said above that  $\vartheta_{f_1}, \dots, \vartheta_{f_N}$  generate  $\Theta_D$ . To prove that they are linearly independent, suppose that  $c_1\vartheta_{f_1} + \dots + c_N\vartheta_{f_N} = 0$ , for some  $c_1, \dots, c_N \in \mathbb{C}$ . Thus  $c_1r_n(f_1) + \dots + c_Nr_n(f_N) = 0$ , for all  $n \geq 0$ .

Fix  $i$ . By Lemma 8(a) we know that there is a prime  $p = p_i \nmid D$  with  $r_p(f_i) > 0$ , and from Lemma 8(b) it follows that  $r_p(f_j) = 0$  for all  $j \neq i$ . We thus have that  $c_i r_p(f_i) = 0$ , so  $c_i = 0$ , and hence  $\{\vartheta_{f_i}\}$  is a basis of  $\Theta_D$ , as claimed.  $\square$

We now introduce another basis of the space  $\Theta_D$ . For this, recall that by Gauss’s theory of composition of forms (cf. Gauss [9]) the set

$$\text{Cl}(D) := Q_D/\text{SL}_2(\mathbb{Z}) = Q_D/\sim$$

has the structure of an abelian group. The identity of  $\text{Cl}(D)$  is the class of the principal form  $1_D$  which is defined by  $1_D(x, y) = x^2 + \varepsilon xy + \frac{\varepsilon - D}{4}y^2$ , for  $D \equiv \varepsilon \pmod{4}$  and  $\varepsilon \in \{0, 1\}$ . Note also that the  $\text{SL}_2(\mathbb{Z})$ -equivalence relation  $\sim$  is related to the previous  $\text{GL}_2(\mathbb{Z})$ -equivalence relation  $\approx$  by

$$(4) \quad f_1 \approx f_2 \iff f_1 \sim f_2 \text{ or } f_1 \sim f_2^{-1},$$

where  $f_2^{-1}$  denotes a representative of the inverse class of  $f_2$  (in the group  $\text{Cl}(D)$ ). From this it follows easily that

$$(5) \quad \overline{h}_D = \frac{1}{2}(g_D + h_D), \quad \text{where } h_D = |\text{Cl}(D)| \text{ and } g_D = [\text{Cl}(D) : \text{Cl}(D)^2].$$

Let  $\chi \in \text{Cl}(D)^* := \text{Hom}(\text{Cl}(D), \mathbb{C}^\times)$  be a character on  $\text{Cl}(D)$ , and put

$$(6) \quad \vartheta_\chi = \frac{1}{w_D} \sum_{f \in \text{Cl}(D)} \chi(f)\vartheta_f \in \Theta_D,$$

where  $w_D = r_1(1_D)$ . (Thus, as is well known,  $w_D = 2$  when  $D < -4$  and  $w_{-4} = 4$  and  $w_{-3} = 6$ .) Note that the terms on the right-hand side are not linearly independent since we sum over  $\text{Cl}(D)$  in place of  $\overline{Q}_D$ . However, by using (4) and noting that  $\chi(f^{-1}) = \overline{\chi(f)}$ , we can re-write (6) in the form

$$(7) \quad \vartheta_\chi = \frac{1}{w_D} \sum_{\bar{f} \in \overline{Q}_D} w(\bar{f})\text{Re}(\chi(\bar{f}))\vartheta_{\bar{f}},$$

where  $w(\bar{f}) = \#\{f \in \text{Cl}(D) : f \approx \bar{f}\}$ . (Thus,  $w(f) = 1$ , if  $f^2 \sim 1_D$  and  $w(f) = 2$  otherwise.) From this expression we see immediately that  $\vartheta_{\chi^{-1}} = \vartheta_\chi$ , and so it is useful to index the  $\vartheta_\chi$ ’s by the set  $\overline{\text{Cl}(D)}^* = \text{Cl}(D)^*/(\chi \mapsto \chi^{-1})$ . We then have:

**Proposition 9.** *If  $f \in Q_D$  is a primitive quadratic form, then*

$$(8) \quad \vartheta_f = \frac{w_D}{h_D} \sum_{\chi \in \overline{\text{Cl}(D)}^*} \overline{\chi}(f)\vartheta_\chi = \frac{w_D}{h_D} \sum_{\chi \in \overline{\text{Cl}(D)}^*} w(\chi)\text{Re}(\chi(f))\vartheta_\chi,$$

where  $w(\chi) = \#\{\chi, \chi^{-1}\}$ . Thus  $\{\vartheta_\chi : \chi \in \overline{\text{Cl}(D)}^*\}$  is also a basis of  $\Theta_D$ .

**Proof.** Since  $\vartheta_{\chi^{-1}} = \vartheta_{\chi}$ , the second identity of (8) is clear. To prove the first identity, recall that the orthogonality relations for group characters imply that

$$\sum_{\chi \in \text{Cl}(D)^*} \chi(f) = \begin{cases} h_D & \text{if } f \sim 1_D, \\ 0 & \text{otherwise.} \end{cases}$$

From this, together with the definition (6) of  $\vartheta_{\chi}$ , it follows that

$$\begin{aligned} w_D \sum_{\chi \in \text{Cl}(D)^*} \bar{\chi}(f) \vartheta_{\chi} &= \sum_{\chi \in \text{Cl}(D)^*} \bar{\chi}(f) \sum_{f_1 \in \text{Cl}(D)} \chi(f_1) \vartheta_{f_1} \\ &= \sum_{\chi \in \text{Cl}(D)^*} \chi(f^{-1}) \sum_{f_1 \in \text{Cl}(D)} \chi(f_1) \vartheta_{f_1} \\ &= \sum_{f_1 \in \text{Cl}(D)} \vartheta_{f_1} \sum_{\chi \in \text{Cl}(D)^*} \chi(f^{-1} f_1) = h_D \vartheta_f, \end{aligned}$$

which proves the first identity of (8) and hence (8) itself.

From (8) it is clear that the set  $\{\vartheta_{\chi} : \chi \in \overline{\text{Cl}(D)^*}\}$  generates the space  $\Theta_D$ . Since  $\text{Cl}(D) \simeq \text{Cl}(D)^*$ , it follows that

$$\#(\overline{\text{Cl}(D)^*}) = \#(\text{Cl}(D)/(f \mapsto f^{-1})) = \#\bar{Q}_D = \dim \Theta_D,$$

where the last two identities follow from (4) and Proposition 7, respectively. Thus, the set  $\{\vartheta_{\chi} : \chi \in \overline{\text{Cl}(D)^*}\}$  is a basis of  $\Theta_D$ . □

We now examine the Fourier coefficients  $a_n(\chi) := a_n(\vartheta_{\chi})$  of the modular form  $\vartheta_{\chi}$  more closely. For this we shall use a basic result due to Dedekind that the class group  $\text{Cl}(D)$  can be identified with the group  $\text{Cl}(\mathfrak{O}_D) = \text{Pic}(\mathfrak{O}_D) = I(\mathfrak{O}_D)/P(\mathfrak{O}_D)$  of classes of invertible (fractional) ideals of the order  $\mathfrak{O}_D$  of discriminant  $D$  in the quadratic field  $K = \mathbb{Q}(\sqrt{D})$ . (Here we use the terminology and notation of Appendix §6.1.) More precisely, if  $L(f) = a\mathbb{Z} + \frac{-b+\sqrt{D}}{2}\mathbb{Z}$  denotes the quadratic lattice in  $K$  associated to the form  $f(x, y) = ax^2 + bxy + cy^2 \in Q_D$ , then the rule  $f \mapsto L(f)$  defines an isomorphism

$$\lambda_D : \text{Cl}(D) \xrightarrow{\sim} \text{Cl}(\mathfrak{O}_D) := I(\mathfrak{O}_D)/P(\mathfrak{O}_D);$$

cf. Cox[6], Theorem 7.7 (p. 137). We observe:

**Proposition 10.** *Let  $\chi \in \text{Cl}(D)^*$ . If  $n \geq 1$ , then the  $n$ -th Fourier coefficient of  $\vartheta_{\chi}$  is given by the formula*

$$(9) \quad a_n(\chi) = \sum_{\mathfrak{a} \in \text{Id}_n(\mathfrak{O}_D)} \chi^*(\mathfrak{a}),$$

where  $\text{Id}_n(\mathfrak{O}_D)$  denotes the set of invertible ideals of  $\mathfrak{O}_D$  of norm  $n$  and

$$\chi^* = \chi \circ \lambda_D^{-1} \in \text{Cl}(\mathfrak{O}_D)^*.$$

Moreover,  $a_0(\chi) = 0$  except when  $\chi = 1$  is trivial, and then  $a_0(1) = \frac{h_D}{w_D}$ .

**Proof.** Recall that if  $f \in Q_D$ , then

$$f(x, y) = ax^2 + bxy + cy^2 = N_K(ax - \beta_f y)/N(L),$$

where  $\beta_f = \frac{-b+\sqrt{D}}{2}$ ,  $N_K$  denotes the field norm and  $N(L) = a$  is the norm of the lattice  $L := L(f)$ ; cf. [6], p. 137. We thus obtain (in the notation of the appendix) that

$$\begin{aligned} (10) \quad \vartheta_f(z) &= \sum_{x,y \in \mathbb{Z}} e^{2\pi i f(x,y)z} \\ &= \sum_{\alpha \in L(f)} e^{2\pi i N_K(\alpha)z/N(L(f))} = 1 + w_D \sum_{\substack{\mathfrak{a} \in Id(\mathfrak{O}_D) \\ \mathfrak{a}L(f) \in P(\mathfrak{O}_D)}} e^{2\pi i N(\mathfrak{a})z}, \end{aligned}$$

where the last equality follows from the fact that  $\mathfrak{a} := \alpha L(f)^{-1}$  is an integral  $\mathfrak{O}_D$ -ideal if and only if  $\alpha \in L(f)$  (where  $\alpha \in K^\times$ ), together with the fact that  $|\mathcal{O}_D^\times| = w_D$ . From this it thus follows from the definition that

$$\begin{aligned} \vartheta_\chi(z) &= \frac{1}{w_D} \sum_{f \in Cl(D)} \chi(f) \left( 1 + w_D \sum_{\substack{\mathfrak{a} \in Id(\mathfrak{O}_D) \\ \mathfrak{a}L(f) \in P(\mathfrak{O}_D)}} e^{2\pi i N(\mathfrak{a})z} \right) \\ &= c_\chi + \sum_{\mathfrak{a} \in Id(\mathfrak{O}_D)} \chi^*(\mathfrak{a}) e^{2\pi i N(\mathfrak{a})z}, \end{aligned}$$

where  $c_\chi = \frac{1}{w_D} \sum_{f \in Cl(D)} \chi(f)$ . This proves (9). Note that  $c_\chi = 0$  except when  $\chi = 1$ ; in the latter case clearly  $c_\chi = \frac{h_D}{w_D}$ . □

**Remark 11.** (a) The above result shows that the  $L$ -function associated to  $\vartheta_\chi$  is

$$L(s, \vartheta_\chi) = \sum_{n \geq 1} \frac{a_n(\chi)}{n^s} = \sum_{\mathfrak{a} \in Id(\mathfrak{O}_D)} \frac{\chi^*(\mathfrak{a})}{N(\mathfrak{a})^s},$$

where  $\chi^* = \chi \circ \lambda_D^{-1}$ . Thus,  $L(s, \vartheta_\chi)$  coincides with the  $L$ -function  $L_{\mathfrak{O}_D}(s, \chi^*)$  defined by the first equation on the bottom p. 280 of the first edition of [21]. (Note, however, that the Euler product given by second equation in [21] is incorrect when  $D$  is not fundamental; this is why Lang restricts his discussion to fundamental discriminants in the second edition.) Similarly,  $L(s, \vartheta_\chi) = L_{\mathcal{F}_\chi}(s)$ , where the latter is as on p. 204 of Antoniadis [1]; in fact, we already have that  $\vartheta_\chi(z) = \mathcal{F}_\chi(z)$ . Here too, the given Euler product is incorrect in general, as Corollary 4 shows. (In fact, the assertion stated there (without proof) that  $\mathcal{F}_\chi$  is always a newform is also incorrect; cf. Corollary 33.)

(b) For later reference we note that for the trivial character  $\chi = 1$  the equations (6) and (9) yield that

$$(11) \quad a_n(1) = \frac{1}{w_D} \sum_{f \in Cl(D)} r_n(f) = \#Id_n(\mathfrak{O}_D).$$

This number will be computed below (cf. Remark 32(b)) and is well-known when  $(n, f_D) = 1$ ; cf. (57).

(c) Note that it follows from the first part of (10) that if  $f \in Q_D$ , then

$$\vartheta_f(z) = \sum_{i=1}^{f_D} \vartheta(f_D z; \alpha_i \sqrt{d_K}, \mathfrak{a}_f, f_D \sqrt{d_K}),$$

where  $\vartheta(z; \rho, \mathfrak{a}, Q\sqrt{d_K}) = \sum_{\mu \equiv \rho \pmod{\mathfrak{a}Q\sqrt{d_K}}} e^{2\pi i z N_K(\mu)/(N(\mathfrak{a})Q|d_K|)}$  is as in Hecke [12],  $\mathfrak{a}_f = L(f)\mathcal{O}_K$ , and  $\{\alpha_i\}$  is a system of coset representatives of  $L(f)/f_D\mathfrak{a}_f$ . Thus,  $\Theta_D$  is a subspace of the space generated by dilations of Hecke’s theta-functions.

By using results about ideals in the ring  $\mathfrak{D}_D$  (cf. Appendix, §6.2), we obtain the following important result.

**Theorem 12.** *If  $\chi \in \text{Cl}(D)^*$ , then the function  $n \mapsto a_n(\chi)$  is multiplicative. Thus  $\vartheta_\chi$  is a normalized eigenfunction with eigenvalue  $a_n(\chi)$  with respect to the Hecke operator  $T_n$  whenever  $(n, f_D) = 1$ .*

**Proof.** Put  $\chi^* = \chi \circ \lambda_D^{-1} \in \text{Cl}(\mathfrak{D}_D)^*$ , and write  $Id_n = Id_n(\mathfrak{D}_D)$ . If  $(m, n) = 1$ , then from (9) together with Proposition 44 of §6.2 we obtain that

$$\begin{aligned} a_{mn}(\chi) &= \sum_{\mathfrak{a} \in Id_{mn}} \chi^*(\mathfrak{a}) \\ &= \sum_{\mathfrak{b} \in Id_m} \sum_{\mathfrak{c} \in Id_n} \chi^*(\mathfrak{bc}) \\ &= \sum_{\mathfrak{b} \in Id_m} \chi^*(\mathfrak{b}) \sum_{\mathfrak{c} \in Id_n} \chi^*(\mathfrak{c}) \\ &= a_m(\chi) a_n(\chi), \end{aligned}$$

which shows that the function  $n \mapsto a_n(\chi)$  is multiplicative. Moreover, since

$$a_1(\chi) = \chi^*(\mathfrak{D}_D) = 1,$$

we see that  $\vartheta_\chi$  is normalized.

From this it follows from Hecke [13], Satz 42, that  $\vartheta_\chi$  is a  $T_n$ -eigenfunction,<sup>2</sup> at least when  $(n, |D|) = 1$ . By using the results of [22], this can be refined to yield the above assertion also for  $(n, f_D) = 1$ , as we shall now show.

Fix  $n \geq 1$  with  $(n, f_D) = 1$ , and consider the function

$$g := (\vartheta_\chi)|_1 T_n - a_n(\chi) \vartheta_\chi \in M_1(|D|, \psi_D).$$

For any  $m \geq 1$  with  $(m, n) = 1$  we have by [22], Lemma 4.5.14, and the above result that the  $m$ -th Fourier coefficient of  $g$  is

$$a_m(g) = a_{nm}(\chi) - a_n(\chi) a_m(\chi) = 0.$$

Thus, since  $\psi_D$  has conductor  $|d_K|$  and since  $\frac{|D|}{|d_K|} = f_D^2$ , we see that  $g$  satisfies the hypothesis of Theorem 4.6.8(1) of [22] (with  $l = n$ ), and so  $g = 0$ . This means that  $\vartheta_\chi$  is a  $T_n$ -eigenfunction with eigenvalue  $a_n(\chi)$ , as asserted.  $\square$

<sup>2</sup>Note that this argument requires “only” the condition that  $a_{mn}(\chi) = a_m(\chi) a_n(\chi), \forall m, n \geq 1$  with  $(m, n) = (n, D) = 1$ . However, one cannot weaken this hypothesis to requiring this condition only for the  $m$ ’s satisfying the extra condition  $(m, D) = 1$ , as easy counterexamples show.

### 3. The Eisenstein and cuspidal parts of $\Theta_D$

By the basic theory of modular forms, the space  $M_k(N, \psi)$  of modular forms of level  $N$ , weight  $k$  and Nebentypus  $\psi$  has a canonical decomposition

$$(12) \quad M_k(N, \psi) = E_k(N, \psi) \oplus S_k(N, \psi)$$

into its Eisenstein part  $E_k(N, \psi)$  and cuspidal part  $S_k(N, \psi)$ ; cf. [22], Theorem 4.7.2 (together with Theorem 2.1.7).

Since  $\Theta_D \subset M_1(|D|, \psi_D)$ , we can define its Eisenstein and cuspidal part by

$$\Theta_D^E := \Theta_D \cap E_1(|D|, \psi_D) \quad \text{and} \quad \Theta_D^S := \Theta_D \cap S_1(|D|, \psi_D),$$

respectively.

We now want to find canonical bases for these spaces. As we shall see, the  $\vartheta_\chi$ 's serve this purpose: it turns out that either  $\vartheta_\chi \in \Theta_D^E$  or  $\vartheta_\chi \in \Theta_D^S$ ; cf. Theorem 14 and Remark 17(a) below.

To verify this, we shall compare the coefficients of  $\vartheta_\chi$  to those of a suitable modular form attached to some *Hecke character* on the ray class group mod  $f_D \mathfrak{D}_K$  of the quadratic field  $K = \mathbb{Q}(\sqrt{D})$ . To achieve this, recall first that the map  $\mathfrak{a} \mapsto \mathfrak{a} \cap \mathfrak{D}_D$  induces an isomorphism

$$\varphi_D : I_K(f_D)/P_{K,\mathbb{Z}}(f_D) \xrightarrow{\sim} \text{Cl}(\mathfrak{D}_D),$$

where  $I_K(f)$  is the group of fractional ideals of  $\mathfrak{D}_K$  which are prime to  $f$  and  $P_{K,\mathbb{Z}}(f)$  is the subgroup of  $I_K(f)$  generated by principal ideals of the form  $\alpha \mathfrak{D}_K$ , where the element  $\alpha \in \mathfrak{D}_K$  satisfies  $\alpha \equiv a \pmod{f \mathfrak{D}_K}$ , for some  $a \in \mathbb{Z}$  with  $(a, f) = 1$ ; cf. Cox[6], Proposition 7.22. We then have:

**Proposition 13.** *Let  $\chi \in \text{Cl}(D)^*$ , and let  $\tilde{\chi} := \chi \circ \lambda_D^{-1} \circ \varphi_D$  be the associated Hecke character on  $I_K(f_D)$ . If  $n \geq 1$ , then*

$$(13) \quad a_n(\chi) = a_n(\tilde{\chi}) := \sum_{\mathfrak{a} \in Id_n(\mathfrak{D}_K)} \tilde{\chi}(\mathfrak{a}), \quad \text{provided that } (n, f_D) = 1.$$

**Proof.** By Proposition 45 we know that the map  $\mathfrak{a} \mapsto \mathfrak{a} \cap \mathfrak{D}_D$  induces a bijection

$$Id_n(\mathfrak{D}_K) \xrightarrow{\sim} Id_n(\mathfrak{D}_D), \quad \text{whenever } (n, f_D) = 1,$$

and so the assertion (13) follows in view of (9). □

In the next sections we shall study the precise relation between  $a_n(\chi)$  and  $a_n(\tilde{\chi})$  also in the case that  $(n, f_D) > 1$ . As we shall see, the formula (13) is in general no longer true in this case, and has to be replaced by the more complicated formula (28) below. Nevertheless, the above relation (13) suffices to derive many useful properties about the  $\vartheta_\chi$ 's. To formulate these in a convenient manner, it is useful to introduce the following notation.

**Notation.** If  $f_1$  and  $f_2$  are two modular forms and if  $N$  an integer, then we write

$$f_1 \sim_N f_2 \stackrel{\text{def}}{\iff} a_n(f_1) = a_n(f_2), \quad \text{for all } n \geq 1 \text{ with } (n, N) = 1.$$

With this notation, we can then reformulate (13) in terms of the modular form  $f(z; \tilde{\chi})$  (cf. [22], p. 183) as follows:

$$(14) \quad \vartheta_\chi(z) \sim_{f_D} \sum_{\mathfrak{a} \in Id(\mathfrak{D}_K) \cap I_K(f_D)} \tilde{\chi}(\mathfrak{a}) e^{2\pi i N(\mathfrak{a})z} \sim_{f_D} f(z; \tilde{\chi}).$$

**Theorem 14.** *Let  $\chi \in \text{Cl}(D)^*$  be a character. If  $\chi$  is not quadratic, i.e., if  $\chi^2 \neq 1$ , then  $\vartheta_\chi$  is a cusp form, and otherwise  $\vartheta_\chi$  is in the Eisenstein space.*

**Proof.** Let  $\tilde{\chi} := \chi \circ \lambda_D^{-1} \circ \varphi_D$  be the Hecke character on  $I_K(f_D)$  associated to  $\chi$ .

Assume first that  $\chi^2 \neq 1$ . In this case we have that  $\tilde{\chi} \neq \psi \circ N_K$ , for any Dirichlet character  $\psi \pmod{f_D}$ . Indeed, if this were the case, then  $\psi$  is necessarily a quadratic character because if  $(a, f_D) = 1$ , then  $\psi(a^2) = \psi(N_K(a\mathfrak{D}_K)) = \tilde{\chi}(a\mathfrak{D}_K) = 1$  since  $a\mathfrak{D}_K \in P_{K, \mathbb{Z}}(f_D)$ . Thus  $\tilde{\chi}$  is also a quadratic character, contrary to the hypothesis.

Since  $\tilde{\chi} \neq \psi \circ N_K$ , we know by [22], Theorem 4.8.2, that  $f(z; \tilde{\chi}) \in S_1(N, \psi_D)$ , where  $N = |d_K| N_K(f_D \mathfrak{D}_K) = |D|$ ; clearly  $f(z; \tilde{\chi})$  is a  $\mathbb{T}(N)$ -eigenfunction. Since  $\vartheta_\chi \sim_{f_D} f(z; \tilde{\chi})$  by (14), it follows from Lemma 15(c) below (together with Theorem 12) that  $\vartheta_\chi$  is also a cusp form.

Now suppose that  $\chi \in \text{Cl}(D)^*$  is a quadratic character. Then by Gauss's genus theory there exist fundamental discriminants  $D_1, D_2$  such that  $D = c^2 D_1 D_2$  for some integer  $c$  and such that  $\tilde{\chi} = \psi_{D_1} \circ N_K = \psi_{D_2} \circ N_K$ ; cf. Weber[34], §104 and §109.

Since the pair  $(\psi_{D_1}, \psi_{D_2})$  satisfies condition (4.7.2)(ii) on p. 176 of [22], there is a form  $f_1 := f_1(z; \psi_{D_1}, \psi_{D_2}) \in E_1(|D_1 D_2|, \psi_{D_1 D_2}) \subset E_1(|D|, \psi_D)$  such that its  $L$ -function is  $L(s, f_1) = L(s, \psi_{D_1}) L(s, \psi_{D_2})$ ; cf. [22], Theorem 4.7.1. We now claim:

$$(15) \quad \vartheta_\chi(z) \sim_D f_1(z; \psi_{D_1}, \psi_{D_2}).$$

Indeed, since  $\tilde{\chi} = \psi_{D_2} \circ N_K$ , it follows from (13) that for  $(n, D) = 1$  we have that

$$(16) \quad \begin{aligned} a_n(\chi) &= a_n(\tilde{\chi}) = \sum_{\mathfrak{a} \in Id_n(\mathfrak{D}_K)} \psi_{D_2}(N(\mathfrak{a})) \\ &= \psi_{D_2}(n) \# Id_n(\mathfrak{D}_K) = \psi_{D_2}(n) \sum_{t|n} \psi_{d_K}(t), \end{aligned}$$

the latter by (57). Since  $\psi_{d_K}(t) = \psi_{D_1}(t) \psi_{D_2}(t)$  and

$$\psi_{D_2}(nt) = \psi_{D_2}\left(t^2 \frac{n}{t}\right) = \psi_{D_2}\left(\frac{n}{t}\right),$$

we thus see that  $a_n(\chi) = \sum_{t|n} \psi_{D_1}(t) \psi_{D_2}\left(\frac{n}{t}\right) = a_n(f_1; \psi_{D_1}, \psi_{D_2})$ , which proves (15). Using this, it follows from Lemma 15(c) below that  $\vartheta_\chi \in E_1(|D|, \psi_D)$ .  $\square$

Above we had used the following general fact about  $\mathbb{T}(N)$ -eigenfunctions which is proved implicitly on p. 179 of [22]. Here, as usual,  $\mathbb{T}(N) \subset \text{End}_{\mathbb{C}}(M_k(N, \psi))$  is the Hecke algebra generated by the Hecke operators  $T_n$  with  $(n, N) = 1$ .

**Lemma 15.** (a) *If  $f \in M_k(N, \psi)$  is a  $\mathbb{T}(N)$ -eigenfunction, then either  $f \in S_k(N, \psi)$  or  $f \in E_k(N, \psi)$ .*

(b) *If  $f_1, f_2 \in M_k(N, \psi)$  are two non-zero  $\mathbb{T}(N)$ -eigenfunctions which have the same  $\mathbb{T}(N)$ -eigenvalues, then either both are cusp forms or both are in  $E_k(N, \psi)$ .*

(c) Let  $f_1, f_2 \in M_k(N, \psi)$  be two normalized  $\mathbb{T}(N)$ -eigenfunctions such that  $f_1 \sim_N f_2$ . Then  $f_1$  is a cusp form if and only if  $f_2$  is a cusp form, and similarly  $f_1 \in E_k(N, \psi)$  if and only if  $f_2 \in E_k(N, \psi)$ .

**Proof.** (a) This is implicitly proven in the course of the proof of Theorem 4.7.2 of [22] on p. 179. For convenience of the reader, we sketch the proof.

If false, then  $f = f_1 + f_2$  with  $f_1 \in S_k(N, \psi)$ ,  $f_2 \in E_k(N, \psi)$  and  $f_i \neq 0$  for  $i = 1, 2$ . Then as [22] (*loc. cit.*),  $f_i$  is a  $\mathbb{T}(N)$ -eigenfunction with the same eigenvalues as  $f$  and there exist normalized eigenfunctions

$$\tilde{f}_1 \in S_k(N, \psi) \text{ and } \tilde{f}_2 = f_k(z; \chi_1, \chi_2) \in E_k(N, \psi)$$

with the same  $\mathbb{T}(N)$ -eigenvalues as  $f$ . Thus  $L(s, \tilde{f}_1) \approx_N L(s, \tilde{f}_2)$ , where  $\approx_N$  means equality except for the Euler factors at the primes  $p \mid N$ . If  $\chi_2$  is trivial, then this is impossible because  $\Gamma(s)L(s, \tilde{f}_1)$  is entire while  $\Gamma(s)L(s, \tilde{f}_2)$  has a pole at  $s = k$ , as is explained in [22], p. 179. In the general case, by considering the twists  $(f_1)_{\tilde{\chi}_2}$  and  $(f_2)_{\tilde{\chi}_2}$ , one obtains a similar contradiction.

(b) If false, then  $f = f_1 + f_2$  is a  $\mathbb{T}(N)$ -eigenfunction which contradicts part (a).

(c) The  $T_n$ -eigenvalue of  $f_i$  is  $a_n(f_1) = a_n(f_2)$  as each  $f_i$  is a normalized eigenfunction. Thus  $f_1$  and  $f_2$  satisfy the hypotheses of (b), and so the assertion follows.  $\square$

**Remark 16.** In the case that  $D = d_K$  is a fundamental discriminant, it follows from the above theorem that all the  $\vartheta_\chi$ 's are *primitive forms* in the sense of section 4 below. Indeed, since in this case  $\psi_D = \psi_{d_K}$  is a primitive character, it follows from the general theory of newforms (primitive forms in the terminology of [22], p. 164) that all  $\mathbb{T}(D)$ -eigenfunctions of  $S_1(|D|, \psi_D)$  are newforms; cf. [22], Lemma 4.6.9(1) and Theorem 4.6.12. In particular, it follows immediately that the  $\vartheta_\chi = f(z, \tilde{\chi})$  are newforms when  $\chi^2 \neq 1$ .

A similar result also holds for the Eisenstein series  $\vartheta_\chi$  when  $\chi^2 = 1$ , but this requires a different argument. Here we use instead the structure theorem of the Eisenstein space  $E_1(|D|, \psi_D)$ ; cf. [22], p. 179. Specialized to the present situation, it yields that

$$(17) \quad E_1(|d_K|, \psi_{d_K}) = \bigoplus_{d_K = D_1 D_2} \mathbb{C} f_1(z; \psi_{D_1}, \psi_{D_2}),$$

where the sum is over all factorizations  $d_K = D_1 D_2$  of  $d_K$  into fundamental discriminants  $D_1, D_2$  (and the factorization  $d_K = D_2 D_1$  is considered to be the same as the factorization  $d_K = D_1 D_2$ ), and  $f_1(z; \psi_{D_1}, \psi_{D_2})$  is as in the above proof. To verify this from formula (4.7.17) of [22], note first that if  $\psi_{d_K} = \psi_1 \psi_2$  is any factorization into characters  $\psi_i$  with conductors  $M_i$  such that  $M_1 M_2 \mid d_K$ , then necessarily  $|d_K| = M_1 M_2$  and  $(M_1, M_2) = 1$ . By adjusting signs, we thus have the factorization  $d_K = D_1 D_2$  into fundamental discriminants (with  $|D_i| = M_i$ ). Moreover, since  $\psi_i$  is necessarily quadratic (because  $(M_1, M_2) = 1$ ), it follows that

$$\psi_i = \psi_{D_i} = \left( \frac{D_i}{\cdot} \right)$$

is the Kronecker-Legendre character. This verifies (17).

From (17) it follows that we can improve the relation (15) to an equality because all  $\mathbb{T}(D)$ -eigenspaces of  $E_1(D, \psi_D)$  are 1-dimensional by (17), and so all the  $\vartheta_\chi$  with  $\chi^2 = 1$  are “primitive” as well.

We observe that it follows from (17) that

$$\Theta_{d_K}^E := \Theta_D \cap E_1(|d_K|, \psi_{d_K}) = E_1(|d_K|, \psi_{d_K}).$$

However, the analogous statement for

$$\Theta_{d_K}^S = \Theta_D \cap S_1(|d_K|, \psi_{d_K})$$

is in general not true, as the discussion of Serre [28], §9, and/or [17] shows. In particular, the table on p. 258 of [28] shows that we have  $\Theta_{-p}^S \neq S_1(p, \psi_{-p})$  for  $p = 139, 163, 211, 227, 283$ .

We are now ready to prove Theorems 1 and 2 of the introduction.

**Proof of Theorem 1.** By Proposition 9 and Theorem 12 we know that the subspace  $\Theta_D$  of  $M_1(|D|, \psi_D)$  has the basis  $\{\vartheta_\chi\}$  which consists of  $\mathbb{T}(D)$ -eigenfunctions, and so it follows that  $\Theta_D$  is a  $\mathbb{T}(D)$ -submodule.

To verify that  $\Theta_D$  has multiplicity one, we have to show that  $\vartheta_{\chi_1}$  and  $\vartheta_{\chi_2}$  belong to different  $\mathbb{T}(D)$ -eigenspaces whenever  $\vartheta_{\chi_1} \neq \vartheta_{\chi_2}$ , i.e., whenever  $\chi_1 \notin \{\chi_2, \chi_2^{-1}\}$ .

For this, we first note that there is an  $f \in Q_D$  such that  $\chi_1(f) \neq \chi_2(f), \bar{\chi}_2(f)$ . Indeed, the hypothesis on  $\chi_1, \chi_2$  implies that  $\exists f_i \in Q_D$  such that  $\chi_1(f_1) \neq \chi_2(f_1)$  and  $\chi_1(f_2) \neq \chi_2^{-1}(f_2) = \bar{\chi}_2(f_2)$ . Now if  $\chi_1(f_1) \neq \bar{\chi}_2(f_1)$ , then we can take  $f = f_1$  and if  $\chi_1(f_2) \neq \chi_2(f_2)$ , then we can take  $f = f_2$ . If neither of these cases holds, then we can take  $f \sim f_1 f_2$  (product in  $\text{Cl}(D)$ ) because here we have that  $\chi_1(f_1) = \bar{\chi}_2(f_1)$  and  $\chi_1(f_2) = \chi_2(f_2)$ , and so

$$\chi_1(f) = \chi_1(f_1)\chi_1(f_2) \neq \chi_2(f_1)\chi_1(f_2) = \chi_2(f_1)\chi_2(f_2) = \chi_2(f),$$

and

$$\chi_1(f) = \chi_1(f_1)\chi_1(f_2) \neq \bar{\chi}_2(f_1)\chi_1(f_2) = \bar{\chi}_2(f_1)\bar{\chi}_2(f_2) = \bar{\chi}_2(f).$$

With  $f$  as above, choose a prime  $p \nmid D$  such that  $r_p(f) > 0$ ; cf. Lemma 8(a). Then by (7) and Lemma 8(b) we see that

$$a_p(\chi_i) = \frac{w(f)}{w_D} \text{Re}(\chi_i(f))r_p(f), \quad \text{for } i = 1, 2.$$

Since  $\chi_1(f) \neq \chi_2(f), \bar{\chi}_2(f)$  and  $|\chi_i(f)| = 1$ , it follows that  $\text{Re}(\chi_1(f)) \neq \text{Re}(\chi_2(f))$ , and so  $a_p(\chi_1) \neq a_p(\chi_2)$ . Since  $a_p(\chi_i)$  is the  $T_p$ -eigenvalue of  $\vartheta_{\chi_i}$  by Theorem 12, we see that  $\vartheta_{\chi_1}$  and  $\vartheta_{\chi_2}$  lie in distinct  $\mathbb{T}(D)$ -eigenspaces, so  $\Theta_D$  has multiplicity one.

The last assertion of Theorem 1 follows immediately from Theorem 14.  $\square$

**Proof of Theorem 2.** Let  $V_E$  and  $V_S$  denote the  $\mathbb{C}$ -subspace of  $\Theta_D$  generated by  $\{\vartheta_\chi : \chi^2 = 1\}$  and by  $\{\vartheta_\chi : \chi^2 \neq 1\}$ , respectively. By Theorem 14 we know that  $V_E \subset \Theta_D^E$  and  $V_S \subset \Theta_D^S$ , and by Proposition 9 we know that  $V_E + V_S = \Theta_D$ . Since  $\Theta_D^E \cap \Theta_D^S = \{0\}$  by (12), it follows that  $V_E = \Theta_D^E$  and  $V_S = \Theta_D^S$  and that hence  $\Theta_D = \Theta_D^E \oplus \Theta_D^S$ .

Let  $\text{Cl}(D)^*[2] = \{ \chi \in \text{Cl}(D)^* : \chi^2 = 1 \}$  denote the group of quadratic characters. By Proposition 9 we know that the set  $\{ \vartheta_\chi : \chi \in \text{Cl}(D)^*[2] \}$  is linearly independent and hence is a basis of  $\Theta_D^E = V_E$ . Thus

$$\dim(\Theta_D^E) = |\text{Cl}(D)^*[2]| = [\text{Cl}(D) : \text{Cl}(D)^2] = g_D,$$

and hence

$$\dim(\Theta_D^S) = \dim(\Theta_D) - \dim(\Theta_D^E) = \bar{h}_D - g_D = \frac{1}{2}(h_D - g_D),$$

the latter by Proposition 7 and equation (5), respectively. This proves (1). □

**Remark 17.** (a) For later reference we note that the above proof of Theorem 2 shows more precisely that

$$\{ \vartheta_\chi : \chi \in \text{Cl}(D)^*[2] \} \text{ is a basis of } \Theta_D^E,$$

and that

$$\{ \vartheta_\chi : \chi \in \overline{\text{Cl}(D)}^*, \chi^2 \neq 1 \} \text{ is a basis of } \Theta_D^S.$$

(b) It follows immediately from (1) that

$$(18) \quad \Theta_D \subset E_1(|D|, \psi_D) \iff \Theta_D^S = \{0\} \iff h_D = g_D.$$

The discriminants  $D < 0$  which satisfy the last condition (or, equivalently, the condition that  $\text{Cl}(D)$  is an elementary abelian 2-group) are called *idoneal discriminants* because a number  $n \geq 1$  is *idoneal* (in the sense of Euler) if and only if  $-4n$  is an idoneal discriminant; cf. [16] for a recent survey about idoneal numbers.

The above assertion (18) can be viewed as an alternate version of Theorem 3 of Kitaoka [18], which states that

$$(19) \quad \vartheta_{1_D} \in E_1(|D|, \psi_D) \iff h_D = g_D.$$

Indeed, if  $h_D = g_D$ , then clearly  $\vartheta_{1_D} \in \theta_D \subset E_1(|D|, \psi_D)$  by (18). To prove the converse, note first that by (8) we have that

$$\vartheta_{1_D} = \frac{1}{w_D} \sum_{\chi \in \overline{\text{Cl}(D)}^*} w(\chi) \vartheta_\chi$$

(because  $\chi(1_D) = 1$  for all  $\chi \in \text{Cl}(D)^*$ ). Thus, by part (a) we see that

$$\vartheta_{1_D} \in E_1(|D|, \psi_D) \iff \chi^2 = 1, \forall \chi \in \text{Cl}(D)^* \iff \text{Cl}(D)^2 = \{1\} \iff h_D = g_D,$$

which proves (19).

The above results immediately imply the following corollary which generalizes the work of Siegel [29] on theta-series attached to forms in  $m = 2k > 4$  variables (cf. [29], particularly pp. 577–581) to the case of binary forms. Note that this gives an alternate proof (for binary forms) of Kitaoka’s extension of Siegel’s work to the case  $m \geq 2$ ; cf. [18], Lemma 1.

**Corollary 18.** (a) Let  $f_1, f_2 \in Q_D$  be two primitive forms. Then  $\vartheta_{f_1} - \vartheta_{f_2}$  is a cusp form if and only if  $f_1$  and  $f_2$  are genus-equivalent, i.e.,

$$(20) \quad \vartheta_{f_1} - \vartheta_{f_2} \in S_1(|D|, \psi_D) \iff \chi(f_1) = \chi(f_2), \forall \chi \in \text{Cl}(D)^*[2] \iff f_1^{-1} f_2 \in \text{Cl}(D)^2.$$

(b) For any  $f \in Q_D$  we have that

$$(21) \quad F(z, f) := \frac{g_D}{h_D} \sum_{f_1 \in \text{Cl}(D)^2} \vartheta_{ff_1}(z) = \frac{w_D}{h_D} \sum_{\chi \in \text{Cl}(D)^*[2]} \chi(f) \vartheta_\chi(z) \in \Theta_D^E.$$

Thus,  $F(z, f) \in \Theta_D^E$  is the Eisenstein component of  $\vartheta_f(z)$ , and  $\vartheta_f(z) - F(z, f) \in \Theta_D^S$  is its cuspidal component.

**Proof.** (a) By (8) we have

$$\vartheta_{f_1} - \vartheta_{f_2} = \frac{h_D}{w_D} \sum_{\chi \in \text{Cl}(D)^*} w(\chi) \text{Re}(\chi(f_1) - \chi(f_2)) \vartheta_\chi,$$

and so it follows from Remark 17(a) that

$$\begin{aligned} \vartheta_{f_1} - \vartheta_{f_2} \in S_1(|D|, \psi_D) &\iff \text{Re}(\chi(f_1) - \chi(f_2)) = 0, \forall \chi \in \text{Cl}(D)^*[2] \\ &\iff \chi(f_1) = \chi(f_2), \forall \chi \in \text{Cl}(D)^*[2] \end{aligned}$$

because all  $\chi \in \text{Cl}(D)^*[2]$  are real-valued. This proves the first equivalence of (20) and hence also (20) because the second equivalence is obvious.

(b) By (8) we have that

$$F(z, f) = \frac{g_D w_D}{h_D^2} \sum_{\chi \in \text{Cl}(D)^*} c(f, \chi) \vartheta_\chi(z)$$

with

$$c(f, \chi) = \sum_{f_1 \in \text{Cl}(D)^2} \bar{\chi}(ff_1) = \bar{\chi}(f) \sum_{f_1 \in \text{Cl}(D)^2} \bar{\chi}(f_1) = \begin{cases} \bar{\chi}(f) \frac{h_D}{g_D} & \text{if } \chi^2 = 1, \\ 0 & \text{otherwise} \end{cases}$$

where the last identity follows from a suitable orthogonality relation and the fact that  $|\text{Cl}(D)^2| = \frac{h_D}{g_D}$ . From this the asserted identity (21) follows immediately.

By (21) and Theorem 14 it is clear that  $F(z, f) \in \Theta_D^E$ . Moreover, by (8) and (21) we have that

$$\vartheta_f(z) - F(z, f) = \frac{w_D}{h_D} \sum_{\chi^2 \neq 1} \bar{\chi}(f) \vartheta_\chi(z) \in \Theta_D^S,$$

the latter by Theorem 14 again. This proves the last assertion.  $\square$

**Remark 19.** The Fourier coefficients  $a_n(F(z, f))$  of the Eisenstein series  $F(z, f)$  are easily determined, at least when  $(n, D) = 1$ . Indeed, we have

$$(22) \quad a_n(F(z, f)) = \frac{w_D g_D}{h_D} \varepsilon(f, n) \# Id_n(\mathfrak{D}_K),$$

where  $\varepsilon(f, n) = 1$  if  $n \equiv f(x, y) \pmod{|D|}$ , for some  $x, y \in \mathbb{Z}$ , and  $\varepsilon(f, n) = 0$  otherwise. To see this, note first that

$$a_n(F(z, f)) = \frac{w_D}{h_D} \sum_{\chi \in \text{Cl}(D)^*[2]} \chi(f) a_n(\chi)$$

by (21). Thus  $a_n(F(z, f)) = 0 = \varepsilon(f, n)$ , if  $Id_n(\mathfrak{D}_K) = \emptyset$  (cf. (9)), so assume that  $\exists \mathfrak{a}_n \in Id_n(\mathfrak{D}_K)$ . If  $f_n \in Q_D$  is such that  $\lambda_D(f_n) = \varphi_D(\mathfrak{a}_n)$ , then there exist  $x, y \in \mathbb{Z}$  such that  $f(x, y) = n$ . Now for any  $\chi \in Cl(D)^*[2]$  we have by (16) that

$$a_n(\chi) = \tilde{\chi}(\mathfrak{a}_n) \# Id_n(\mathfrak{D}_K) = \chi(f_n) \# Id_n(\mathfrak{D}_K),$$

so we obtain that

$$a_n(F(z, f)) = \frac{w_D}{h_D} \# Id_n(\mathfrak{D}_K) \sum_{\chi \in Cl(D)^*[2]} \chi(ff_n).$$

By an orthogonality relation we have that  $\sum_{\chi^2=1} \chi(ff_n) = |Cl(D)^*[2]| = g_D$  if  $ff_n \in Cl(D)^2$  and equals 0 otherwise. By genus theory (cf. [6], §3.B), this means that this sum equals  $g_D \varepsilon(f, n)$ , and so (22) follows.

As another application of the above results we compute the trace  $\text{tr}(T_n|V)$  of the Hecke operator  $T_n$  on the vector spaces  $V = \Theta_D^E$  and  $V = \Theta_D^S$ .

**Corollary 20.** *If  $(n, f_D) = 1$ , then the traces of the Hecke operator  $T_n$  on the spaces  $\Theta_D^E$  and  $\Theta_D^S$  are given by*

$$(23) \quad \begin{cases} \text{tr}(T_n|\Theta_D^E) &= \frac{h_D}{w_D} a_n(F(z, 1_D)) \\ \text{tr}(T_n|\Theta_D^S) &= \frac{h_D}{2w_D} (r_n(1_D) - a_n(F(z, 1_D))). \end{cases}$$

**Proof.** By Remark 17(a), Theorem 12, and (21) we have that

$$\text{tr}(T_n|\Theta_D^E) = \sum_{\chi \in Cl(D)^*[2]} a_n(\chi) = \frac{h_D}{w_D} a_n(F(z, 1_D)).$$

This clearly proves the first equation of (23). In order to prove the second equation, put  $X = \{ \chi \in Cl(D)^* : \chi^2 \neq 1 \}$ . Then by Remark 17(a) and Theorem 12 we have

$$\text{tr}(T_n|\Theta_D^S) = \sum_{\chi \in X} a_n(\chi) = \frac{1}{2} \sum_{\chi^2 \neq 1} a_n(\chi) = \frac{1}{2} \sum_{\chi} a_n(\chi) - \frac{1}{2} \sum_{\chi^2=1} a_n(\chi).$$

Since the first sum equals  $\frac{h_D}{w_D} r_n(1_D)$  by (8) and since (as above) the second sum equals  $\frac{h_D}{w_D} a_n(F(z, 1_D))$  by (21), the formula (23) follows. □

### 4. Primitive characters and primitive forms

We now want to study the relation between the theta-series  $\vartheta_\chi$  and the modular form  $f(z; \tilde{\chi})$  attached to the associated Hecke character  $\tilde{\chi} = \chi \circ \lambda_D^{-1} \circ \varphi_D$  in more detail. For this, it is useful to introduce the following terminology.

**Definition.** If  $\chi \in Cl(D)^*$  is a character on the class group  $Cl(D)$ , then we say that  $\chi$  is *primitive* if we have that  $\text{Ker}(\bar{\pi}_{D, D/c^2}) \not\subset \text{Ker}(\chi)$ , for all divisors  $c \mid f_D$  with  $c > 1$ . Here  $\bar{\pi}_{D, D/c^2} : Cl(D) \rightarrow Cl(D/c^2)$  is the homomorphism induced by the map  $\bar{\pi}_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}} : Cl(\mathfrak{D}_D) \rightarrow Cl(\mathfrak{D}_{D/c^2})$  which is defined in Remark 39 of the Appendix; in other words,  $\bar{\pi}_{D, D'} = \lambda_{D'}^{-1} \circ \bar{\pi}_{\mathfrak{D}_D, \mathfrak{D}_{D'}} \circ \lambda_D$ , for  $D' = D/c^2$ .

Moreover, the conductor  $f_\chi$  of  $\chi \in \text{Cl}(D)^*$  is defined by

$$f_\chi = \gcd(f : f \mid f_D \text{ and } \text{Ker}(\bar{\pi}_{D, f^2 d_K}) \subset \text{Ker}(\chi)).$$

Thus, if  $\chi \in \text{Cl}(D)^*$  is primitive, then clearly  $f_\chi = f_D$ . Moreover, the converse is also true, as the following result shows.

**Proposition 21.** *Let  $\chi \in \text{Cl}(D)^*$  be a character with conductor  $f_\chi$ , and let  $f \mid f_D$ . Then*

$$(24) \quad \text{Ker}(\bar{\pi}_{D, f^2 d_K}) \subset \text{Ker}(\chi) \iff f_\chi \mid f.$$

Thus, if  $D_\chi := f_\chi^2 d_K$ , then  $\chi = \chi_{pr} \circ \bar{\pi}_{D, D_\chi}$ , for a unique character  $\chi_{pr} \in \text{Cl}(D_\chi)^*$ , and  $\chi_{pr}$  is primitive.

**Proof.** From the definition of  $D_\chi$  and (49) we have that  $\mathfrak{D}_{D_\chi} = \prod_f \mathfrak{D}_{f^2 d_K}$ , where the product is over all  $f \mid f_D$  with  $\text{Ker}(\bar{\pi}_{D, f^2 d_K}) \subset \text{Ker}(\chi)$ . It thus follows from Corollary 41 that  $\text{Ker}(\bar{\pi}_{D, D_\chi}) \subset \text{Ker}(\chi)$ , and so the assertion (24) is obvious. Moreover, since  $\bar{\pi}_{D, D_\chi} : \text{Cl}(D) \rightarrow \text{Cl}(D_\chi)$  is surjective (cf. Proposition 38), there is a unique character  $\chi_{pr}$  on  $\text{Cl}(D_\chi)$  such that  $\chi = \chi_{pr} \circ \bar{\pi}_{D, D_\chi}$ . Finally,  $\chi_{pr}$  is primitive because if  $\text{Ker}(\bar{\pi}_{D_\chi, D_\chi/c^2}) \subset \text{Ker}(\chi_{pr})$ , for some  $c \mid f_{D_\chi}$ , then

$$\text{Ker}(\bar{\pi}_{D, (f_\chi/c)^2 d_K}) = \bar{\pi}_{D, D_\chi}^{-1}(\text{Ker}(\bar{\pi}_{D_\chi, D_\chi/c^2})) \subset \bar{\pi}_{D, D_\chi}^{-1}(\text{Ker}(\chi_{pr})) = \text{Ker}(\chi),$$

so  $f_\chi \mid \frac{f_\chi}{c}$  by (24) and hence  $c = 1$ , i.e.,  $\chi_{pr}$  is primitive.  $\square$

Using the results of the Appendix, we can now compute the Fourier coefficients  $a_n(\chi)$  of  $\vartheta_\chi$  in the case that  $n \mid f_D^2$ . This is a *key result* that will be used in the next section to determine the local factors of the  $L$ -series  $L(s, \vartheta_\chi)$ .

**Proposition 22.** *Let  $\chi \in \text{Cl}(D)^*$  be a character on  $\text{Cl}(D)$  with conductor  $f_\chi$ , and put  $\bar{f}_\chi = f_D/f_\chi$ . If  $n \mid f_D^2$ , then*

$$(25) \quad a_n(\chi) = \begin{cases} c \prod_{p \mid c} \left(1 - \frac{1}{p} \psi_{D/c^2}(p)\right) & \text{if } n = c^2 \text{ and } c \mid \bar{f}_\chi, \\ 0 & \text{otherwise.} \end{cases}$$

**Proof.** If  $n$  is not a square, then  $\text{Id}_n(\mathfrak{D}_D) = \emptyset$  by Proposition 48, so by (9) we have that  $a_n(\chi) = 0$  in this case. Thus, assume that  $n = c^2$ , and put  $\chi^* = \chi \circ \lambda_D^{-1}$ . Then it follows from (9) and Proposition 48 that

$$a_n(\chi) = \sum_{\mathfrak{a} \in c \text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}})} \chi^*(\mathfrak{a}) = \sum_{\mathfrak{a} \in \text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}})} \chi^*(\mathfrak{a}).$$

Now from (52) and (24) we see that

$$\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}})P(\mathfrak{D}_D) \leq \text{Ker}(\chi^*) \iff f_\chi \mid \frac{f_D}{c} \iff c \mid \bar{f}_\chi.$$

If this is not the case, i.e., if  $\chi^*$  is non-trivial on  $\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}})$ , then this sum equals 0 by an orthogonality relation. On the other hand, if  $c \mid \bar{f}_\chi$ , then from the above discussion we see that  $a_n(\chi) = |\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}})|$  and so the assertion follows by using (54).  $\square$

We now turn our attention to the modular form  $f(z; \tilde{\chi})$ , where, as before,  $\tilde{\chi}$  is defined by  $\tilde{\chi} = \chi \circ \lambda_D^{-1} \circ \varphi_D$ . To determine its properties, we need to know the conductor  $\text{cond}(\tilde{\chi})$  of the Hecke character  $\tilde{\chi}$ ; here  $\text{cond}(\tilde{\chi}) \in \text{Id}(\mathfrak{O}_K)$  is as defined on p. 91 of [22]. The following result is due to Hasse[10]:

**Theorem 23.** *The conductor of the Hecke character  $\tilde{\chi} = \chi \circ \lambda_D^{-1} \circ \varphi_D$  is*

$$\text{cond}(\tilde{\chi}) = f_\chi \mathfrak{O}_K .$$

*In particular,  $\chi$  is primitive if and only if  $\tilde{\chi}$  is a primitive Hecke character.*

**Proof.** The first assertion is Satz 2 of Hasse[10], and the second clearly follows from the first. □

From the above theorem it follows easily that if  $\chi \in \text{Cl}(D)^*$  is a primitive character, then  $f(z; \tilde{\chi})$  is a *primitive form* in the sense of the following definition.

**Definition.** A modular form  $f \in M_k(N, \psi)$  is called a *primitive form* if either  $f \in S_k(N, \psi)$  is a normalized newform of some level  $M|N$  (so  $f$  is a primitive form in the sense of [22], §4.6) or if  $f = f_k(z; \psi_1, \psi_2)$  is one of the Eisenstein series defined on p. 178 of [22].

**Remark 24.** It follows from the theory of newforms (cf. [22], §4.6) and the theory of Eisenstein forms (cf. [22], §4.7) that each  $\mathbb{T}(N)$ -eigenspace of  $M_k(N, \psi)$  contains a unique primitive form. Thus, there is natural bijection between primitive forms and  $\mathbb{T}(N)$ -eigenspaces of  $M_k(N, \psi)$ .

**Corollary 25.** *If  $\chi \in \text{Cl}(D)^*$  is a primitive character, then  $f(z; \tilde{\chi})$  is a primitive form of level  $|D|$ .*

**Proof.** Suppose first that  $\chi^2 \neq 1$ . Then from the proof of Theorem 14 we know that  $\tilde{\chi} \neq \chi' \circ N_K$  for all Dirichlet characters  $\chi'$ , and by Theorem 23 we know that  $\tilde{\chi}$  is a primitive Hecke character. It thus follows from Theorem 4.8.2 of [22] that  $f(z; \tilde{\chi})$  is a primitive cusp form (of level  $|D|$ ).

Now suppose that  $\chi^2 = 1$ . Since  $\tilde{\chi}$  is a primitive Hecke character by Theorem 23 and since  $f(z; \tilde{\chi})$  is not a cusp form by (the proof of) Theorem 14, it follows from the last part of the proof of Theorem 4.8.2 of [22] that there exist Dirichlet characters  $\chi_1, \chi_2$  with  $\chi_1 \chi_2 = \psi_D$  such that  $L(s, f(z; \tilde{\chi})) = L(s, \chi_1)L(s, \chi_2)$ . Since the latter equals  $L(s, f_1(z; \chi_1, \chi_2))$  by [22], Theorem 4.7.1, we conclude that  $f(z; \tilde{\chi}) = f_1(z; \chi_1, \chi_2)$ , and hence  $f(z; \tilde{\chi})$  is primitive of level  $|D|$ . □

**Remark 26.** If  $\chi \in \text{Cl}(D)^*[2]$  is a primitive *quadratic* character, then by combining the above proof with that of Theorem 14, we obtain the formula

$$(26) \quad f(z; \tilde{\chi}) = f_1(z; \psi_{D_1}, \psi_{D_2}), \quad \text{where } \tilde{\chi} = \psi_{D_1} \circ N_K = \psi_{D_2} \circ N_K,$$

and  $D = D_1 D_2$  is a suitable factorization of  $D$  into fundamental discriminants. Indeed, by the proof of Corollary 25 we know that (26) holds for some pair of characters  $\chi_1, \chi_2$ , and the proof of Theorem 14, particularly equations (15) and (16), show that  $\chi_i = \psi_{D_i}$  for  $i = 1, 2$ , where  $D = D_1 D_2$  is the fundamental factorization associated to  $\chi$ .

We can now prove Theorem 3 of the introduction.

**Proof of Theorem 3.** (a) This follows immediately from Proposition 21.

(b) Since  $\chi_{pr} \in \text{Cl}(D_\chi)^*$  is primitive by part (a), it follows from Corollary 25 that  $f(z; \tilde{\chi}_{pr}) \in M_1(|D_\chi|, \psi_{D_\chi})$  is a primitive form of level  $|D_\chi|$ . By Theorem 12 and (14) we know that  $\vartheta_{\chi_{pr}}$  is a normalized  $\mathbb{T}(D_\chi)$ -eigenfunction which has the same eigenvalues as  $f(z; \tilde{\chi}_{pr})$ . Since  $f(z; \tilde{\chi}_{pr})$  is primitive of level  $|D_\chi|$ , its associated eigenspace in  $M_1(|D_\chi|, \psi_{D_\chi})$  is one-dimensional (cf. [22], Theorems 4.6.12 and 4.7.2) and so

$$(27) \quad \vartheta_{\chi_{pr}}(z) = f(z; \tilde{\chi}_{pr})$$

because both forms are normalized. Thus  $\vartheta_{\chi_{pr}}$  is primitive and  $L(s, \vartheta_{pr}) = L(s, \tilde{\chi}_{pr})$ .

Now by Theorem 12 we know that  $\vartheta_\chi$  is a  $\mathbb{T}(D)$ -eigenfunction which lies in the eigenspace defined by the primitive form  $\vartheta_{\chi_{pr}}(z) = f(z; \tilde{\chi}_{pr})$  because by (14) we have that  $\vartheta_\chi \sim_{f_D} f(z; \tilde{\chi}) \sim_{f_D} f(z; \tilde{\chi}_{pr})$ . Since  $D/D_\chi = (f_D/f_\chi)^2 = \bar{f}_\chi^2$ , it follows from the description of the  $\mathbb{T}(D)$ -eigenspaces of  $M_1(|D|, \psi_D)$  given in Corollary 4.6.20 and (implicitly) in Theorem 4.7.2 of [22], that there exist constants  $c_n(\chi) \in \mathbb{C}$  with  $n \mid \bar{f}_\chi^2$  such that (2) holds. Thus, if we put  $c_n(\chi) = 0$  when  $n \nmid \bar{f}_\chi^2$ , then we have

$$(28) \quad a_m(\chi) = \sum_{\substack{n \mid \bar{f}_\chi^2 \\ n \mid m}} c_n(\chi) a_{m/n}(\chi_{pr}) = \sum_{n \mid m} c_n(\chi) a_{m/n}(\chi_{pr}), \quad \text{for all } m \geq 1,$$

and so it follows that  $L(s, \vartheta_\chi) = (\sum_{n \geq 1} c_n(\chi) n^{-s}) L(s, \vartheta_{\chi_{pr}})$ ; cf. [11], Theorem 284. Thus (3) holds. Moreover, it follows that  $n \mapsto c_n(\tilde{\chi})$  is multiplicative because both  $L(s, \vartheta_\chi)$  and  $L(s, \vartheta_{\chi_{pr}})$  have Euler products by Theorem 12, and hence so does

$$(29) \quad C(s, \chi) := \sum_{n \geq 1} \frac{c_n(\chi)}{n^s} = \frac{L(s, \vartheta_\chi)}{L(s, \vartheta_{\chi_{pr}})} = \frac{L(s, \vartheta_\chi)}{L(s, \tilde{\chi}_{pr})}.$$

This means that  $n \mapsto c_n(\chi)$  is multiplicative.

Finally, we note that it follows from (29) that all the  $c_n(\chi)$  are real because the Fourier coefficients  $a_n(\chi)$  and  $a_n(\chi_{pr})$  are real; cf. (7).  $\square$

## 5. The Dirichlet series $C(s, \chi)$

In the previous section we had seen that each theta-series  $\vartheta_\chi$  can be expressed as a linear combination of “shifted” modular forms  $f(nz; \tilde{\chi}_{pr}) = \vartheta_{\chi_{pr}}(nz)$  associated to the primitive Hecke character  $\tilde{\chi}_{pr}$ ; cf. Theorem 3. We now want to obtain precise formulae for the coefficients  $c_n(\chi)$  of this linear combination. We thus study the (finite) Dirichlet series  $C(s, \chi)$  of equation (29) in more detail.

As in Hardy and Wright [11], §17.4, we let  $F_p(s)$  denote the  $p$ -Euler factor of a Dirichlet series  $F(s)$  at the prime  $p$ , i.e.,

$$F_p(s) = \sum_{k \geq 0} a_{p^k} p^{-ks}, \quad \text{if } F(s) = \sum_{n \geq 1} a_n n^{-s}.$$

The following proposition gives the relation between the  $p$ -Euler factors  $C_p(s, \chi)$  and  $L_p(s, \chi)$  of the Dirichlet series  $C(s, \chi)$  and  $L(s, \chi) := L(s, \vartheta_\chi)$ .

**Proposition 27.** *If  $\chi \in \text{Cl}(D)^*$ , then for any prime  $p$  we have that*

$$(30) \quad C_p(s, \chi) = L_p(s, \chi)(1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s}).$$

**Proof.** By taking  $m = p^k$  (for  $k \geq 0$ ) in (28) we see that

$$L_p(s, \chi) = C_p(s, \chi)L_p(s, \chi_{pr}),$$

and so (30) follows because

$$(31) \quad L_p(s, \chi_{pr}) = (1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s})^{-1}.$$

Indeed, since  $\vartheta_{\chi_{pr}} = f(z; \tilde{\chi}_{pr})$  is a primitive form of level  $|D_\chi|$  by Theorem 3(b), equation (31) follows from Corollary 4.6.22 and (4.7.16) of [22].  $\square$

We are now ready to determine  $C(s, \chi)$  explicitly.

**Theorem 28.** *Let  $\chi \in \text{Cl}(D)^*$  be a character on  $\text{Cl}(D)$  of conductor  $f_\chi|f_D$ , and let  $\chi_{pr} \in \text{Cl}(D_\chi)$  be the associated primitive character on  $\text{Cl}(D_\chi)$ , where  $D_\chi = f_\chi^2 d_K$ . Then*

$$(32) \quad C(s, \chi) = \prod_{p|\bar{f}_\chi} C_p(s, \chi),$$

where  $\bar{f}_\chi = f_D/f_\chi$ . Moreover, if  $p|\bar{f}_\chi$  and if  $p^{\bar{e}_p}||\bar{f}_\chi$  is the highest power of  $p$  dividing  $\bar{f}_\chi$ , then

$$(33) \quad C_p(s, \chi) = \frac{1 - p^{(1-2s)\bar{e}_p}}{1 - p^{1-2s}} \left( 1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s} \right) + \left( 1 - \frac{1}{p}\psi_{D_\chi}(p) \right) p^{(1-2s)\bar{e}_p};$$

in other words, we have for  $k \geq 1$  that

$$(34) \quad c_{p^k}(\chi) = \begin{cases} -a_p(\chi_{pr})p^{(k-1)/2} & \text{if } k \equiv 1 \pmod{2} \text{ and } k < 2\bar{e}_p, \\ p^{k/2} + p^{k/2-1}\psi_{D_\chi}(p) & \text{if } k \equiv 0 \pmod{2} \text{ and } k < 2\bar{e}_p, \\ p^{\bar{e}_p} & \text{if } k = 2\bar{e}_p, \\ 0 & \text{if } k > 2\bar{e}_p. \end{cases}$$

**Proof.** The first assertion (32) follows immediately from the multiplicativity of the  $c_n(\chi)$ 's; cf. Theorem 3(b). To prove (33), we first observe that if  $0 \leq k \leq 2\bar{e}_p$ , then we obtain from (25) that

$$(35) \quad a_{p^k}(\chi) = \begin{cases} 0 & \text{if } k \equiv 1 \pmod{2}, \\ p^{k/2} & \text{if } k \equiv 0 \pmod{2} \text{ and } k < 2\bar{e}_p, \\ p^{\bar{e}_p} \left( 1 - \frac{1}{p}\psi_{D_\chi}(p) \right) & \text{if } k = 2\bar{e}_p. \end{cases}$$

Indeed, if  $k$  is odd, then this is clear, so assume that  $k$  is even. Then (25) yields that  $a_{p^k}(\chi) = p^{k/2} \left( 1 - \frac{1}{p}\psi_{D/p^k}(p) \right)$ . Now if  $k < 2\bar{e}_p$ , then  $p|\frac{D}{p^k}$  because  $p^{2\bar{e}_p}|D$ . Thus  $\psi_{D/p^k}(p) = 0$ , and hence (35) holds in this case. On the other hand, if  $k = 2\bar{e}_p$ , then

$\frac{D}{p^k} = D_\chi f_1^2$ , where  $f_1 = \bar{f}_\chi p^{-\bar{e}_p}$ . Since  $(f_1, p) = 1$ , we have that  $\psi_{D/p^k}(p) = \psi_{D_\chi}$ , and so (35) holds in all cases.

Note that by using the identity

$$\sum_{k=0}^{\bar{e}_p-1} p^k X^2 = \frac{1 - (pX^2)^{\bar{e}_p}}{1 - pX^2}$$

we can re-write (35) in the form

$$(36) \quad \sum_{k=0}^{2\bar{e}_p} a_{p^k}(\chi) X^k = \frac{1 - (pX^2)^{\bar{e}_p}}{1 - pX^2} + \left(1 - \frac{1}{p} \psi_{D_\chi}(p)\right) (pX^2)^{\bar{e}_p}.$$

Thus, viewing (30) as an identity of power series in  $X = p^{-s}$ , it follows from (30) and (36) that

$$C_p(s, \chi) \equiv \frac{1 - (pX^2)^{\bar{e}_p}}{1 - pX^2} \left(1 - a_p(\chi_{pr})X + \psi_{D_\chi}(p)X^2\right) + \left(1 - \frac{1}{p} \psi_{D_\chi}(p)\right) (pX^2)^{\bar{e}_p} \pmod{X^{2\bar{e}_p+1}}.$$

Now since  $C_p(s, \chi)$  is a polynomial of degree  $\leq 2\bar{e}_p$  (because  $c_{p^k}(\chi) = 0$  by definition when  $k > 2\bar{e}_p$ ), and since the same is true for the right hand side, it follows that these two polynomials are equal, and so (33) follows (by replacing  $X$  by  $p^{-s}$ ). The last assertion (34) follows immediately from (33) by comparing the coefficients in  $X = p^{-s}$ .  $\square$

**Corollary 29.** *In the situation of Theorem 28 we have that*

$$(37) \quad c_{\bar{f}_\chi^2}(\chi) = \bar{f}_\chi.$$

*Thus,  $\vartheta_\chi$  has exact level  $|D|$ , and  $C(s, \chi) = 1$  if and only if  $\chi$  is primitive. In particular,  $\vartheta_\chi$  is a primitive form if and only if  $\chi$  is a primitive character.*

**Proof.** Since  $\bar{f}_\chi = \prod_{p|\bar{f}_\chi} p^{\bar{e}_p}$  and  $c_n(\chi)$  is multiplicative, we have by (34) that

$$c_{\bar{f}_\chi^2}(\chi) = \prod_{p|\bar{f}_\chi} c_{p^{2\bar{e}_p}}(\chi) = \prod_{p|\bar{f}_\chi} p^{\bar{e}_p} = \bar{f}_\chi,$$

which proves (37). From this and (2) it follows that  $\vartheta_\chi$  has exact level  $|D|$ .

If  $\chi$  is primitive, i.e., if  $\bar{f}_\chi = 1$ , then clearly  $C(s, \chi) = 1$  by definition (cf. (3)). Conversely, if  $C(s, \chi) = 1$ , i.e., if  $c_n(\chi) = 0$  for all  $n > 1$ , then (37) forces that  $\bar{f}_\chi = 1$ , which means that  $\chi$  is primitive.

To prove the last assertion, recall first that if  $\chi = \chi_{pr}$  is a primitive character, then  $\vartheta_\chi$  is a primitive form by Theorem 3(b). Conversely, suppose that  $\vartheta_\chi$  is a primitive form. Then  $\vartheta_\chi = \vartheta_{\chi_{pr}}$  because both are in the same  $\mathbb{T}(D)$ -eigenspace of  $M_1(|D|, \psi_D)$ , and so  $C(s, \chi) = L(s, \chi)/L(s, \chi_{pr}) = L(s, \vartheta_\chi)/L(s, \vartheta_{\chi_{pr}}) = 1$ , which means that  $\chi$  is primitive.  $\square$

**Remark 30.** (a) It follows from equations (8) and (3) that if  $f \in Q_D$ , then its (Epstein) zeta-function  $Z_f(s)$  of Remark 6(b) can be written as a sum of the  $L$ -functions  $L(s, \chi) := L(s, \vartheta_\chi)$  and the Hecke  $L$ -functions  $L(s, \chi_{pr}) = L(s, \tilde{\chi}_{pr})$  in the following way:

$$(38) \quad Z_f(s) = \frac{w_D}{h_D} \sum_{\chi \in \text{Cl}(D)^*} \overline{\chi(f)} L(s, \chi) = \frac{w_D}{h_D} \sum_{\chi} \overline{\chi(f)} C(s, \chi) L(s, \chi_{pr});$$

this generalizes the well-known relation (cf. [7]) for  $Z_f$  when  $D = d_k$  is a fundamental discriminant. However, the above relation and Corollary 29 show that if  $D \neq d_K$  is not a fundamental discriminant, then  $Z_f$  is never a linear combination of the associated Hecke  $L$ -functions because in that case the factor  $C(s, 1)$  is not a constant since the trivial character  $\chi = 1$  is not primitive; cf. Example 35 below.

(b) Similarly (and equivalently), by combining (8) with (2), we obtain in view of (32) and (35) the following explicit expression of the theta series  $\vartheta_f$  in terms of the (extended) Atkin-Lehner basis:

$$(39) \quad \vartheta_f(z) = \frac{w_D}{h_D} \sum_{\chi \in \text{Cl}(D)^*} \overline{\chi(f)} \vartheta_\chi(z) = \frac{w_D}{h_D} \sum_{\chi} \overline{\chi(f)} \sum_{n|\bar{f}_\chi} \vartheta_{\chi_{pr}}(nz).$$

As was mentioned in the introduction, Theorem 28 yields immediately a formula for the  $L$ -function  $L(s, \chi) = L(s, \vartheta_\chi)$ .

**Corollary 31.** *If  $\chi \in \text{Cl}(D)^*$ , then the  $L$ -function  $L(s, \chi)$  of  $\vartheta_\chi$  has the Euler product*

$$(40) \quad L(s, \chi) = \prod_p L_p(s, \chi)$$

where for  $p \nmid \bar{f}_\chi$  the  $p$ -Euler factor  $L_p(s, \chi)$  is

$$(41) \quad \begin{aligned} L_p(s, \chi) &= \left(1 - a_p(\chi)p^{-s} + \psi_D(p)p^{-2s}\right)^{-1} \\ &= \left(1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s}\right)^{-1}, \end{aligned}$$

whereas for  $p \mid \bar{f}_\chi$  it is given by

$$(42) \quad L_p(s, \chi) = \frac{1 - p^{(1-2s)\bar{e}_p}}{1 - p^{1-2s}} + \frac{\left(1 - \frac{1}{p}\psi_{D_\chi}(p)\right)p^{(1-2s)\bar{e}_p}}{1 - a_p(\chi_{pr})p^{-s} + \psi_{D_\chi}(p)p^{-2s}}.$$

**Proof.** Since the  $a_n(\chi)$ 's are multiplicative (cf. Theorem 12), it follows that  $L(s, \chi)$  has an Euler product (40). Since  $C_p(s, \chi) = 1$  when  $p \nmid \bar{f}_\chi$ , it follows from (3) that  $L_p(s, \chi) = L_p(s, \chi_{pr})$ , so in particular  $a_p(\chi) = a_p(\chi_{pr})$ . Moreover, since  $D = D_\chi \bar{f}_\chi^2$ , we see that  $\psi_D(p) = \psi_{D_\chi}(p)$ , so the second equality of (41) holds, and hence the first equality follows from (31).

The formula (42) follows immediately from (30) and (33). □

**Remark 32.** (a) The above Corollary 31 shows that the second identity of the equation on the bottom of p. 280 of Lang [21] is incorrect when  $\chi$  is not a primitive character.

(b) An alternate way of writing formula (42) is follows: if  $p^{\bar{e}_p} || \bar{f}_\chi$ , then

$$a_{p^k}(\chi) = \begin{cases} 0 & \text{if } k \equiv 1 \pmod{2} \text{ and } k < 2\bar{e}_p, \\ p^{k/2} & \text{if } k \equiv 0 \pmod{2} \text{ and } k < 2\bar{e}_p, \\ p^{e_p} \left(1 - \frac{1}{p} \psi_{D_\chi}(p)\right) a_{p^{k-2\bar{e}_p}}(\chi_{pr}) & \text{if } k \geq 2\bar{e}_p. \end{cases}$$

(c) Corollary 31 can be viewed as a partial generalization of some of the main results of the articles of Sun and Williams [30, 31]. Indeed, for a negative discriminant  $D < 0$ , the function  $F(A, n)$  of [30] is essentially the same as the function  $a_n(\chi)$  above. More precisely, if we put  $\chi_A(K) = e^{2\pi i[K, A]}$ , where  $[K, A]$  (which depends on a choice of a “basis” of  $\text{Cl}(D)$ ) is as defined on p. 143 of [30], then a comparison of formula (6) with the formula for  $F(A, n)$  on p. 144 of [30] shows that

$$a_n(\chi_A) = F(A, n), \quad \forall A \in \text{Cl}(D), \quad n \geq 1.$$

In particular, Corollary 31 generalizes Theorem 5.3 of [31] (for  $D < 0$ ) to the non-cyclic case, and gives a succinct general formula for it (and also for the many special cases discussed in §8 of [30]). In addition, Corollary 31 generalizes (for  $D < 0$ ) Theorem 4.3 of [30] because  $a_n(\chi_{D,0}) = N(n, D)/w(D)$  in the notation of Example 35(b) and that of [30].

As another application of Theorem 28, we determine when  $\vartheta_\chi$  is an eigenfunction under the *full* Hecke algebra  $\mathbb{T}_{|D|} = \langle T_n : n \geq 1 \rangle$  of level  $|D|$ .

**Corollary 33.** *If  $\chi \in \text{Cl}(D)^*$ , then  $\vartheta_\chi$  is a  $\mathbb{T}_{|D|}$ -eigenfunction if and only if  $\chi$  is primitive.*

**Proof.** If  $\chi = \chi_{pr}$  is primitive, then by (31) the  $L$ -function of  $\vartheta_\chi$  has the Euler product

$$(43) \quad L(s, \chi) = \prod_p \left(1 - a_p(\chi)p^{-1} + \psi_D(p)p^{-2s}\right)^{-1}.$$

Thus, by a theorem of Hecke (*cf.* [22], Theorem 4.5.16), it follows that  $\vartheta_\chi$  is a  $\mathbb{T}_{|D|}$ -eigenfunction.

Conversely, suppose that  $\vartheta_\chi$  is a  $\mathbb{T}_{|D|}$ -eigenfunction. Then by Hecke’s Theorem again, we know that its  $L$ -function  $L(s, \chi)$  has an Euler product (43). If  $\chi$  is not primitive, then  $\bar{f}_\chi = f_D/f_\chi > 1$ , so there exists a prime  $p | \bar{f}_\chi$ . Then by (35) we know that  $a_p(\chi) = 0$ , so the  $p$ -Euler factor of  $\vartheta_\chi$  is trivial, *i.e.*,  $L_p(s, \chi) = 1$ . (Recall that  $p | \bar{f}_\chi | D$ .) Thus by (30) we see that  $C_p(s, \chi) = 1 - a_p(\chi_{pr})p^{-1} + \psi_{D_\chi}(p)p^{-2s}$ , *i.e.*, that

$$c_p(\chi) = -a_p(\chi), \quad c_{p^2}(\chi) = \psi_{D_\chi}(p) \quad \text{and} \quad c_{p^k}(\chi) = 0, \quad \forall k > 2.$$

Comparing this to (34) shows that  $\bar{e}_p \leq 1$ . But if  $\bar{e}_p = 1$ , then by (34)

$$c_{p^2}(\chi) = p \neq \psi_{D_\chi}(p),$$

a contradiction. Thus  $\bar{f}_\chi = 1$ , and hence  $\chi$  is primitive. □

The above corollary allows us to determine when the spaces  $\Theta_D$ ,  $\Theta_D^E$  and  $\Theta_D^S$  are  $\mathbb{T}_{|D|}$ -modules.

**Proposition 34.** (a) *The space  $\Theta_D$  is a  $\mathbb{T}_{|D|}$ -module if and only if its Eisenstein space  $\Theta_D^E$  is a  $\mathbb{T}_{|D|}$ -module if and only if  $D$  is a fundamental discriminant.*

(b) *The space  $\Theta_D^S$  of cusp forms is a  $\mathbb{T}_{|D|}$ -module if and only if every proper discriminant divisor  $D'$  of  $D$  idoneal, i.e., if and only if  $D$  satisfies the following condition:*

$$(44) \quad g_{D/p^2} = h_{D/p^2}, \text{ for all primes } p \text{ such that } p^2 \mid D \text{ and } D/p^2 \equiv 0, 1 \pmod{4}.$$

**Proof.** (a) If  $D$  is a fundamental discriminant, then every  $\chi \in \text{Cl}(D)^*$  is primitive, and hence by Corollary 33 (and Proposition 9)  $\Theta_D$  has a basis consisting of  $\mathbb{T}_{|D|}$ -eigenforms and hence is a  $\mathbb{T}_{|D|}$ -module.

Next, suppose that  $\Theta_D$  is a  $\mathbb{T}_{|D|}$ -module. Then we have:

$$(45) \quad \vartheta_\chi \text{ is a } \mathbb{T}_{|D|}\text{-eigenfunction, } \forall \chi \in \text{Cl}(D)^*.$$

To see this, note first  $f := \vartheta_\chi$  is a  $\mathbb{T}(D)$ -eigenfunction by Theorem 12. Consider  $n \geq 1$  and put  $f_n := f|T_n$ . By hypothesis,  $f_n \in \Theta_D$ . Since  $T_n$  commutes with the operators in  $\mathbb{T}(D)$ ,  $f_n$  is the same  $\mathbb{T}(D)$ -eigenspace as  $f$ . By multiplicity 1 (cf. Theorem 1), it follows that  $f_n = c_n f$ , for some  $c_n \in \mathbb{C}$ . Thus  $f = \vartheta_\chi$  is a  $\mathbb{T}_{|D|}$ -eigenfunction, as claimed.

Thus, if  $\Theta_D$  is a  $\mathbb{T}_{|D|}$ -eigenspace, then it follows from (45) that  $\Theta_D^E = \sum_{\chi^2=1} \mathbb{C}\vartheta_\chi$  is also a  $\mathbb{T}_{|D|}$ -eigenspace.

Now assume that  $\Theta_D^E$  is a  $\mathbb{T}_{|D|}$ -eigenspace. Then the above proof of (45) yields that for each quadratic  $\chi$ , the series  $\vartheta_\chi$  is a  $\mathbb{T}_{|D|}$ -eigenfunction, and hence by Corollary 33 we obtain that all quadratic characters are primitive. In particular, the trivial character  $\chi = 1$  is primitive, which can happen only when  $D$  is fundamental; cf. Example 35 below.

(b) By the same argument as in part (a) we see that  $\Theta_D^S$  is a  $\mathbb{T}_{|D|}$ -module if and only if every non-quadratic  $\chi \in \text{Cl}(D)^*$  is primitive. But this means that for each prime  $p$  as in (44) we have that  $\text{Cl}(D/p^2)^*$  has only quadratic characters, and so  $\text{Cl}(D/p^2)$  is an elementary abelian 2-group, that is to say,  $g_{D/p^2} = h_{D/p^2}$ . Thus (44) holds (and conversely).  $\square$

We now illustrate the main results of this paper by working out some special cases. The first example examines  $\vartheta_\chi$  when  $\chi = 1 \in \text{Cl}(D)^*$  is the trivial character.

**Example 35.** (a) If  $D = d_K < 0$  is a *fundamental* discriminant, then the theta function of associated to the trivial character  $\chi_{d_K,0} = 1 \in \text{Cl}(d_K)^*$  is

$$\vartheta_K(z) := \vartheta_{\chi_{d_K,0}}(z) = f_1(z; 1, \psi_{d_K}) \text{ with } L(s, \vartheta_K) = \zeta(s)L(s, \psi_{d_K}) = \zeta_K(s);$$

i.e., its associated  $L$ -function is just the Dedekind  $\zeta$ -function of  $K = \mathbb{Q}(\sqrt{D})$ . In particular, the  $T_p$ -eigenvalue of  $\vartheta_K$  is  $a_p(\vartheta_K) = 1 + \left(\frac{d_K}{p}\right)$ , for all primes  $p$ .

(b) If  $D = d_K f_D^2$  is any negative discriminant, then the trivial character  $\chi_{D,0} = 1$  on  $\text{Cl}(D)$  has conductor  $f_{\chi_{D,0}} = 1$ , and its associated primitive character is

$$(\chi_{D,0})_{pr} = \chi_{d_K,0} = 1 \in \text{Cl}(d_K)^*$$

Thus, by (2), (3) and part (a) we have that

$$\vartheta_{\chi_{D,0}}(z) = \sum_{n|f_D^2} c_n(\chi_{D,0})\vartheta_K(nz) \quad \text{and} \quad L(s, \vartheta_{\chi_{D,0}}) = C(s, \chi_{D,0})\zeta_K(s),$$

where the coefficients  $c_n(\chi_{D,0})$  of the finite Dirichlet series  $C(s, \chi_{D,0})$  are given explicitly by Theorem 28.

We thus obtain an expression for  $a_n(\chi_{0,D}) = \#Id_n(\mathfrak{O}_D)$  for all  $n \geq 1$ . In particular, the formula of Remark 32(b) complements the expression obtained in the Appendix (cf. Proposition 48). Thus, if  $p^{e_p} || f_D$ , then we have

$$\begin{aligned} \#I_{p^k}(\mathcal{O}_D) &= p^{e_p} \left(1 - \frac{1}{p}\psi_{d_K}(p)\right) \#I_{p^{k-2e_p}}(\mathfrak{O}_K) \\ &= p^{e_p} \left(1 - \frac{1}{p}\psi_{d_K}(p)\right) \sum_{i=0}^{k-2e_p} \psi_{d_K}(p)^i, \end{aligned}$$

if  $k \geq 2e_p$ . (For  $k < 2e_p$  we have that  $\#I_{p^k}(\mathcal{O}_D) = 0$  and that  $\#I_{p^k}(\mathcal{O}_D) = p^{k/2}$ , for  $k$  odd and even, respectively, as can also be deduced from the Appendix §6.2.)

We now illustrate what happens when the class number  $h_D$  is small.

**Example 36.** (a)  $h_D = 1$ . In this case  $\text{Cl}(D) = \{1_D\}$  and  $\text{Cl}(D)^* = \{\chi_{D,0}\}$ . Thus, by (6) we have that  $\vartheta_{1_D} = w_D \vartheta_{\chi_{D,0}}$ . If  $D = d_K$  is a fundamental discriminant (i.e., if  $-D = 3, 4, 7, 8, 11, 19, 43, 67, 163$ ), then we have by Example 35(a) that

$$\vartheta_{\chi_{D,0}} = \vartheta_K \quad \text{and} \quad L(s, \chi_{D,0}) = \zeta_K(s).$$

In the remaining cases (i.e.,  $-D = 12, 16, 27, 28$ ) this formula is no longer true. Indeed, since  $f = f_D \in \{2, 3\}$  is here a prime, we obtain from Theorem 28 that

$$C(s, \chi_{D,0}) = 1 - a_D f^{-s} + f^{1-2s}, \quad \text{where } a_D := a_f(\chi_{d_K,0}) = 1 + \left(\frac{d_K}{f}\right),$$

(i.e.,  $a_D = 0, 1, 1, 2$ , respectively), and so

$$\begin{cases} \vartheta_{\chi_{D,0}}(z) = \vartheta_K(z) - a_D \vartheta_K(fz) + f \vartheta_K(f^2z), \\ L(s, \vartheta_{\chi_{D,0}}) = (1 - a_D f^{-s} + f^{1-2s})\zeta_K(s). \end{cases}$$

(b)  $h_D = 2$ . Here  $\text{Cl}(D) = \{1_D, cl(f)\}$  and  $\text{Cl}(D)^* = \{\chi_{D,0}, \chi\}$ . Since  $D < -4$ , we have that  $w_D = 2$  and hence it follows from (6) that

$$\vartheta_{\chi_{D,0}} = \frac{1}{2}(\vartheta_{1_D} + \vartheta_f) \quad \text{and} \quad \vartheta_\chi = \frac{1}{2}(\vartheta_{1_D} - \vartheta_f).$$

Since  $g_D = h_D = 2$ , we know by genus theory that  $D = D_1 D_2 c^2$ , where  $D_1 < 0$  and  $D_2 > 0$  are fundamental discriminants (with  $(D_1, D_2) = 1$ ) and  $c \geq 1$ , and that the associated Hecke character  $\tilde{\chi}$  satisfies  $\tilde{\chi}(\mathfrak{p}) = \psi_{D_i}(N_K(\mathfrak{p}))$ , for  $i = 1, 2$  and  $\mathfrak{p} \nmid D$ . Thus,  $f_\chi = D_1 D_2$  and  $f_\chi = c$ , and

$$\vartheta_{\chi_{pr}}(z) = f_1(z; \psi_{D_1}, \psi_{D_2}) \quad \text{and} \quad L(s, \vartheta_{\chi_{pr}}) = L(s, \psi_{D_1})L(s, \psi_{D_2}).$$

In particular, the  $T_p$ -eigenvalue of  $\vartheta_\chi$  is  $a_p(\chi) = \left(\frac{D_1}{p}\right) + \left(\frac{D_2}{p}\right)$ , for  $p \nmid D$ . Note that if  $\chi$  is primitive, i.e., if  $c = 1$ , then  $\vartheta_\chi = \vartheta_{\chi_{pr}}$ . If  $c > 1$ , then a check of all 29  $D$ 's with  $h_D = 2$  shows that necessarily  $D = -60$ , and then  $\vartheta_\chi$  is worked out in Example 37(b) below.

(c)  $h_D = 3$ . Here  $\text{Cl}(D) = \{1_D, cl(f), cl(f)^{-1}\}$  and  $\text{Cl}(D)^* = \{\chi_{D,0}, \chi, \chi^{-1}\}$ . Since  $\text{Re}(\chi(f)) = \text{Re}\left(\frac{-1 \pm \sqrt{-3}}{2}\right) = -\frac{1}{2}$ , we have by (7) that

$$\vartheta_{\chi_{D,0}} = \frac{1}{2}(\vartheta_{1_D} + 2\vartheta_f) \quad \text{and} \quad \vartheta_\chi = \vartheta_{\chi^{-1}} = \frac{1}{2}(\vartheta_{1_D} - \vartheta_f).$$

By Remark 17(a) we know that  $\Theta_D^E = \mathbb{C}\vartheta_{\chi_{D,0}}$  and that  $\Theta_D^S = \mathbb{C}\vartheta_\chi$ ; in particular,  $\vartheta_\chi$  is a cusp form and a  $\mathbb{T}(D)$ -eigenfunction; cf. Theorem 1. Moreover, Corollary 29 shows that  $\vartheta_\chi$  is a newform of level  $|D|$  if and only if  $\chi$  is primitive, i.e., if and only there is no  $c > 1$  such that  $h_{D/c^2} = 3$ . This condition holds for 23 of the cases with  $h_D = 3$  (i.e., for  $-D = 23, 31, 44, 59, 76, 83, 92, 107, 108, 124, 139, 172, 211, 243, 268, 283, 307, 331, 379, 499, 547, 643, 652, 883, 907$ ) but fails for the two cases

$$-D = 92 = 2^2 \cdot 23 \quad \text{and} \quad -D = 124 = 2^2 \cdot 31.$$

We complement the above discussion by working out two numerical examples.

**Example 37.** (a)  $D = -23$ .

Here  $\text{Cl}(D) = \{[1, 1, 6], [2, 1, 3], [2, -1, 3]\}$ , where  $[a, b, c]$  denotes the equivalence class of the form  $ax^2 + bxy + cy^2$ , and so  $h_D = 3$ . Since  $D$  is a fundamental discriminant, we have by Examples 35(a) and 36(c) that  $\vartheta_{\chi_{D,0}} = \vartheta_K = f_1(\cdot; 1, \psi_{-23})$  and  $\vartheta_\chi$  are primitive forms of level 23, and that  $\vartheta_\chi$  is a cusp form. (In fact,  $N = |D| = 23$  is the smallest level for which  $\Theta_D^S \neq \{0\}$ .) Moreover,

$$\vartheta_{\chi_{D,0}} = \vartheta_K = \frac{1}{2}(\vartheta_{[1,1,6]} + 2\vartheta_{[2,1,3]}) \quad \text{and} \quad \vartheta_\chi = \frac{1}{2}(\vartheta_{[1,1,6]} - \vartheta_{[2,1,3]}),$$

and from this (or from (8)) we obtain that

$$\vartheta_{[1,1,6]} = \frac{2}{3}(\vartheta_K + 2\vartheta_\chi) \quad \text{and} \quad \vartheta_{[2,1,3]} = \frac{2}{3}(\vartheta_K - \vartheta_\chi).$$

(b)  $D = -60 = -15 \cdot 2^2$ .

Here  $\text{Cl}(D) = \{[1, 0, 15], [3, 0, 5]\}$ , so  $h_D = 2$  and  $\text{Cl}(D)^* = \{\chi_{D,0}, \chi\}$ . Since  $h_{d_K} = h_{-15} = 2$ , we see that  $f_{\chi_{D,0}} = f_\chi = 1$  and so both  $\chi_{D,0}$  and  $\chi$  are imprimitive with  $\bar{f}_{\chi_{D,0}} = \bar{f}_\chi = 2$ . The theta-functions of the associated primitive characters are  $\vartheta_K = f_1(\cdot; 1, \psi_{-15})$  and  $\vartheta_{\chi_{pr}} = f_1(\cdot; \psi_{-3}, \psi_5)$ , respectively. Since

$$a_2(\vartheta_K) = 1 + \left(\frac{-15}{2}\right) = 2 \quad \text{and} \quad a_2(\chi_{pr}) = \left(\frac{-3}{2}\right) + \left(\frac{5}{2}\right) = -2,$$

it follows from Theorem 28 that

$$C(s, \chi_{D,0}) = 1 - 2 \cdot 2^{-s} + 2 \cdot 2^{-2s}$$

and that

$$C(s, \chi) = 1 + 2 \cdot 2^{-s} + 2 \cdot 2^{-2s}.$$

Thus,

$$\vartheta_{\chi_{D,0}}(z) = \vartheta_K(z) - 2\vartheta_K(2z) + 2\vartheta_K(4z)$$

and

$$\vartheta_\chi(z) = \vartheta_{\chi_{pr}}(z) + 2\vartheta_{\chi_{pr}}(2z) + 2\vartheta_{\chi_{pr}}(4z),$$

and the associated  $L$ -functions are

$$L(s, \chi_{D,0}) = (1 - 2^{1-s} + 2^{1-2s})\zeta_K(s) \quad \text{and} \quad L(s, \chi) = (1 + 2^{1-s} + 2^{1-2s})L(s, \chi_{pr}),$$

where  $L(s, \chi_{pr}) = L(s, \psi_{-3})L(s, \psi_5)$ .

## 6. Appendix: Ideals of quadratic orders

The purpose of this appendix is to collect some well-known results about ideals in an order  $\mathfrak{D}_D$  of an imaginary quadratic field  $K$ , and to extend these to obtain the results which were used in this article.

### 6.1. Lattices in an imaginary quadratic field

Let  $K = \mathbb{Q}(\sqrt{d_K})$  be an imaginary quadratic field of discriminant  $d_K < 0$ , and let  $\mathfrak{O}_K = \mathbb{Z} + \omega_K\mathbb{Z}$  denote its ring of integers, where  $\omega_K = \frac{1}{2}(d_K + \sqrt{d_K})$ . For any  $f \geq 1$ , put  $D = f^2d_K$  and

$$\mathfrak{D}_D = \mathbb{Z} + f\omega_K\mathbb{Z} = \mathbb{Z} + \omega_D\mathbb{Z}, \quad \text{where } \omega_D = \frac{1}{2}(D + \sqrt{D}).$$

It is immediate that  $\mathfrak{D}_D$  is a subring of  $\mathfrak{O}_K$  of index  $f_{\mathfrak{D}_D} := [\mathfrak{O}_K : \mathfrak{D}_D] = f$ . The ring  $\mathfrak{D}_D$  is called the *order of discriminant  $D$*  (or *of conductor  $f$* ), and it is well-known that every subring  $R$  of  $\mathfrak{O}_K$  with quotient field  $K$  is of this form, i.e.,  $R = \mathfrak{D}_D$  with  $D = f_R^2d_K$ , where  $f_R = [\mathfrak{O}_K : R]$ . (For such and other basic facts about orders see [34], §90-113, [3], §II.7, [21], §8.1, [6], §II.7.)

Let  $\text{Lat}(K)$  denote the set of all *lattices* in  $K$ , i.e., the set of all finitely generated  $\mathbb{Z}$ -submodules of  $K$  which contain some  $\mathbb{Q}$ -basis of  $K$ .

If  $L \in \text{Lat}(K)$  is a lattice, then its *associated order* is  $\mathfrak{D}(L) = \{\lambda \in K : \lambda L \subset L\}$ . It is easy to see that  $\mathfrak{D}(L)$  is an *order* of  $K$ , i.e.,  $\mathfrak{D}(L) = \mathfrak{D}_D$ , for some  $D = f^2d_K$ .

The *norm* of the lattice  $L$  is by defined by  $N(L) = |\det(T)|$ , where  $T \in \text{GL}_2(\mathbb{Q})$  is such that  $T(\mathfrak{D}(L)) = L$ ; cf. [3], §II.6. Note that if  $L \subset \mathfrak{O}_K$ , then

$$(46) \quad [\mathfrak{O}_K : L] = [\mathfrak{O}_K : \mathfrak{D}(L)]N(L).$$

If  $L_1, L_2 \in \text{Lat}(K)$  are two lattices, then the product (module)  $L_1L_2$  is again a lattice. By [3], §II.7, Ex. 6 and 10, its order and norm are given by the formulae

$$(47) \quad \mathfrak{D}(L_1L_2) = \mathfrak{D}(L_1)\mathfrak{D}(L_2) \quad \text{and} \quad N(L_1L_2) = N(L_1)N(L_2).$$

In particular, we see that the set  $I(\mathfrak{D}_D) = \{L \in \text{Lat}(K) : \mathfrak{D}(L) = \mathfrak{D}_D\}$  is closed under multiplication. In fact,  $I(\mathfrak{D}_D)$  is a group with unit element  $\mathfrak{D}_D$ : it can be identified with the group of invertible  $\mathfrak{D}_D$ -submodules of  $K$ ; cf. Lang [21], p. 91. Thus, if  $P(\mathfrak{D}_D) = \{\lambda\mathfrak{D}_D : \lambda \in K^\times\}$  denotes the subgroup of principal  $\mathfrak{D}_D$ -modules, then the quotient

$$\text{Cl}(\mathfrak{D}_D) = \text{Pic}(\mathfrak{D}_D) = I(\mathfrak{D}_D)/P(\mathfrak{D}_D)$$

is called the *class group* (or *Picard group*) of the order  $\mathfrak{D}_D$ , and its cardinality (or group order)  $h_D := |\text{Cl}(\mathfrak{D}_D)|$  is called the *class number* of  $\mathfrak{D}_D$ .

If  $\mathfrak{O}_{D_i}$  are two orders of  $K$  with discriminants  $D_i = f_{D_i}^2 d_K$ , then

$$(48) \quad \mathfrak{O}_{D_1} \supset \mathfrak{O}_{D_2} \quad \text{if and only if} \quad f_{D_1} \mid f_{D_2},$$

and hence

$$(49) \quad \mathfrak{O}_{D_1} \cap \mathfrak{O}_{D_2} = \mathfrak{O}_{\text{lcm}(f_{D_1}, f_{D_2})^2 d_K} \quad \text{and} \quad \mathfrak{O}_{D_1} \mathfrak{O}_{D_2} = \mathfrak{O}_{\text{gcd}(f_{D_1}, f_{D_2})^2 d_K}.$$

**Proposition 38.** *If  $R$  and  $R'$  are two orders of  $K$  with  $R \subset R'$ , then the rule  $L \mapsto LR'$  defines a surjective homomorphism  $\pi_{R,R'} : I(R) \rightarrow I(R')$  with kernel*

$$(50) \quad \text{Ker}(\pi_{R,R'}) = \{ L \in I(R) : L \subset R' \text{ and } [R' : L] = [R' : R] \}.$$

**Proof.** If  $L \in I(R)$ , then by (47) we have

$$\mathfrak{O}(LR') = \mathfrak{O}(L)\mathfrak{O}(R') = RR' = R',$$

so  $LR' \in I(R')$ . Thus, the rule  $L \mapsto LR'$  defines a map  $\pi = \pi_{R,R'} : I(R) \rightarrow I(R')$ . Moreover,  $\pi$  is a homomorphism because

$$\pi(L_1)\pi(L_2) = L_1R'L_2R' = L_1L_2R'R' = L_1L_2R' = \pi(L_1L_2).$$

To prove (50), let  $L \in \text{Ker}(\pi)$ . Then  $LR' = R'$ . Since  $R'$  is an order, we have that

$$N(R') = [\mathfrak{O}(R') : R'] = [R' : R'] = 1,$$

and so by (47) we obtain that

$$N(L) = N(L)N(R') = N(LR') = N(R') = 1.$$

Thus, since  $\mathfrak{O}(L) = R$ , we see that

$$[R' : R] = [R' : R]N(L) = [R' : R][R : L] = [R' : L].$$

Moreover, since  $L = L \cdot 1 \subset LR' = R'$  it follows that  $L \subset R'$ , and hence

$$L \in \mathcal{K} := \{ L \in I(R) : L \subset R' \text{ and } [R' : L] = [R' : R] \}.$$

Conversely, if  $L \in \mathcal{K}$ , then

$$N(L) = [R : L] = [R : L]/[R' : R] = 1,$$

and hence

$$N(LR') = N(L)N(R') = 1 \cdot 1 = 1.$$

Now since

$$\mathfrak{O}(LR') = \mathfrak{O}(L)\mathfrak{O}(R') = RR' = R',$$

we have that  $1 = N(LR') = [R' : LR']$ . But since  $L \subset R'$ , we have  $LR' \subset R'$  and so this forces  $LR' = R'$ . Thus  $L \in \text{Ker}(\pi)$ , which proves (50).

It remains to show that  $\pi$  is surjective. Let  $L' \in I(R')$ . Then by [21], Theorem 5 (p. 93), there exists  $\lambda \in K^\times$  such that  $\lambda L' + fR' = R'$ , where  $f = [\mathfrak{O}_K : R]$ . We then have that  $L := \lambda L' \cap R \in I(R)$  and that  $LR' = \lambda L'$  by applying [21], Theorem 4 (p. 92), to  $R'$  and  $R$ . (See also Proposition 48 below.) Thus  $\pi(\lambda^{-1}L) = (\lambda^{-1}L)R' = L'$ , and so  $\pi$  is surjective.  $\square$

**Remark 39.** Since  $\pi_{R,R'}(\lambda R) = \lambda R'$ , for all  $\lambda \in K^\times$ , it is clear that  $\pi_{R,R'}$  induces a surjective homomorphism

$$(51) \quad \bar{\pi}_{R,R'} : \text{Cl}(R) = I(R)/P(R) \rightarrow \text{Cl}(R') = I(R')/P(R').$$

Moreover, it is immediate that

$$(52) \quad \text{Ker}(\bar{\pi}_{R,R'}) = \text{Ker}(\pi_{R,R'})P(R)/P(R)$$

because if  $LP(R) \in \text{Ker}(\bar{\pi}_{R,R'})$ , then  $L = \lambda R'$ , for some  $\lambda \in K^\times$ , and then

$$LP(R) = (\lambda^{-1}L)(\lambda R)P(R) \in \text{Ker}(\pi_{R,R'})P(R)/P(R),$$

so (52) follows.

The following result, which gives the order and a presentation of  $\text{Ker}(\pi_{R,R'})$ , can be viewed as a refinement of the well-known results which give a presentation and the order of  $\text{Ker}(\bar{\pi}_{R,R'})$ ; cf. Lang [21], p. 95 and Cox [6], p. 147. Note that while the formula for  $|\text{Ker}(\bar{\pi}_{R,R'})|$  involves the unit index  $[R^\times : (R')^\times]$ , this is not the case for that of  $|\text{Ker}(\pi_{R,R'})|$ ; cf. (54).

**Corollary 40.** *In the above situation put  $\bar{f} = [R' : R]$ . Then the map defined by the rule  $\lambda \mapsto L_\lambda := \mathbb{Z}\lambda + \bar{f}R'$  induces the exact sequence*

$$(53) \quad 0 \rightarrow (\mathbb{Z}/\bar{f}\mathbb{Z})^\times \rightarrow (R'/\bar{f}R')^\times \xrightarrow{L_{R',\bar{f}}} \text{Ker}(\pi_{R,R'}) \rightarrow 0.$$

Thus, if  $D'$  denotes the discriminant of  $R' = \mathfrak{O}_{D'}$  and if  $\psi_{R'}(p) = \left(\frac{D'}{p}\right)$  denotes the associated Legendre-Kronecker symbol, then

$$(54) \quad |\text{Ker}(\pi_{R,R'})| = \bar{f} \prod_{p|\bar{f}} \left(1 - \frac{1}{p} \psi_{R'}(p)\right),$$

**Proof.** First note that if  $\lambda \in R'$ , then  $L_\lambda/\bar{f}R' \leq R'/\bar{f}R'$  is the cyclic subgroup generated by  $\bar{\lambda} = \lambda + \bar{f}R'$ , and hence  $L_\lambda = L_{\bar{\lambda}}$  depends only on  $\bar{\lambda}$ . Moreover, since  $\lambda\bar{f}R' = \bar{f}R'$ , if  $\bar{\lambda} \in (R'/\bar{f}R')^\times$ , it is clear that  $L_\lambda L_{\lambda'} = L_{\lambda\lambda'}$ , and so the rule  $\lambda \mapsto L_\lambda$  defines a homomorphism  $L_{R',\bar{f}} : (R'/\bar{f}R')^\times \rightarrow \text{Lat}(K)$ . Next we show that  $\text{Im}(L_{R',\bar{f}}) \subset \text{Ker}(\pi_{R,R'})$ . For this, note first that  $L_\lambda$  is clearly an  $R$ -module because  $R = \mathbb{Z} + \bar{f}R'$ . Moreover, since  $\lambda\lambda' \equiv 1 \pmod{\bar{f}R'}$ , for some  $\lambda' \in R'$ , we see that  $L_\lambda L_{\lambda'} = L_1 = R$ , and so  $L_\lambda \in I(R)$ . In addition,  $L_\lambda R' = R'$  because  $L_\lambda R'$  is an  $R'$ -ideal which contains 1, and so  $L_\lambda \in \text{Ker}(\pi_{R,R'})$ .

To show that  $L_{R',\bar{f}}$  is surjective, let  $L \in \text{Ker}(\pi_{R,R'})$ . Then by (50) we know that  $L \subset R'$  and that  $[R' : L] = \bar{f}$ , so  $L \supset \bar{f}R'$  and

$$[L : \bar{f}R'] = \frac{[R' : \bar{f}R']}{[R' : L]} = \frac{\bar{f}^2}{\bar{f}} = \bar{f}.$$

Since  $LR' = R'$ , we see that  $L \not\subset pR'$ , for any prime  $p|\bar{f}$ , and hence  $L/\bar{f}R'$  is cyclic, i.e.,  $L = \mathbb{Z}\lambda + \bar{f}R'$  for some  $\lambda \in R'$ . Since  $|L/\bar{f}R'| = \bar{f}$ , this means that  $\lambda$  has order  $\bar{f}$  in  $R'/\bar{f}R' \simeq \mathbb{Z}/\bar{f}\mathbb{Z} \times \mathbb{Z}/\bar{f}\mathbb{Z}$ . Then  $\lambda = a + b\bar{f}\omega_{D'}$  with  $a, b \in \mathbb{Z}$  and  $(a, b, \bar{f}) = 1$ , where  $D'$  is the discriminant of  $R' = \mathfrak{O}_{D'}$ , and so it follows easily that  $\lambda \in (R'/\bar{f}R')^\times$ . Thus  $L = L_\lambda = L_{R',\bar{f}}(\lambda)$ , and hence  $L_{R',\bar{f}}$  is surjective.

Next we observe that  $\bar{\lambda} \in \text{Ker}(L_{R', \bar{f}}) \Leftrightarrow L_{\bar{\lambda}} = R = \mathbb{Z} + \bar{f}R' \Leftrightarrow \bar{\lambda} = n + \bar{f}R'$ , for some  $n \in \mathbb{Z}$  with  $(n, \bar{f}) = 1$ . Thus, since the map  $\mathbb{Z}/\bar{f}\mathbb{Z} \rightarrow R'/\bar{f}R'$  is injective, it follows that the sequence (53) is exact.

It remains to verify (54). Applying (53) with  $R' = \mathfrak{O}_K$  (and hence  $\bar{f} = f := f_R$ ) yields that

$$|\text{Ker}(\pi_{R, \mathfrak{O}_K})| = \frac{\phi(f\mathfrak{O}_K)}{\phi(f)} = f \prod_{p|f} \left(1 - \frac{1}{p} \psi_{\mathfrak{O}_K}(p)\right),$$

where the second formula is deduced as in Lang [21], p. 95. (Here, as in Lang [21],  $\phi(f\mathfrak{O}_K) = |(\mathfrak{O}_K/f\mathfrak{O}_K)^\times|$ .) From this (and the surjectivity of  $\pi_{R, R'}$ ) it follows that for any  $R' \supset R$  we have

$$\begin{aligned} |\text{Ker}(\pi_{R, R'})| &= \frac{|\text{Ker}(\pi_{R, \mathfrak{O}_K})|}{|\text{Ker}(\pi_{R', \mathfrak{O}_K})|} \\ &= \bar{f} \prod_{p|f_R, p \nmid f_{R'}} \left(1 - \frac{1}{p} \psi_{d_K}(p)\right) \\ &= \bar{f} \prod_{p|\bar{f}} \left(1 - \frac{1}{p} \psi_{R'}(p)\right). \quad \square \end{aligned}$$

**Corollary 41.** *If  $R \subset R_1 \cap R_2$  and  $R_3 = R_1 R_2$ , then*

$$\text{Ker}(\pi_{R, R_3}) = \text{Ker}(\pi_{R, R_1}) \cdot \text{Ker}(\pi_{R, R_2}) \quad \text{and} \quad \text{Ker}(\bar{\pi}_{R, R_3}) = \text{Ker}(\bar{\pi}_{R, R_1}) \cdot \text{Ker}(\bar{\pi}_{R, R_2}).$$

**Proof.** By (52) it is enough to prove the first assertion. Moreover, since

$$\text{Ker}(\pi_{R, R_i}) = \pi_{R, R_1 \cap R_2}^{-1}(\text{Ker}(\pi_{R_1 \cap R_2, R_i})), \quad \text{for } i = 1, 2, 3,$$

it suffices to verify the formula for  $R = R_1 \cap R_2$ . For this, write

$$f = [\mathfrak{O}_K : R], \quad f_i = [\mathfrak{O}_K : R_i] \quad \text{and} \quad \bar{f}_i = \frac{f}{f_i} = [R_i : R], \quad \text{for } i = 1, 2, 3.$$

Then by (49) we have that  $f_3 = (f_1, f_2)$ . Since  $R = R_1 \cap R_2$ , we have that  $f = \frac{f_1 f_2}{f_3}$  by (49), so  $\bar{f}_3 = \bar{f}_1 \bar{f}_2$  and  $(\bar{f}_1, \bar{f}_2) = 1$ . It thus follows from (54) that

$$|\text{Ker}(\pi_{R, R_3})| = |\text{Ker}(\pi_{R, R_1})| \cdot |\text{Ker}(\pi_{R, R_2})|.$$

Thus, the assertion follows once we have shown that  $\text{Ker}(\pi_{R, R_1}) \cap \text{Ker}(\pi_{R, R_2}) = \{R\}$ . Now if  $L \in \text{Ker}(\pi_{R, R_1}) \cap \text{Ker}(\pi_{R, R_2})$ , then  $L \subset R_1 \cap R_2 = R$ . But since

$$[R : L] = N(L) = 1$$

(the latter because  $LR_i = R_i$ ), it follows that  $L = R$ , and hence

$$\text{Ker}(\pi_{R, R_1}) \cap \text{Ker}(\pi_{R, R_2}) = \{R\},$$

as claimed. □

## 6.2. The ideal theory of $\mathfrak{D}_D$

We now study the ideals of the order  $R = \mathfrak{D}_D \subset K$  more closely. For this, we shall use the following general result which is applicable here because  $\mathfrak{D}_D$  is a noetherian domain of (Krull) dimension 1; cf. Neukirch [23], p. 77.

**Proposition 42.** *Let  $A$  be a noetherian domain of dimension 1, and let  $\mathfrak{a} \subset A$  be a non-zero ideal of  $A$ . If  $\mathfrak{m} \in \max(A)$  is any maximal ideal of  $A$  with  $\mathfrak{a} \subset \mathfrak{m}$ , then  $\mathfrak{a}(\mathfrak{m}) := \mathfrak{a}A_{\mathfrak{m}} \cap A$  is an  $\mathfrak{m}$ -primary ideal of  $A$ . In particular, if  $|A/\mathfrak{m}| < \infty$ , then  $|A/\mathfrak{a}(\mathfrak{m})| = |A/\mathfrak{m}|^s$ , for some integer  $s = s_{\mathfrak{m}}(\mathfrak{a}) \geq 1$ . Moreover, we have*

$$(55) \quad \mathfrak{a} = \bigcap_{\mathfrak{m}} \mathfrak{a}(\mathfrak{m}) = \prod_{\mathfrak{m}} \mathfrak{a}(\mathfrak{m}).$$

Thus  $\mathfrak{a}$  is an invertible ideal if and only if  $\mathfrak{a}(\mathfrak{m})$  is invertible for all maximal ideals  $\mathfrak{m}$ .

**Proof.** Since  $A_{\mathfrak{m}}$  is a local 1-dimensional noetherian domain, every proper ideal of  $A_{\mathfrak{m}}$  is primary, and hence  $\mathfrak{a}(\mathfrak{m})$  is primary ideal of  $A$  with radical  $\mathfrak{m}$ . This proves the first assertion. From this the second assertion is an immediate consequence because by [4], p. 265, we know that the  $A$ -module  $M = A/\mathfrak{a}(\mathfrak{m})$  has a composition series  $\{M_i\}$  with  $M_i/M_{i+1} \simeq A/\mathfrak{m}$ , and so the assertion follows (with  $s = \text{length}(M)$ ).

The first identity of (55) is true in any domain; cf. [4], p. 89. Next, if  $\mathfrak{m}$  and  $\mathfrak{m}'$  are two distinct maximal ideals of  $A$ , then  $\mathfrak{a}(\mathfrak{m}) \not\subset \mathfrak{m}'$  and hence  $\mathfrak{a}(\mathfrak{m})A_{\mathfrak{m}'} = A_{\mathfrak{m}'}$ . Thus, if we put  $\mathfrak{a}' = \prod \mathfrak{a}(\mathfrak{m})$ , then  $\mathfrak{a}'A_{\mathfrak{m}} = \mathfrak{a}(\mathfrak{m})A_{\mathfrak{m}} = \mathfrak{a}A_{\mathfrak{m}}$ , for every  $\mathfrak{m} \in \max(A)$ , and so  $\mathfrak{a}' = \mathfrak{a}$ , which proves the second equality of (55).

Finally, if  $\mathfrak{a}(\mathfrak{m})$  is invertible for all  $\mathfrak{m} \in \max(A)$ , then  $\mathfrak{a}$  is invertible by (55). Conversely, suppose that  $\mathfrak{a}$  is invertible and fix  $\mathfrak{m} \in \max(A)$  with  $\mathfrak{m} \supset \mathfrak{a}$ . Then  $\mathfrak{a}(\mathfrak{m})A_{\mathfrak{m}} = \mathfrak{a}A_{\mathfrak{m}}$  is principal, and for every  $\mathfrak{m}' \neq \mathfrak{m}$  we have  $\mathfrak{a}(\mathfrak{m})A_{\mathfrak{m}'} = A_{\mathfrak{m}'}$ . Thus  $\mathfrak{a}(\mathfrak{m})$  is locally principal and hence invertible; cf. [4], p. 117.  $\square$

**Notation.** Let  $Id(R) = \{L \in I(R) : L \subset R\}$  denote the set of invertible  $R$ -ideals. Moreover, for any integer  $n \geq 1$ , let  $Id_n(R) = \{\mathfrak{a} \in Id(R) : N(\mathfrak{a}) = n\}$  denote the set of invertible  $R$ -ideals of norm  $n$ , and let  $Id(R, n) = \{\mathfrak{a} \in Id(R) : (N(\mathfrak{a}), n) = 1\}$  denote the set of invertible  $R$ -ideals whose norm is relatively prime to  $n$ .

**Corollary 43.** *Let  $\mathfrak{a} \in Id(R)$  be an invertible  $R$ -ideal, and  $\mathfrak{m} \in \max(R)$  be a maximal ideal, where  $R = \mathfrak{D}_D$ . Then  $\mathfrak{a}(\mathfrak{m}) \in Id(R)$  is also invertible and has norm  $N(\mathfrak{a}(\mathfrak{m})) = p^s$ , for some  $s$ , where  $p \mid \mathfrak{m}$ , i.e.,  $p$  is the unique prime number with  $p \in \mathfrak{m}$ . Furthermore,*

$$\mathfrak{a} = \prod_{\mathfrak{m}} \mathfrak{a}(\mathfrak{m}) \quad \text{and hence} \quad N(\mathfrak{a}) = \prod_{\mathfrak{m}} N(\mathfrak{a}(\mathfrak{m})).$$

In particular, if  $\mathfrak{m} \in \text{supp}(\mathfrak{a}) := \{\mathfrak{m} \in \max(R) : \mathfrak{a} \subset \mathfrak{m}\}$ , and  $p \mid \mathfrak{m}$ , then  $p \mid N(\mathfrak{a})$ .

**Proof.** If  $\mathfrak{m} \in \max(R)$ , then  $R/\mathfrak{m}$  is a finite field of characteristic  $p$ , where  $p \in \mathfrak{m}$ . Thus, since  $R$  is a 1-dimensional noetherian ring and since  $N(\mathfrak{a}) = [R : \mathfrak{a}]$ , whenever  $\mathfrak{a} \in Id(R)$ , the corollary follows immediately from Proposition 42 (together with (47)).  $\square$

We now want to study the set  $Id_n(R)$  of invertible ideals of norm  $n$  of  $R$  more closely. First we note:

**Proposition 44.** *If  $(m, n) = 1$ , then the map  $(\mathfrak{a}, \mathfrak{b}) \mapsto \mathfrak{a}\mathfrak{b}$  induces a bijection*

$$Id_m(R) \times Id_n(R) \xrightarrow{\sim} Id_{mn}(R).$$

**Proof.** If  $(\mathfrak{a}, \mathfrak{b}) \in Id_m(R) \times Id_n(R)$ , then  $\mathfrak{a}\mathfrak{b} \in Id_{mn}(R)$  by (47). Suppose now that  $\mathfrak{c} \in Id_{mn}(R)$ , and put, for an integer  $k$ ,

$$\text{supp}_k(\mathfrak{c}) := \text{supp}(\mathfrak{c} + kR) = \{ \mathfrak{m} \in \text{supp}(\mathfrak{c}) : p \mid \mathfrak{m} \Rightarrow p \mid k \}$$

and  $\mathfrak{c}_k = \prod_{\mathfrak{m} \in \text{supp}_k(\mathfrak{c})} \mathfrak{a}(\mathfrak{m})$ . Then by Corollary 43 we have

$$\text{supp}(\mathfrak{c}) = \text{supp}(\mathfrak{c})_m \dot{\cup} \text{supp}(\mathfrak{c})_n,$$

and so  $\mathfrak{c} = \mathfrak{c}_m \mathfrak{c}_n$ . Since

$$(N(\mathfrak{c}_m), n) = (N(\mathfrak{c}_n), m) = 1,$$

we see that  $(\mathfrak{c}_m, \mathfrak{c}_n) \in Id_m(R) \times Id_n(R)$ , and so the map is surjective. It is injective because  $\mathfrak{c}$  is uniquely determined by its local components  $\mathfrak{c}(\mathfrak{m})$ .  $\square$

Thus, by the above result, it is enough to study the sets  $Id_n(R)$  for prime powers  $n = p^r$ . If  $p$  is prime to the conductor  $f_R$  of  $R$ , i.e., if  $p \nmid f_R = [\mathfrak{D}_K : R]$ , then  $Id_n(R)$  is essentially the same as  $Id_n(\mathfrak{D}_K)$ , as the next (well-known) result shows.

**Proposition 45.** *The rule  $\mathfrak{a} \mapsto \mathfrak{a} \cap R$  induces an injection*

$$\tilde{\varphi}_R : Id(\mathfrak{D}_K, f_R) \hookrightarrow Id(R)$$

with image  $Id(R, f_R)$ , and we have that

$$(56) \quad \tilde{\varphi}_R(Id_n(\mathfrak{D}_K)) = Id_n(R), \quad \text{for all } n \geq 1, (n, f_R) = 1.$$

Thus

$$(57) \quad \#Id_n(R) = \#Id_n(\mathfrak{D}_K) = \sum_{d|n} \psi_{d_K}(d), \quad \text{if } (n, f_R) = 1.$$

**Proof.** The first assertion follows from [21], Theorem 4 (p. 92). Moreover, since that theorem also asserts that  $\tilde{\varphi}_R(\mathfrak{a})\mathfrak{D}_K = \mathfrak{a}$ , if  $\mathfrak{a} \in Id(\mathfrak{D}_K, f_R)$ , it follows from (47) that  $N(\tilde{\varphi}_R(\mathfrak{a})) = N(\tilde{\varphi}_R(\mathfrak{a}))N(\mathfrak{D}_K) = N(\mathfrak{a})$ . Thus (56) holds, and hence the first equality of (57) follows. The second equality of (57) is well-known; cf. Weber [34], p. 345.  $\square$

We now turn to study the ideals of  $R = \mathfrak{D}_D$  with norm dividing the conductor  $f_R = f_D$ . Here we first prove the following result which is closely related to Theorem 3.1.17 of Schertz [25].

**Proposition 46.** *If  $p \mid f_D$  is a prime divisor of the conductor  $f_D = [\mathfrak{D}_K : \mathfrak{D}_D]$  of  $\mathfrak{D}_D$ , then  $\mathfrak{m}_p := p\mathfrak{D}_D/p^2$  is the unique maximal ideal of  $\mathfrak{D}_D$  containing  $p$ . Moreover,  $|\mathfrak{D}_D/\mathfrak{m}_p| = p$  and  $\mathfrak{m}_p$  is an  $\mathfrak{D}_D$ -ideal which is not invertible, i.e.,  $\mathfrak{m}_p \notin Id(\mathfrak{D}_D)$ . In particular,  $Id_p(\mathfrak{D}_D) = \emptyset$ .*

**Proof.** Since  $\mathfrak{m}_p = p(\mathbb{Z} + (f_D/p)\omega_K\mathbb{Z}) = p\mathbb{Z} + f_D\omega_K\mathbb{Z}$ , it immediately follows that  $\mathfrak{m}_p \subset \mathfrak{D}_D = \mathbb{Z} + f_D\omega_K\mathbb{Z}$  and that  $[\mathfrak{D}_D : \mathfrak{m}_p] = p$ . Clearly,  $\mathfrak{m}_p$  is a principal  $\mathfrak{D}_D/p^2$ -ideal, and hence a fortiori an  $\mathfrak{D}_D$ -ideal. Thus,  $\mathfrak{m}_p$  is a maximal ideal of  $\mathfrak{D}_D$  and  $\mathfrak{D}(\mathfrak{m}_p) = \mathfrak{D}_D/p^2$ , so  $\mathfrak{m}_p$  cannot be invertible as an  $\mathfrak{D}_D$ -ideal.

Now suppose that  $\mathfrak{m} \supset p\mathfrak{D}_D$  is a maximal ideal. Then (by Cohen-Seidenberg)  $\mathfrak{m} = \mathfrak{m}\mathfrak{D}_{D/p^2} \cap \mathfrak{D}_D \supset p\mathfrak{D}_{D/p^2} \cap \mathfrak{D}_D = \mathfrak{m}_p$ , and so  $\mathfrak{m} = \mathfrak{m}_p$ . Thus  $\mathfrak{m}_p$  is the unique maximal ideal containing  $p$ .

Finally, if  $\mathfrak{a} \in Id_p(\mathfrak{D}_D)$ , then  $[\mathfrak{D}_D : \mathfrak{a}] = N(\mathfrak{a}) = p$ , so  $\mathfrak{a}$  is a maximal ideal of  $\mathfrak{D}_D$  with  $p \in \mathfrak{a}$ , and hence  $\mathfrak{a} = \mathfrak{m}_p$ . But  $\mathfrak{m} \notin Id(\mathfrak{D}_D)$ , contradiction. Thus  $Id_p(\mathfrak{D}_D) = \emptyset$ .  $\square$

**Corollary 47.** *Suppose that  $c \mid f_D = [\mathfrak{D}_K : \mathfrak{D}_D]$ . If  $\mathfrak{a} \in Id_n(\mathfrak{D}_D)$  and  $c^2 \mid n$ , then  $\mathfrak{a}\mathfrak{D}_{D/c^2} = c\mathfrak{b}$ , for some  $\mathfrak{b} \in Id_{n/c^2}(\mathfrak{D}_{D/c^2})$ .*

**Proof.** Suppose first that  $c = p$ . Then by Corollary 43 and Proposition 46 we know that  $\mathfrak{a} \subset \mathfrak{m}_p = p\mathfrak{D}_{D/p^2}$ , and so  $\mathfrak{b}' := \mathfrak{a}\mathfrak{D}_{D/p^2} \subset p\mathfrak{D}_{D/p^2}$ . Thus  $\mathfrak{b} := \frac{1}{p}\mathfrak{b}' \subset \mathfrak{D}_{D/p^2}$ , and hence  $\mathfrak{b} \in Id_{n/p^2}(\mathfrak{D}_{D/p^2})$  and  $\mathfrak{a}\mathfrak{D}_{D/p^2} = p\mathfrak{b}$ . Thus the assertion holds for  $c = p$ .

To prove the general case, induct on  $c$ . Since the assertion is trivial for  $c = 1$ , we may assume that there is a prime  $p \mid c$ . Put  $D' = D/c^2$ ,  $\bar{D} = D/p^2$  and  $\bar{n} = n/p^2$ . Then, by what was just shown,  $\mathfrak{a}\mathfrak{D}_{\bar{D}} = p\bar{\mathfrak{a}}$  with  $\bar{\mathfrak{a}} \in Id_{\bar{n}}(\mathfrak{D}_{\bar{D}})$ . Put  $\bar{c} = c/p$ . Then  $\bar{c} \mid \frac{f}{p} = [\mathfrak{D}_K : \mathfrak{D}_{\bar{D}}]$  and  $\bar{c}^2 \mid \bar{n}$ , so by induction  $\bar{\mathfrak{a}}\mathfrak{D}_{D'} = \bar{c}\mathfrak{b}$  with  $\mathfrak{b} \in Id_{\bar{n}/\bar{c}^2}(\mathfrak{D}_{D'})$ . Thus  $\mathfrak{a}\mathfrak{D}_{D'} = (p\bar{\mathfrak{a}})\mathfrak{D}_{D'} = p\bar{c}\mathfrak{b} = c\mathfrak{b}$ , and so the assertion also holds for  $c$ .  $\square$

As we shall now see, the above results show that when  $n \mid f_D^2$ , there is close connection between  $Id_n(\mathfrak{D}_D)$  and  $\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/n}})$ .

**Proposition 48.** *Suppose that  $n \mid f_D^2 = D/d_K$ . If  $n$  is not a square, then we have that  $Id_n(\mathfrak{D}_D) = \emptyset$ , whereas for  $n = c^2$  we have the formula*

$$Id_{c^2}(\mathfrak{D}_D) = c\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D/c^2}}).$$

**Proof.** Suppose first that  $n$  is not a square, i.e., suppose that there is a prime  $p$  such that  $p^{2r+1} \parallel n$ , for some  $r \geq 0$ . If  $\mathfrak{a} \in Id_n(\mathfrak{D}_D)$ , then  $\mathfrak{a}_p := \mathfrak{a}(\mathfrak{m}_p) \in Id_{p^{2r+1}}(\mathfrak{D}_D)$ , and hence, putting  $D' = D/p^{2r}$ , we have by Corollary 47 that  $\mathfrak{a}_p\mathfrak{D}_{D'} = p^r\mathfrak{b}$ , for some  $\mathfrak{b} \in Id_p(\mathfrak{D}_{D'})$ . But since  $p \mid [\mathfrak{D}_K : \mathfrak{D}_{D'}]$ , this contradicts Proposition 46. Thus  $Id_n(\mathfrak{D}_D) = \emptyset$ .

Now suppose that  $n = c^2$ , and put  $D' = D/c^2$ . If  $\mathfrak{a} \in Id_n(\mathfrak{D}_D)$ , then by Corollary 47 we have that  $\mathfrak{a}\mathfrak{D}_{D'} = c\mathfrak{b}$  with  $\mathfrak{b} \in Id_1(\mathfrak{D}_{D'})$ , i.e.,  $\mathfrak{a}\mathfrak{D}_{D'} = c\mathfrak{D}_{D'}$ , and hence  $\mathfrak{a} = cL$  with  $L = \frac{1}{c}\mathfrak{a} \in \text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D'}})$ . Thus  $Id_n(\mathfrak{D}_D) \subset c\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D'}})$ .

Conversely, if  $L \in \text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D'}})$ , then  $L \subset \mathfrak{D}_{D'}$  (cf. (50)), so  $cL \subset c\mathfrak{D}_{D'} \subset \mathfrak{D}_D$ , the latter because  $c = [\mathfrak{D}_{D'} : \mathfrak{D}_D]$ . Thus,  $cL \in Id_n(\mathfrak{D}_D)$  because

$$N(cL) = c^2N(L) = c^2 = n,$$

and hence  $Id_n(\mathfrak{D}_D) = c\text{Ker}(\pi_{\mathfrak{D}_D, \mathfrak{D}_{D'}})$ , as claimed.  $\square$

*Acknowledgments.* First of all, I would like to thank the referee for the many detailed comments and references which improved the exposition of the paper. In addition, I thank Norm Hurt for his comments and notes on the first version of this paper, and for drawing my attention to the article of Sun and Williams. Finally, I would like to gratefully acknowledge receipt of funding from the Natural Sciences and Engineering Research Council of Canada (NSERC).

## REFERENCES

- [1] J. Antoniadis, *Diedergruppe und Reziprozitätsgesetz*, J. Reine Angew. Math. **377** (1987), 197–209.
- [2] D. Buell, *Binary Quadratic Forms*, Springer-Verlag, New York, 1989.
- [3] Z. Borevich, I. Shafarevich, *Number Theory*, Academic Press, New York, 1966.
- [4] N. Bourbaki, *Commutative Algebra, Ch. I-VII*. Addison-Wesley, Reading, 1972.
- [5] J. Buchmann, U. Vollmer, *Binary Quadratic Forms*, Springer-Verlag, Berlin, 2007.
- [6] D. Cox, *Primes of the Form  $x^2 + ny^2$ : Fermat, Class Field Theory and Complex Multiplication*, Wiley, New York, 1989.
- [7] H. Davenport and H. Heilbronn, *On the zeros of certain Dirichlet series I*, J. London Math. Soc. **11** (1936), 181–185. Reprinted in *The Collected Papers of Hans Arnold Heilbronn*, Wiley-Interscience, New York, 1988, pp. 272–276.
- [8] G. Lejeune Dirichlet, *Recherches sur diverses applications de l'analyse infinitésimale à la théorie des nombres*, J. Reine Angew. Math. **19** (1939), 324–369; **21** (1940), 1–12, 134–155. Reprinted in *Werke I*, Reimer, Berlin, 1889, pp. 413–496.
- [9] C. F. Gauss, *Untersuchungen über höhere Arithmetik*. Translation (1889) of *Disquisitiones Arithmeticae* (1801) by H. Maser. Reprint: Chelsea Publ. Co., New York, 1981.
- [10] H. Hasse, *Bemerkungen zu den Ring- und Strahlklasseneinteilungen in quadratischen Zahlkörpern*, Math. Nachr. **4** (1951), 322–327.
- [11] G. Hardy and E. Wright, *An Introduction to the Theory of Numbers* (Fourth Ed.), Oxford U. Press, Glasgow, 1968.
- [12] E. Hecke, *Zur Theorie der elliptischen Modulfunktionen*, Math. Ann. **97** (1926), 210–242. Reprinted in *Math. Werke*, Vandenhoeck & Ruprecht, Göttingen, 1983, pp. 428–460.
- [13] E. Hecke, *Über Modulfunktionen und die Dirichletschen Reihen mit Eulerscher Produktentwicklung I, II*, Math. Ann. **114** (1937), 1–28, 316–351. Reprinted in *Math. Werke*, Vandenhoeck & Ruprecht, Göttingen, 1983, pp. 644–707.
- [14] E. Hecke, *Analytische Arithmetik der positiven quadratischen Formen*, Kgl. Danske Videnskabernes Selskab. Math.-fysiske Meddelelser **XIII**, **12** (1940). Reprinted in *Math. Werke*, Vandenhoeck & Ruprecht, Göttingen, 1983, pp. 789–918.
- [15] H. Hijikata, A. Pizer and T. Schemanske, *The basis problem for modular forms on  $\Gamma_0(N)$* , Mem. Amer. Math. Soc. **82** (1989), no. 418, vi+159 pp.
- [16] E. Kani, *Idoneal numbers and some generalizations*, Ann. Sci. Math. Québec **35** (2011), no. 2, 197–227.
- [17] E. Kani, *Binary theta series and CM forms*, Intern. J. Number Theory (to appear).
- [18] Y. Kitaoka, *A note on Hecke operators and theta-series*, Nagoya Math. J. **42** (1971), 189–195.
- [19] Y. Kitaoka, *Representations of quadratic forms and their application to Selberg's zeta functions*, Nagoya Math. J. **63** (1976), 153–162.
- [20] L. Kronecker, *Zur Theorie der elliptischen Funktionen*, Sitzungsber. Königl. Preuss. Akad. Wiss. 1883–1889. Reprinted in *Werke*, vol. IV, Teubner, Leipzig, 1897, pp. 347–495.
- [21] S. Lang, *Elliptic Functions*, Addison-Wesley, Reading, MA, 1972.
- [22] T. Miyake, *Modular Forms*, Springer-Verlag, Berlin, 1989.

- [23] J. Neukirch, *Algebraische Zahlentheorie*, Springer-Verlag, Berlin, 1992.
- [24] J. Pöhler, *Über Primzahldarstellungen durch binäre quadratische Formen*, Math. Ann. **141** (1960), 239–241.
- [25] R. Schertz, *Complex Multiplication*, Cambridge U. Press, Cambridge, 2010.
- [26] B. Schoeneberg, *Das Verhalten mehrfacher Thetareihen bei Modulusubstitutionen*, Math. Ann. **116** (1939), 511–523.
- [27] B. Schoeneberg, *Elliptic Modular Functions*, Springer-Verlag, Berlin, 1974.
- [28] J-P. Serre, *Modular forms of weight one and Galois representations*, in Algebraic Number Fields: *L*-functions and Galois properties (Proc. Sympos., Univ. Durham, Durham, 1975), pp. 193–268. Academic Press, London, 1977. Reprinted in *Œuvres/Collected Papers III*, Springer-Verlag, Berlin, 1986, pp. 292–367.
- [29] C.-L. Siegel, *Über die analytische Theorie der quadratischen Formen*, Ann. Math. **36** (1935), 527–606. Reprinted in *Gesam. Abh. I*, Springer-Verlag, Berlin, 1966, pp. 322–405.
- [30] Z.-H. Sun and K. S. Williams, *On the number of representations of  $n$  by  $ax^2 + bxy + cy^2$* , Acta Arith. **122** (2006), no. 2, 101–171.
- [31] Z.-H. Sun and K. S. Williams, *Ramanujan identities and Euler products for a type of Dirichlet series*, Acta Arith. **122** (2006), no. 4, 349–393.
- [32] H. Weber, *Beweis des Satzes, dass jede eigentlich primitive quadratische Form unendlich viele Primzahlen darzustellen fähig ist*, Math. Ann. **20** (1882), no. 3, 301–329.
- [33] H. Weber, *Zahlentheoretische Untersuchungen aus dem Gebiete der elliptischen Funktionen III*, Nachrichten Königl. Gesell. Wiss. Univ. Göttingen **7** (1893), 245–264.
- [34] H. Weber, *Lehrbuch der Algebra*, Second edition, Vol. III, F. Vieweg und Sohn, Braunschweig, 1908. Reprint: Chelsea Pub. Co., New York, 1961.