

1. In each of the following cases, either prove that the given subset  $W$  is a subspace of  $V$ , or show why it is not a subspace of  $V$ .

- (a)  $V = C^\infty(\mathbb{R})$ ,  $W$  is the set of functions  $f \in V$  for which  $\lim_{x \rightarrow \infty} f(x) = 0$ .
- (b)  $V = C^\infty(\mathbb{R})$ ,  $W$  is the set of functions  $f \in V$  for which  $f(4) = 1$ .
- (c)  $V = M_{2 \times 2}(\mathbb{R})$ ,  $W$  is the set of matrices whose square is the zero matrix (i.e., those matrices  $A$  with  $A^2$  the zero matrix).
- (d)  $V = \mathbb{R}^\infty$ ,  $W$  is the subset of vectors  $(x_1, x_2, x_3, \dots)$  for which  $x_3 = x_2 + x_1$ ,  $x_4 = x_3 + x_2$ ,  $x_5 = x_4 + x_3$ ,  $\dots$ , and in general  $x_{n+2} = x_{n+1} + x_n$  for all  $n \geq 1$ .

2. In each of the following cases, either prove that the vectors are linearly independent, or show that they are linearly dependent.

- (a) The vectors  $v_1 = (1, 2)$  and  $v_2 = (3, 6)$  in  $W_2$ .
- (b) The vectors  $v_1 = \ln(x^2 + 1)$ ,  $v_2 = \ln(x^4 + 4x^2 + 3)$ , and  $v_3 = \ln(x^2 + 3)$  in  $C^\infty(\mathbb{R})$ .
- (c) The vectors  $v_1 = e^x$ ,  $v_2 = e^{3x}$ , and  $v_3 = \cos(x)$  in  $C^\infty(\mathbb{R})$ .

3. In each of the following cases, either prove that the given rule is a linear transformation, or show why it isn't.

- (a)  $T : \mathbb{R}^\infty \rightarrow \mathbb{R}^\infty$ ,  $T$  is the rule "shift to the right":  $T(x_1, x_2, x_3, \dots) = (0, x_1, x_2, x_3, \dots)$ .
- (b)  $T : M_{2 \times 3}(\mathbb{R}) \rightarrow \mathbb{R}^2$ ,  $T$  sends the matrix  $M$  to  $M\vec{w}$  where  $\vec{w} = (1, 2, 3)$  (the output is a vector in  $\mathbb{R}^2$ ).
- (c)  $T : W_2 \rightarrow \mathbb{R}^2$ ,  $T(x, y) = (x, y)$ .
- (d)  $T : W_2 \rightarrow \mathbb{R}^2$ ,  $T(x, y) = (\ln(x), \ln(y))$ .
- (e)  $T : C^\infty(\mathbb{R}) \rightarrow \mathbb{R}$ ,  $T(f) = \int_1^3 f \cdot \sin(x) dx$ .

The vector spaces used on the previous page are

$C^\infty(\mathbb{R})$ : The vector space of functions from  $\mathbb{R}$  to  $\mathbb{R}$  with infinitely many derivatives. Addition and scalar multiplication are addition and scalar multiplication of functions.

$\mathbb{R}^\infty$ : The set of infinite sequences  $(x_1, x_2, x_3, \dots)$  of numbers in  $\mathbb{R}$ , where addition and scalar multiplication are coordinate-wise.

$M_{m \times n}(\mathbb{R})$ : The set of  $m \times n$  matrices with entries in  $\mathbb{R}$ , where addition is addition of matrices and scalar multiplication is multiplying all entries of the matrix by that number.

$W_2$ : The following “weird” vector space:  $W_2$  is the set of all pairs  $(x, y)$  with both  $x > 0$  and  $y > 0$  real numbers, with addition defined as multiplication of the coordinates:

$$(x_1, y_1) + (x_2, y_2) = (x_1x_2, y_1y_2)$$

and scalar multiplication defined as exponentiation of the coordinates:

$$c \cdot (x, y) = (x^c, y^c),$$

for any  $(x_1, y_1), (x_2, y_2) \in W_2, c \in \mathbb{R}$ .

REMINDER: The zero vector in  $W_2$  is  $(1, 1)$ .

All of these vector spaces are vector spaces over  $\mathbb{R}$ .