

AN ANALOGUE OF LIOUVILLE'S THEOREM AND AN APPLICATION TO CUBIC SURFACES

DAVID MCKINNON AND MIKE ROTH

ABSTRACT. We prove a strong analogue of Liouville's Theorem in Diophantine approximation for points on arbitrary algebraic varieties. We use this theorem to prove a conjecture of the first author for cubic surfaces in \mathbb{P}^3 .

1. INTRODUCTION

The famous theorem of K.F. Roth (see for example [HS, Part D]) gives a sharp upper bound on how well an irrational algebraic number can be approximated by rational numbers. In [MR], the authors prove an analogue of Roth's Theorem for algebraic points on arbitrary algebraic varieties. In this paper we generalize, in the sense of [MR], Liouville's approximation theorem to arbitrary varieties, as well as giving an extension involving the stable base locus.

The point of view of [MR] is that the Roth and Liouville theorems are examples of "local Bombieri-Lang phenomena" whereby local positivity of a line bundle influences local accumulation of rational points. Specifically, given a variety X , an algebraic point $x \in X$, and an ample line bundle L on X , these theorems are expressed as inequalities between $\epsilon_x(L)$, the *Seshadri constant*, measuring local positivity of L near x , and $\alpha_x(L)$, an invariant measuring how well we can approximate x by rational points.

Roth's Theorem is usually thought of as stronger than Liouville's, but if the locus being approximated is defined over the ground field, Liouville's Theorem is strictly better. On \mathbb{P}^1 one gains a factor of two. For arbitrary varieties, however, moving past the Seshadri constant into the non-nef part of the big cone can provide even larger gains. We need this improvement for our application in §4 where we verify a conjecture of the first author for cubic surfaces in \mathbb{P}^3 .

In §2 we review the definitions and elementary properties of α_x and ϵ_x . In §3 we prove the generalized Liouville theorem (Theorem 3.3). We close the paper in §4 by computing α_x and ϵ_x for an arbitrary nef line bundle and rational point, not on a line, on a smooth cubic

2010 *Mathematics Subject Classification*. Primary 14G05; Secondary 14G40.

Key words and phrases. Seshadri constants, Diophantine approximation, rational points.

David McKinnon was partially supported by an NSERC research grant.

Mike Roth was partially supported by an NSERC research grant.

surface (where the lines are also rational); we then use this to verify Conjecture 3.2 from [McK].

Acknowledgements

We thank Kevin Hare for useful computations performed in the initial stages of this project. We are grateful to the referees of this paper who provided many useful suggestions on the exposition and advice on strengthening one of the results. Finally, we wish to acknowledge an intellectual debt to Mike Nakamaye, who has long advocated the point of view that Seshadri constants are diophantine.

2. ELEMENTARY PROPERTIES OF α AND ϵ

In this section, we give a brief overview of the properties of α and ϵ used in this paper. For a more detailed discussion of α , see [MR]. For a more detailed discussion of ϵ , there are many good references – see for example [Laz, chap. 5]. Proofs of all of the facts listed below can be found in [MR].

Let k be a number field, and X a projective variety over $\text{Spec}(k)$ (i.e., a reduced subscheme of some finite-dimensional projective space \mathbb{P}_k^r).

The constant α_x . In order to motivate the definition of α_x it is helpful to recall the classical case of approximation on the line. For a point $x \in \mathbb{R}$ the *approximation exponent* τ_x of x is the unique extended real number $\tau_x \in (0, \infty]$ such that the inequality

$$\left| x - \frac{a}{b} \right| \leq \frac{1}{b^{\tau_x + \delta}}$$

has only finitely many solutions $a/b \in \mathbb{Q}$ whenever $\delta > 0$ (respectively has infinitely solutions $a/b \in \mathbb{Q}$ whenever $\delta < 0$). The approximation exponent measures a certain tension between our ability to closely approximate x by rational numbers (the distance term $|x - a/b|$) and the complexity (the $1/b$ term) of the number required to make this approximation. In this notation the 1844 theorem of Liouville [L] is that $\tau_x \leq d$ for $x \in \mathbb{R}$ algebraic of degree d over \mathbb{Q} .

To generalize τ_x to arbitrary projective varieties over $\text{Spec}(k)$ for a number field k , we replace the function $|x - a/b|$ by a distance function $d_v(x, \cdot)$ depending on a place v of k , and measure the complexity of a point via a height function $H_L(\cdot)$ depending on an ample line bundle L . For an introduction to the theory of heights the reader is referred to any one of [BG, Chap. 2], [HS, Part B], [La, Chap. III], or [Se, Chap. 2]. Unless otherwise specified all height functions in this paper are multiplicative, relative to k , and come from line bundles on X defined over k . In this paper we use the following normalizations. The absolute values are normalized with respect to k : if v is a finite place of k , π a uniformizer of the corresponding maximal ideal, and κ the residue field then $\|\pi\|_v = 1/\#\kappa$; if v is an infinite place corresponding to an embedding $i: k \hookrightarrow \mathbb{C}$ then $\|x\|_v = \|i(x)\|^{m_v}$ for all $x \in k$, where

$m_v = 1$ or 2 depending on whether v is real or complex. The heights are then normalized so that for a point $x = [x_0 : \cdots : x_n] \in \mathbb{P}^n(k)$, the height with respect to $\mathcal{O}_{\mathbb{P}^n}(1)$ is

$$H(x) = \prod_v \max(\|x_0\|_v, \dots, \|x_n\|_v)$$

where the product ranges over all the places v of k .

In order to define a distance function we fix a place v of k and extension (which we also call v) to \bar{k} .

If v is archimedean. We choose a distance function on $X(\bar{k})$ by choosing an embedding $X \hookrightarrow \mathbb{P}_k^r$ and pulling back (via v) the function on $\mathbb{P}^r(\mathbb{C}) \times \mathbb{P}^r(\mathbb{C})$ given by the formula

$$d_v(x, y) = \left(1 - \frac{|\sum_{i=0}^r x_i \bar{y}_i|^2}{(\sum_{i=0}^r |x_i|^2)(\sum_{j=0}^r |y_j|^2)} \right)^{[k_v:\mathbb{R}]/2}$$

where $x = [x_0 : \cdots : x_r]$, and $y = [y_0 : \cdots : y_r]$ are points of $\mathbb{P}^r(\mathbb{C})$, and $|\cdot|$ is the absolute value on \mathbb{C} extending the usual absolute value on \mathbb{R} , i.e., such that $|3 + 4\sqrt{-1}| = 5$.

If v is non-archimedean. We choose a distance function on $X(\bar{k})$ by choosing an embedding $X \hookrightarrow \mathbb{P}_k^r$ and pulling back the distance function on $\mathbb{P}^r(\bar{k})$ given by the formula

$$d_v(x, y) = \frac{\max_{0 \leq i < j \leq r} (\|x_i y_j - x_j y_i\|_v)}{\max_{0 \leq i \leq r} (\|x_i\|_v) \max_{0 \leq j \leq r} (\|y_j\|_v)}$$

where $x = [x_0 : \cdots : x_r]$, and $y = [y_0 : \cdots : y_r]$ are points of $\mathbb{P}^r(\bar{k})$.

These definitions are standard in Arakelov theory, albeit here we have normalized with respect to k rather than \mathbb{Q} . (See for instance [BG, §2.8] where a distance function $\delta_v(\cdot, \cdot)$ is defined for each place v ; the distance functions are related by $d_v(\cdot, \cdot) = \delta_v(\cdot, \cdot)^{[k:\mathbb{Q}]}$.)

Two real valued functions g and g' with the same domain are called *equivalent* if there are positive constants $c \leq C$ such that $cg \leq g' \leq Cg$ for all values of the domain. The distance functions defined above depend on the choice of embedding into projective space, but by [MR, Proposition 2.4] any two embeddings give equivalent distance functions on $X(k_v) \times X(k_v)$. (We may restrict ourselves to points of $X(k_v)$ since any point of $X(\bar{k})$ that does not lie in $X(k_v)$ cannot be approximated by k -rational points; see also the remark on page 5.) It follows from the definition of α_x below that equivalent distance functions produce the same value of α_x ; thus our definition of α_x does not depend on the projective embedding chosen to define d_v . A more geometric definition of distance, and the proof that it is equivalent to the distance formulae above, may be found in §6.

Definition 2.1. Let X be a projective variety, $x \in X(\bar{k})$, L a line bundle on X . For any sequence $\{x_i\} \subset X(k)$ of distinct points with $d_v(x, x_i) \rightarrow 0$ (which we denote by $\{x_i\} \rightarrow x$), we set

$$A(\{x_i\}, L) = \left\{ \gamma \in \mathbb{R} \mid d_v(x, x_i)^\gamma H_L(x_i) \text{ is bounded from above} \right\}.$$

Remarks. (a) It follows easily from the definition that if $A(\{x_i\}, L)$ is nonempty then it is an interval unbounded to the right, i.e., if $\gamma \in A(\{x_i\}, L)$ then $\gamma + \delta \in A(\{x_i\}, L)$ for any $\delta > 0$.

(b) If $\{x'_i\}$ is a subsequence of $\{x_i\}$ then $A(\{x_i\}, L) \subseteq A(\{x'_i\}, L)$.

Definition 2.2. If $A(\{x_i\}, L)$ is empty we set $\alpha_x(\{x_i\}, L) = \infty$. Otherwise we set $\alpha_x(\{x_i\}, L)$ to be the infimum of $A(\{x_i\}, L)$. We call $\alpha_x(\{x_i\}, L)$ the approximation constant of $\{x_i\}$ with respect to L .

As $i \rightarrow \infty$ we have $d_v(x, x_i) \rightarrow 0$. We thus expect that $d_v(x, x_i)^\gamma H_L(x_i)$ goes to 0 for large γ and to ∞ for small γ . The number $\alpha_x(\{x_i\}, L)$ marks the transition point between these two behaviours.

By remark (b) above if $\{x'_i\}$ is a subsequence of $\{x_i\}$ then $\alpha_x(\{x'_i\}, L) \leq \alpha_x(\{x_i\}, L)$. Thus we may freely replace a sequence with a subsequence when trying to establish lower bounds.

Definition 2.3. Let k be a number field, X a projective variety over $\text{Spec}(k)$, L a line bundle on X , and $x \in X(\bar{k})$. Then $\alpha_x(L)$ is defined to be the infimum of all approximation constants of sequences of points in $X(k)$ converging to x . If no such sequence exists then set $\alpha_x(L) = \infty$.

To see the connection with the usual approximation exponent on \mathbb{P}^1 , suppose that L is an ample line bundle. We may define an approximation constant $\tau_x(L)$ by simply extending the definition on \mathbb{P}^1 , namely by defining $\tau_x(L)$ to be the unique extended real number $\tau_x(L) \in [0, \infty]$ such that the inequality

$$d_v(x, y) < \frac{1}{H_L(y)^{\tau_x(L) + \delta}}$$

has only finitely many solutions $y \in X(k)$ whenever $\delta > 0$ (respectively has infinitely many solutions $y \in X(k)$ whenever $\delta < 0$). Then [MR, Proposition 2.11] implies that $\alpha_x(L) = \frac{1}{\tau_x(L)}$. In particular the theorem of Liouville becomes $\alpha_x(\mathcal{O}_{\mathbb{P}^1}(1)) \geq \frac{1}{d}$ for $x \in \mathbb{R}$ of degree d over \mathbb{Q} , and it is this type of lower bound that we wish to generalize to arbitrary varieties. The choice of using the reciprocal of τ is justified by the resulting formal similarity with the Seshadri constant, and more natural behaviour when we vary L (see, for example, Proposition 2.9).

We need two results on α_x before continuing onto the Seshadri constant. First, we will need to know how to calculate α_x in one simple case.

Lemma 2.4. Let $x \in \mathbb{P}^n(k)$. Then $\alpha_{x, \mathbb{P}^n}(\mathcal{O}_{\mathbb{P}^n}(1)) = 1$.

Proof: This is Lemma 2.13 from [MR]. \square

Second, it will be useful to know how the approximation constant changes when we change the field k . Let K/k be a finite extension with $K \subset \bar{k}$, and set $X_K := X \times_k K$. A point $y \in X(\bar{k})$ is a map $\text{Spec}(\bar{k}) \rightarrow X$ over $\text{Spec}(k)$, and factors through the map $X_K \rightarrow X$, i.e., such

a point x gives a point of $X_K(\bar{k})$. We thus have a canonical identification $X_K(\bar{k}) = X(\bar{k})$ which we use implicitly in the paragraphs below. We use the notation that $\alpha_x(\{x_i\}, L)_K$ (respectively $\alpha_x(L)_K$) denotes the approximation constant of a sequence (resp. point x) computed on X_K with respect to K . This means that when computing α , we use the height H_L relative to K and normalize d_v relative to K (we define d_v on X_K using the same embedding used to define d_v on X). If $d = [K:k]$ and $m_v = [K_v:k_v]$ (where K_v and k_v denote the completions of K and k with respect to v) then this means simply that $H_L(x_i)_K = H_L(x_i)_k^d$ and $d_v(x, x_i)_K = d_v(x, x_i)_k^{m_v}$.

Proposition 2.5. *Let X be a variety over $\text{Spec}(k)$, $x \in X(\bar{k})$ any point, L a line bundle on X , and $\{x_i\} \rightarrow x$ a sequence of points in $X(k)$ approximating x . Let K be any finite extension of k and set $X_K := X \times_k K$. The base change of $\{x_i\}$ gives a sequence $\{y_i\}$ of K -points of X_K approximating x . Set $m_v = [K_v:k_v]$, and let $d = [K:k]$. Then*

$$\alpha_x(\{y_i\}, L)_K = \frac{d}{m_v} \alpha_x(\{x_i\}, L)_k.$$

In particular, we have the bound $\alpha_x(L)_K \leq \frac{d}{m_v} \alpha_x(L)_k$.

Proof: The claim that $\alpha_x(\{x_i\}, L)_K = \frac{d}{m_v} \alpha_x(\{x_i\}, L)_k$ follows immediately from the equalities $H_L(\cdot)_K = H_L(\cdot)_k^d$ and $d_v(\cdot, \cdot)_K = d_v(\cdot, \cdot)_k^{m_v}$. The inequality $\alpha_x(L)_K \leq \frac{d}{m_v} \alpha_x(L)_k$ then follows since the sequences in $X_K(K)$ approximating x which come from sequences $\{x_i\}$ in $X(k)$ are a subset of all the sequences in $X_K(K)$ approximating x . \square

Remark. Let x be a point of $X(\bar{k})$ and let K be the field of definition of x . If $K \not\subseteq k_v$, or equivalently, $K_v \neq k_v$ then it will be impossible to find a sequence of points of $X(k)$ converging (in terms of d_v) to x . For example, when v is archimedean this happens when $k_v = \mathbb{R}$ and $K_v = \mathbb{C}$. Thus, if we can approximate x by points of $X(k)$ we may assume that $K_v = k_v$ and so $m_v = 1$.

The following result (appearing in [MR] as Theorem 2.16 and in [McK] as Theorem 2.8, although the latter version is incorrectly stated) is obtained by combining the Roth and Dirichlet theorems for approximation on \mathbb{P}^1 , as well as the local information about the singularity type. It shows how to calculate α_x on any singular k -rational curve.

Theorem 2.6. *Let C be any singular k -rational curve and $\varphi: \mathbb{P}^1 \rightarrow C$ the normalization map. Then for any ample line bundle L on C , and any $x \in C(\bar{k})$ we have the equality:*

$$\alpha_{x,C}(L) = \min_{q \in \varphi^{-1}(x)} d/r_q m_q$$

where $d = \deg(L)$, m_q is the multiplicity of the branch of C through x corresponding to q , and

$$r_q = \begin{cases} 0 & \text{if } \kappa(q) \not\subseteq k_v \\ 1 & \text{if } \kappa(q) = k \\ 2 & \text{otherwise.} \end{cases}$$

Here $\kappa(q)$ means the residue field of the point q , and we use $r_q = 0$ as a shorthand for $d/r_q m_q = \infty$.

The Seshadri constant. The Seshadri constant was introduced by Demailly in [D] for the purposes of measuring the local positivity of a line bundle.

Definition 2.7. *Let X be a projective variety, x a point of X , and L a nef line bundle on X . The Seshadri constant, $\epsilon_x(L)$, is defined to be*

$$\epsilon_x(L) := \sup \{ \gamma \geq 0 \mid \pi^*L - \gamma E \text{ is nef} \}$$

where $\pi: \tilde{X} \rightarrow X$ is the blowup of X at x , with exceptional divisor E .

A basic property of the Seshadri constant is that if L is ample (as opposed to just being nef) then $\pi^*L - \gamma E$ is itself ample on \tilde{X} for all rational $\gamma \in (0, \epsilon_x(L))$. An argument for this appears in the original paper of Demailly [D] defining the Seshadri constant¹. \square

In the discussion of Conjecture 4.2 below we will need the following alternate characterization of the Seshadri constant:

Proposition 2.8. *With the same setup as definition 2.7,*

$$\epsilon_x(L) = \inf_{x \in C \subseteq X} \left\{ \frac{(L \cdot C)}{\text{mult}_x(C)} \right\}$$

where the infimum is taken over all reduced irreducible curves C passing through x .

Proof: This is [Laz, Proposition 5.1.5]. \square

In order to indicate the parallels between α_x and ϵ_x , and for use below, we list a few of their formal properties here.

Proposition 2.9. *Let X be a projective variety over $\text{Spec}(k)$, $x \in X(\bar{k})$, and let L be any nef line bundle on X .*

- (a) *For any positive integer m , $\alpha_x(mL) = m\alpha_x(L)$ and $\epsilon_x(mL) = m\epsilon_x(L)$. (Thus α and ϵ also make sense for nef \mathbb{Q} -divisors.)*
- (b) *α_x and ϵ_x are concave functions of L : for any positive rational numbers a and b , and any \mathbb{Q} -divisors L_1 and L_2 (again defined over k , but with the exception of the case that $\{\alpha_x(L_1), \alpha_x(L_2)\} = \{-\infty, \infty\}$) we have*

$$\alpha_x(aL_1 + bL_2) \geq a\alpha_x(L_1) + b\alpha_x(L_2)$$

and

$$\epsilon_x(aL_1 + bL_2) \geq a\epsilon_x(L_1) + b\epsilon_x(L_2)$$

where for the last inequality we assume that L_1 and L_2 are nef.

- (c) *If Z is a subvariety of X over $\text{Spec}(k)$ then for any point $z \in Z(\bar{k})$ we have $\alpha_z(L|_Z) \geq \alpha_{z,X}(L)$ and $\epsilon_z(L|_Z) \geq \epsilon_z(L)$.*

¹This property appears as the statement “ $F_{p,q}$ is ample whenever $p > q/\epsilon_x(L)$ ” on page 98 of [D].

(d) If Y is also a variety over $\text{Spec}(k)$, $x \in X(k)$, $y \in Y(k)$ and L_X and L_Y are nef line bundles on X and Y respectively then

$$\alpha_{x \times y, X \times Y}(L_X \boxplus L_Y) = \min(\alpha_{x, X}(L_X), \alpha_{y, Y}(L_Y))$$

and

$$\epsilon_{x \times y, X \times Y}(L_X \boxplus L_Y) = \min(\epsilon_{x, X}(L_X), \epsilon_{y, Y}(L_Y)).$$

Note that by $L_X \boxplus L_Y$ we mean the line bundle $pr_X^* L_1 + pr_Y^* L_2$ on $X \times Y$, where pr_X and pr_Y are the projections. We prefer additive notation for line bundles since this is in line with the behaviour of α_x and ϵ_x , and hence use $L_X \boxplus L_Y$ rather than $L_1 \boxtimes L_2$.

Proof: All the proofs follow from elementary arguments using the definitions. For the statements about α_x see [MR, Proposition 2.14], and for the statements about ϵ_x see [MR, Proposition 3.4]. \square

3. A LIOUVILLE LOWER BOUND FOR α

In this section, as in the previous one, we fix a number field k and let X be a projective variety over $\text{Spec}(k)$.

Lemma 3.1. *Let x be a point of $X(k)$, and $\pi: \tilde{X} \rightarrow X$ the blow up of X at x with exceptional divisor E . Choose an embedding $\varphi: X \hookrightarrow \mathbb{P}^n$ so that $x \mapsto [1:0:\dots:0]$. Let Z_0, \dots, Z_n be the coordinates on \mathbb{P}^n and define functions u_i , $i = 1, \dots, n$ on the open subset where $Z_0 \neq 0$ by $u_i = Z_i/Z_0$.*

For each place v of k , define a function $e_v: X(k_v) \rightarrow \mathbb{R}_{\geq 0}$ by

$$e_v(y) = \begin{cases} 1 & \text{if } Z_0(y) = 0, \\ \min(1, \max(\|u_1(y)\|_v, \dots, \|u_n(y)\|_v)) & \text{if } Z_0(y) \neq 0. \end{cases}$$

Then

- (a) $0 \leq e_v \leq 1$ for all places v .
- (b) $e_v(\cdot)$ is equivalent to $d_v(x, \cdot)$ as a function on $X(k_v)$.
- (c) For $y \in X(k)$, $y \neq x$, we have $H_E(y) = (\prod_w e_w(y))^{-1}$.

Proof: Part (a) is clear from the definition. We now prove (b). By [MR, Proposition 2.4], distance functions associated to different projective embeddings of X are equivalent on $X(k_v) \times X(k_v)$, so we may assume that d_v has been defined using the embedding φ . If v is non-archimedean, then the formula for the distance function in the non-archimedean case and the fact that x is sent to $[1:0:\dots:0]$ give

$$d_v(x, y) = \frac{\max(\|Z_1(y)\|_v, \|Z_2(y)\|_v, \dots, \|Z_r(y)\|_v)}{\max(\|Z_0(y)\|_v, \|Z_1(y)\|_v, \dots, \|Z_r(y)\|_v)} \text{ for all } y \in X(\bar{k}).$$

For $y \in X(k_v)$, this is equal to $e_v(y)$.

In the case that v is archimedean, we may further assume that $k_v = \mathbb{C}$, since the functions to be compared transform the same way under field extensions. From the formula for the distance in the archimedean case and the fact that x is sent to $[1:0:\cdots:0]$ we obtain

$$\begin{aligned} d_v(x, y) &= 1 - \frac{|y_0|^2}{|y_0|^2 + \cdots + |y_r|^2} = \frac{|y_1|^2 + \cdots + |y_r|^2}{|y_0|^2 + |y_1|^2 + \cdots + |y_r|^2} \\ &= \frac{\|y_1\|_v + \cdots + \|y_r\|_v}{\|y_0\|_v + \|y_1\|_v + \cdots + \|y_r\|_v}. \end{aligned}$$

For $y \in U(k_v)$, $y_0 \neq 0$, and $u_j(y) = y_j/y_0$ for $j = 1, \dots, r$. Thus $d_v(x, y) = \frac{\|u_1(y)\|_v + \cdots + \|u_r(y)\|_v}{1 + \|u_1(y)\|_v + \cdots + \|u_r(y)\|_v}$; it is then elementary to check that

$$\frac{1}{r} d_v(x, y) \leq \max(\|u_1(y)\|_v, \dots, \|u_n(y)\|_v) \leq 2 d_v(x, y),$$

for all $y \in U(k_v)$. For $y \in X(k_v) \setminus U(k_v)$ we have $1 = d_v(x, y) = e_v(y)$, and thus $d_v(x, \cdot)$ is equivalent to $e_v(\cdot)$ on $X(k_v)$.

In (c) we are considering points $y \in X(k)$, $y \neq x$ also to be points of $\tilde{X}(k)$ via the birational map π . To prove (c) it suffices, by using the functoriality of heights under pullback, to consider the case that $X = \mathbb{P}^n$. Then the blow up $\tilde{\mathbb{P}}^n$ of \mathbb{P}^n at x is a subvariety of $\mathbb{P}^n \times \mathbb{P}^{n-1}$ and $\mathcal{O}_{\tilde{\mathbb{P}}^n}(E)$ is the restriction of $\mathcal{O}_{\mathbb{P}^n \times \mathbb{P}^{n-1}}(1, -1)$ to $\tilde{\mathbb{P}}^n$. From this description of $\mathcal{O}_{\tilde{\mathbb{P}}^n}(E)$ we obtain the formula

$$H_E(y) = \prod_w \frac{\max(\|Z_0(y)\|_w, \|Z_1(y)\|_w, \dots, \|Z_n(y)\|_w)}{\max(\|Z_1(y)\|_w, \dots, \|Z_n(y)\|_w)}$$

from which (c) follows easily. \square

For the following lemma, we will need an additional definition. The stable base locus of a line bundle L is the intersection of the base loci of mL as $m \rightarrow \infty$. For details, see Definition 2.1.20 of [Laz].

Lemma 3.2. *Suppose that $x \in X(k)$ and let $\pi: \tilde{X} \rightarrow X$ be the blow up at x with exceptional divisor E . Let L be a line bundle on X and $\gamma > 0$ a rational number such that $L_\gamma := \pi^*L - \gamma E$ is in the effective cone of \tilde{X} . Let B' be the stable base locus of L_γ and set $B = \pi(B')$.*

Then there is a positive real constant M (depending only on x and L) such that for any sequence of k -points $\{x_i\} \rightarrow x$ with all points of $\{x_i\}$ outside of B , we have

$$H_L(x_i) d_v(x, x_i)^\gamma \geq M$$

In particular, $\alpha(\{x_i\}, L) \geq \gamma$.

Proof: Let $U = \tilde{X} \setminus B'$. Since B' is the stable base locus of L_γ there is a constant c (depending only on x and L) so that $H_{L_\gamma}(y) \geq c$ for all $y \in U(k)$. Applying Lemma 3.1 we then have

$$\begin{aligned} c \leq H_{L_\gamma}(x_i) &= H_L(x_i) H_E(x_i)^{-\gamma} \stackrel{3.1(c)}{=} H_L(x_i) \left(\prod_w e_w(x_i) \right)^\gamma \\ &\stackrel{3.1(a)}{\leq} H_L(x_i) e_v(x_i)^\gamma. \end{aligned}$$

By Lemma 3.1(b) $d_v(x, x_i)$ and $e_v(x_i)$ are equivalent functions on $X(k)$ and therefore $H_L(x_i)d_v(x, x_i)^\gamma \geq M$ for some positive constant M , again depending only on x and L .

For any $\delta > 0$ we thus have $H_L(x_i)d_v(x, x_i)^{\gamma-\delta} \geq Md_v(x, x_i)^{-\delta}$ and so conclude that $\gamma - \delta \notin A(\{x_i\}, L)$ since $c'd_v(x, x_i)^{-\delta} \rightarrow \infty$ as $i \rightarrow \infty$. Therefore $\gamma \leq \alpha(\{x_i\}, L)$. \square

The main result of this section is the following implication of Lemma 3.2.

Theorem 3.3 (Liouville-type theorem). *Let X be an algebraic variety over $\text{Spec}(k)$, $x \in X(\bar{k})$ any point, and set $d = [K : k]$ where K is the field of definition of x .*

*Set $X_K := X \times_k K$, let \tilde{X} be the blowup of X_K at x with exceptional divisor E , and set π to be the composite $\pi: \tilde{X} \rightarrow X_K \rightarrow X$. Let L be a nef line bundle on X , and $\gamma > 0$ a rational number such that $L_\gamma := \pi^*L - \gamma E$ is in the effective cone of \tilde{X} . Finally let B' be the stable base locus of L_γ and set $B = \pi(B')$. Then there is a positive real constant M such that for all $y \in X(k) - B(k)$, we have $H_L(y)d_v(x, y)^{\gamma/d} \geq M$, and*

- (a) *For any sequence $\{x_i\} \rightarrow x$ of k -points approximating x , if infinitely many points of $\{x_i\}$ are outside B then $\alpha(\{x_i\}, L) \geq \gamma/d$.*
- (b) *If $\alpha_x(L) < \gamma/d$ then $x \in B$ and $\alpha_x(L) = \alpha_x(L|_B)$.*
- (c) *If $x \in B$ and $\alpha_x(L|_B) \geq \gamma/d$ then $\alpha_x(L) \geq \gamma/d$.*

Furthermore, there is a subvariety Y of X such that $x \in Y$ and for all $y \in X(k)$, we have $H_L(y)d_v(x, y)^{\gamma/d} \geq M$, provided that $(L - \gamma E)|_Y$ is in the effective cone of Y .

Proof: Let $\{x_i\}$ be a sequence approximating x . If infinitely many x_i lie outside of B then we may pass to the subsequence of points outside of B , which can only approximate the point x better than the sequence as a whole. To prove part (a) we may therefore assume that all points of $\{x_i\}$ lie outside B . Base changing $\{x_i\}$ we obtain a sequence $\{y_i\}$ in $X_K(K)$ approximating $x \in X_K(K)$. Applying Lemma 3.2 to X_K we conclude that $H_L(y_i)d_v(x, y_i)^\gamma \geq M$ and $\alpha(\{y_i\}, L)_K \geq \gamma$. Since there is a sequence of k -points approximating x we conclude by the remark on page 5 that (in the notation of Proposition 2.5) $m_v = 1$. Therefore by Proposition 2.5, $H_L(y_i)d_v(x, y_i)^{\gamma/d} \geq M$ and $\alpha(\{x_i\}, L)_k = \frac{1}{d}\alpha(\{y_i\}, L)_K \geq \gamma/d$, proving (a).

If $\alpha_x(L) < \gamma/d$ then there must be a sequence $\{x_i\}$ approximating x such that $\alpha(\{x_i\}, L) < \gamma/d$. By part (a) this implies that all but finitely many x_i lie in B . Thus $x \in B$ since B is closed. Since omitting finitely many elements of a sequence does not change the approximation constant we may assume that all x_i are contained in B . Since $\alpha_{x, X}(L)$ is the infimum of the approximation constants for sequences $\{x_i\}$ with $\alpha(\{x_i\}, L) < \gamma/d$ we conclude that $\alpha_x(L) = \alpha_x(L|_B)$ proving (b).

If $\alpha_x(L) < \gamma/d$ then part (b) along with the hypothesis for part (c) lead to an immediate contradiction. Thus, under the hypotheses of part (c), $\alpha_x(L) \geq \gamma/d$.

The final remark is obvious in cases (a) and (c), and in case (b), we may replace X with B and L with $L|_B$ and apply Theorem 3.3 again. Iterating this, we deduce the desired result. \square

Remark. Theorem 3.3 still holds if we replace B by the Zariski closure of $B(k)$. This has the added advantage that every component of B is then absolutely irreducible (see [MR, Lemma 2.17]).

Corollary 3.4. *Suppose that L is a nef line bundle on X such that $\pi^*L - \gamma E$ has empty stable base locus for all rational $\gamma \in (0, \epsilon_x(L))$. Then $\alpha_x(L) \geq \epsilon_x(L)/d$.*

Proof: For all rational $\gamma \in (0, \epsilon_x(L))$ the stable base locus of $\pi^*L - \gamma E$ is empty by hypothesis, and thus by Theorem 3.3(a) we conclude that $\alpha_x(L) \geq \gamma/d$ for any such γ , and hence that $\alpha_x(L) \geq \epsilon_x(L)/d$. \square .

Corollary 3.5. *If L is an ample line bundle, then $\alpha_x(L) \geq \epsilon_x(L)/d$.*

Proof: For all rational $\gamma \in (0, \epsilon_x(L))$ the line bundle $\pi^*L - \gamma E$ is ample on \tilde{X} . (See the discussion after Definition 2.7 for a proof of this.) In particular, the stable base locus of $\pi^*L - \gamma E$ is empty, so the result follows by Corollary 3.4. \square

Remark. If $X = \mathbb{P}^1$ then Corollary 3.5 and the fact that $\epsilon_x(\mathcal{O}_{\mathbb{P}^1}(1)) = 1$ give $\alpha_x(\mathcal{O}_{\mathbb{P}^1}(1)) \geq 1/d$. Thus on \mathbb{P}^1 Corollary 3.5 amounts to the classic Liouville bound $\tau_x \leq d$. Liouville's original result also includes an explicit relation between the height and distance of approximating rational points, and Theorem 3.3 has this feature as well. For this reason we consider Theorem 3.3 and its corollaries to be “Liouville bounds” for α_x .

The effective cone is usually larger than the nef cone, and in general the parts of Theorem 3.3 imply a much stronger lower bound for $\alpha_x(L)$ than Corollary 3.4. We will use this in the next section to compute α for the cubic surface, but give a brief illustration now by calculating α for rational points of a non-split quadric surface in \mathbb{P}^3 . (A split quadric surface is isomorphic to $\mathbb{P}^1 \times \mathbb{P}^1$, and $\alpha_x(\mathcal{O}_{\mathbb{P}^1 \times \mathbb{P}^1}(a, b)) = \min(a, b)$ when $a, b > 0$, as implied by Proposition 2.9(d) and computed in both [McK, Theorem 3.1] and [MR, §2; Example (b) just before Lemma 2.17].)

Example. Let X be a smooth quadric surface in \mathbb{P}^3 defined over k , and set $L = \mathcal{O}_{\mathbb{P}^3}(1)|_X$. We assume that no lines on X are defined over k . Let x be a k -point of X . By intersecting with a (rationally defined) hyperplane we may find a conic C passing through x such that C is isomorphic to \mathbb{P}^1 over k . By Lemma 2.4 and Proposition 2.9(a,c), we therefore have $\alpha_{x,X}(L) \leq \alpha_{x,C}(L|_C) = \alpha_{x,\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(2)) = 2$. Since x lies on a line (over \bar{k}), we have $\epsilon_x(L) = 1$, and applying Corollary 3.4 we obtain $\alpha_x(L) \geq 1$. Thus $1 \leq \alpha_x(L) \leq 2$, i.e., Corollary 3.4 does not give enough information to determine $\alpha_x(L)$ in this case.

However, let $\pi: \tilde{X} \rightarrow X$ be the blow up of X at x with exceptional divisor E . Then $\pi^*L - 2E$ is effective with base locus the proper transform of the two lines passing through x . In particular the image B of this base locus is the union of the two lines of ruling passing through x . Since (by assumption) neither of these lines is defined over k , x is the only k -point of B . Thus by Theorem 3.3(a) if $\{x_i\}$ is any sequence of k -points approximating x then $\alpha(\{x_i\}, L) \geq 2$, and in particular $\alpha_x(L) \geq 2$. Thus $\alpha_x(L) = 2$ for all k -points of X .

Since X is non-split the Picard group of X (over k) has rank one with generator L . Thus the above computation and the homogeneity in Proposition 2.9(a) determines α for all $x \in X(k)$ and all ample line bundles on X defined over k .

4. THE CUBIC SURFACE

In this section, we will compute α_x and ϵ_x for all k -rational points x on the blowup X of \mathbb{P}^2 at six k -rational points in general position.

To begin, we will recall some notions from [McK].

Definition 4.1. *A sequence $\{x_i\} \rightarrow x$ whose approximation constant is equal to $\alpha_x(L)$ (if such a sequence exists) is called a sequence of best approximation to x . A curve C passing through x is called a curve of best approximation (with respect to L) if C contains a sequence of best approximation to x .*

In other words, if C is a curve of best approximation to x on X , then the rational points on C approximate x roughly as well as the rational points on X approximate x .

Note also that for a point x on a curve C and an ample divisor L on C , there is always a sequence $\{x_i\}$ such that $\alpha_x(\{x_i\}, L) = \alpha_x(L)$. Thus, in particular, C is a curve of best approximation if and only if $\alpha_x(L|_C) = \alpha_x(L)$.

In the example of the non-split quadric — and in many others considered in [McK] — there is always a curve of best approximation to x . In [McK, §4] it is shown that if Vojta's main conjectures (see [V] for statements) are true, then $\alpha_x(L)$ finite implies that $\alpha_x(L)$ is computed on a subvariety $V \subseteq X$ of negative Kodaira dimension (possibly X itself, if X has negative Kodaira dimension). Since varieties of negative Kodaira dimension are (again, conjecturally) covered by rational curves, one is led to the following further prediction ([McK, Conjecture 2.7]):

Conjecture 4.2. *Let X be an algebraic variety defined over k , and L any ample divisor on X . Let x be any k -rational point on X and assume that there is a rational curve defined over k passing through x . Then there exists a curve C (necessarily rational) of best approximation to x on X with respect to L .*

In [McK], the first author proves this conjecture in many cases, and shows that in many others it follows from Vojta's Conjecture. Those proofs use a slightly different definition of α , but the proofs do not essentially change in the new setting.

The Seshadri-constant analogue of a curve of best approximation is called a *Seshadri curve* (cf. Proposition 2.8):

Definition 4.3. *Let L be a nef divisor on an algebraic variety X , and $x \in X$ any point. A Seshadri curve for x with respect to L is a curve C such that $\epsilon_{x,X}(L) = (L \cdot C) / \text{mult}_x(C)$.*

In all currently known examples, there exists a Seshadri curve for x with respect to L , but it is conjectured that this is not always the case. In particular, it is possible that the Seshadri constant might sometimes be irrational (see [Laz, Remark 5.1.13]).

It is useful to know that for a fixed curve C , the set of line bundles for which C is a curve of best approximation form a subcone of the Néron-Severi group, and similarly for the property of being a Seshadri curve.

Proposition 4.4. *Let X be a variety over $\text{Spec}(k)$, and let $x \in X(k)$ be any k -rational point. Let D_1 and D_2 be nef divisors on X with height functions H_1 and H_2 bounded below by a positive constant in some neighbourhood of x . Let a_1 and a_2 be non-negative integers, and let $D = a_1D_1 + a_2D_2$.*

- (a) *If C is a curve of best approximation for D_1 and D_2 , then C is also a curve of best approximation for D .*
- (b) *If C is a Seshadri curve for x with respect to D_1 and D_2 , then C is also a Seshadri curve for x with respect to D .*

Proof: Part (a) appears as [McK, Corollary 3.2]. To prove part (b), note that Proposition 2.9(b) implies the estimate

$$\epsilon_x(a_1D_1 + a_2D_2) \geq a_1\epsilon_x(D_1) + a_2\epsilon_x(D_2).$$

On the other hand, the hypotheses of part (b) give

$$\begin{aligned} \frac{C \cdot D}{\text{mult}_x(C)} &= \frac{C \cdot (a_1D_1 + a_2D_2)}{\text{mult}_x(C)} = \frac{a_1(C \cdot D_1)}{\text{mult}_x(C)} + \frac{a_2(C \cdot D_2)}{\text{mult}_x(C)} \\ &= a_1\epsilon_x(D_1) + a_2\epsilon_x(D_2). \end{aligned}$$

Thus, by Proposition 2.8, $a_1\epsilon_x(D_1) + a_2\epsilon_x(D_2)$ is an upper bound for $\epsilon_x(D)$. Therefore $\epsilon_x(a_1D_1 + a_2D_2) = a_1\epsilon_x(D_1) + a_2\epsilon_x(D_2)$ and C is a Seshadri curve for D , proving (b). \square

We are now ready to begin the proof of the main result of this section. Before we state and prove the general result, we will illustrate the fundamental techniques in the case $L = -K$.

Theorem 4.5. *Let X be a smooth cubic surface in \mathbb{P}^3 defined over k , and isomorphic over k to the blowup of \mathbb{P}^2 at six k -rational points in general position. Let $x \in X(k)$ be any k -rational point, and let C_x be the curve of intersection of X with the tangent plane to X at*

x. Then

$$\epsilon_x(-K) = \begin{cases} 1 & \text{if } x \text{ lies on one of the 27 lines of } X; \\ \frac{3}{2} & \text{otherwise,} \end{cases}$$

while

$$\alpha_x(-K) = \begin{cases} 1 & \text{if } x \text{ lies on one of the 27 lines of } X; \\ \frac{3}{2} & \text{if } x \text{ is not on one of the 27 lines, and if either} \\ & \quad \circ C_x \text{ is cuspidal at } x, \text{ or} \\ & \quad \circ C_x \text{ is nodal at } x \text{ with tangent lines having slopes} \\ & \quad \text{in } k_v \text{ but not } k; \\ 2 & \text{otherwise.} \\ & \text{(i.e., } C_x \text{ is nodal at } x, \text{ and the slopes of the tangent} \\ & \text{lines are in } k \text{ or not in } k_v.) \end{cases}$$

Proof: Set $L = -K = \mathcal{O}_{\mathbb{P}^3}(1)|_X$, and let x be a point of $X(k)$. If x lies on a line ℓ then by Proposition 2.9(c) we have $\epsilon_{x,\ell}(L|_\ell) \geq \epsilon_{x,X}(L) \geq \epsilon_{x,\mathbb{P}^3}(\mathcal{O}_{\mathbb{P}^3}(1))$. Since $\epsilon_{x,\ell}(L|_\ell) = \epsilon_{x,\mathbb{P}^3}(\mathcal{O}_{\mathbb{P}^3}(1)) = 1$, we conclude that $\epsilon_x(L) = 1$. Similarly (using Proposition 2.9(c) again and the fact that $\alpha_{x,\ell}(L|_\ell) = \alpha_{x,\mathbb{P}^3}(\mathcal{O}_{\mathbb{P}^3}(1)) = 1$ by Lemma 2.4) we conclude that $\alpha_x(L) = 1$.

We now suppose that x does not lie on a line. Let $\pi: \tilde{X} \rightarrow X$ be the blowup of X at x , with exceptional divisor E . Then C_x is a Seshadri curve for x with respect to L . To see this, note first that C_x satisfies $C_x \cdot L / \text{mult}_x(C_x) = 3/2$, so $\epsilon_x(L) \leq 3/2$. Conversely, if $a > 3/2$, then $\pi^*L - aE$ is not nef, because $(\pi^*L - aE)(\pi^*L - 2E) = 3 - 2a < 0$ and $\pi^*L - 2E$ is the class of the proper transform of C_x . Thus, $\epsilon_x(L) \geq 3/2$, and so $\epsilon_x(L) = 3/2$, and C_x is a Seshadri curve for x with respect to L .

We now turn to the computation of α . The stable base locus of $\pi^*L - 2E$ is \tilde{C}_x , the proper transform of C_x . Hence by Theorem 3.3(b) either $\alpha_x(L) \geq 2$ or $\alpha_x(L) = \alpha_{x,C_x}(L|_{C_x})$ (note that $d = 1$). By intersecting X with a hyperplane containing x and one of the lines, we produce a k -rational conic passing through x , and approximating on the conic gives us $2 \geq \alpha_x(L)$. We therefore conclude that $\alpha_x(L) = \min(2, \alpha_{x,C_x}(L|_{C_x}))$.

The curve C_x is singular at x , and since x does not lie on a line, C_x is also irreducible. In particular, C_x is an irreducible curve of geometric genus zero, and since x is defined over k , C_x is birational to \mathbb{P}^1 over k , via projection from x in the tangent plane.

Applying Theorem 2.6 to C_x , we find that

$$\alpha_{x,C_x}(L|_{C_x}) = \begin{cases} \frac{3}{2} & \text{if } C_x: \quad \circ \text{ is cuspidal, or} \\ & \quad \circ \text{ is nodal and the tangent lines have slopes in } k_v \\ & \quad \text{but not in } k; \\ 3 & \text{if } C_x \text{ is nodal and the tangent lines have slopes in } k; \\ \infty & \text{if } C_x \text{ is nodal and the tangent lines do not have} \\ & \text{slopes in } k_v, \end{cases}$$

and this implies the stated values of $\alpha_x(L)$ above. \square

We now treat the case of a general nef divisor D . In what follows, we assume that the point x does not lie on a (-1) -curve on X . We begin with a calculation of the Seshadri constant ϵ . To do this, we will need some notation.

Let $\varphi: X \rightarrow \mathbb{P}^2$ be the blowing down map, and let E_1, \dots, E_6 be the exceptional divisors of φ . We define the following linear equivalence classes on X :

- $L = \varphi^* \mathcal{O}(1)$
- $L_i = L - E_i$, the strict transform of a line through $P_i = \varphi(E_i)$
- $L_{ij} = 2L - (\sum E_n) + E_i + E_j$ (where $i \neq j$), the strict transform of a conic through the four points P_n with $n \neq i, j$
- $B_i = 3L - (\sum E_n) - E_i$, the strict transform of a cubic curve through all six points P_n , with a node at P_i .

Let h be the class of a hyperplane in the anticanonical embedding $X \subset \mathbb{P}^3$. For any line ℓ on X , the hyperplanes containing ℓ give (after removing ℓ) a base-point-free pencil on X . If $x \in X$ does not lie on a line then the unique curve in this pencil through x is smooth and irreducible. The classes $\{L_i, L_{ij}, B_i\}$ defined above are the 27 pencils coming from the lines. Recall that for any point x on X we use C_x for the intersection of X with its tangent plane at x (so C_x has class h). If x does not lie on a line, then C_x is a plane cubic curve with one double point, at x .

Theorem 4.6. *Let x be a point on X that does not lie on a (-1) -curve, and let D be a nef divisor on X . The Seshadri constant $\epsilon_x(D)$ is equal to $\min\{D \cdot L_i, D \cdot L_{ij}, D \cdot B_i, (D \cdot h)/2\}$.*

Proof: The nef cone Γ of X has 99 generators, which are listed in §5, Table 1. Let S be the set of 27 divisor classes $\{L_i, L_{ij}, B_i\}$ as i and j range over all possible values, and for each element C in S , we define the subcone $\Gamma(C)$ by:

$$\Gamma(C) = \left\{ D \in \Gamma \mid D \cdot C = \min_{C' \in S} \{D \cdot C'\} \text{ and } D \cdot C \leq (D \cdot h)/2 \right\}.$$

Further define the subcone $\Gamma(h)$ to be:

$$\Gamma(h) = \left\{ D \in \Gamma \mid (D \cdot h)/2 \leq \min_{C' \in S} \{D \cdot C'\} \right\}.$$

It is clear that Γ is the union of these 28 subcones. To prove Theorem 4.6, it suffices to show that for every subcone $\Gamma(C)$, with $C \in S$, the curve through x in the pencil corresponding to C is a Seshadri curve for x with respect to D for all $D \in \Gamma(C)$ (respectively, in the case of the subcone $\Gamma(h)$, that C_x is a Seshadri curve for x with respect to D for all $D \in \Gamma(h)$). By Proposition 4.4(b) it further suffices to prove this for D a generator of the cone $\Gamma(C)$ (respectively $\Gamma(h)$).

The fundamental group of the space of all smooth cubic surfaces acts via monodromy on the Néron-Severi lattice of X . This monodromy action preserves the hyperplane class h and acts transitively on the classes of the lines. Thus, up to monodromy action, there are

only two of these subcones: $\Gamma(L_1)$ and $\Gamma(h)$. Generators for each of these subcones can be found in §5. Let $F = F_{x,L_1}$ be the unique curve in the pencil L_1 passing through x . For each generator D of $\Gamma(L_1)$, it is straightforward to verify that F is a Seshadri curve for x with respect to D . These verifications also appear in §5. Each generator G of $\Gamma(h)$ is also a generator of one of the other twenty-seven subcones $\Gamma(C)$, and for each such G , we have $G \cdot C = (G \cdot h)/2 = (G \cdot C_x)/\text{mult}_x(C_x)$. Thus, since C is a Seshadri curve for x with respect to G , it follows that C_x is also a Seshadri curve for x with respect to G , and so C_x is a Seshadri curve for every element of $\Gamma(h)$. \square

The next step is to calculate α_x for a point on a cubic surface.

Theorem 4.7. *Let $x \in X(k)$ be a point that does not lie on a (-1) -curve, and let D be a nef divisor on X . If the tangent curve C_x is a cuspidal cubic, or a nodal cubic whose tangent lines at x are defined over k_v but not defined over k , then $\alpha_x(D) = \epsilon_x(D)$. Otherwise, $\alpha_x(D) = \min\{D \cdot L_i, D \cdot L_{ij}, D \cdot B_i\}$.*

Proof: Suppose that D is in one of the cones $\Gamma(C)$ for $C \in S$, and let $F_{x,C}$ be the element of the pencil corresponding to C which passes through x . Since $F_{x,C}$ is a smooth k -rational curve, we have

$$D \cdot C \stackrel{2.6}{=} \alpha_x(D|_{F_{x,C}}) \stackrel{2.9(c)}{\geq} \alpha_x(D) \stackrel{3.4}{\geq} \epsilon_x(D) \stackrel{4.6}{=} D \cdot C,$$

where, reading from left to right, the equalities and inequalities are given by Theorem 2.6, Proposition 2.9(c), Corollary 3.4, and Theorem 4.6 respectively. (Note that the blowup of X at a point not on a line is a del Pezzo surface of degree two, and so every nef divisor on the blowup is semiample (see Theorem 5.1.2.1 of [ADHL]), and thus has nonempty stable base locus. In particular the hypotheses of Corollary 3.4 are satisfied.) Thus $\alpha_x(D) = D \cdot C$ and $F_{x,C}$ is a curve of best approximation with respect to D .

Now suppose that $D \in \Gamma(h)$. If C_x is cuspidal, or nodal with tangent lines having slopes in k_v but not k , then Theorem 2.6 gives $\alpha_x(D|_{C_x}) = D \cdot C_x/2 = D \cdot C_x/\text{mult}_x(C_x)$. By Theorem 4.6, $\epsilon_x(D) = D \cdot C_x/2$, and so as above we conclude that $\alpha_x(D) = D \cdot C_x = \epsilon_x(D)$, and that C_x is a curve of best approximation for D .

We now assume that C_x is nodal and the slopes of the tangent lines are in k or not in k_v . The codimension one faces of $\Gamma(h)$ (i.e., the facets) occur where one of the inequalities defining $\Gamma(h)$ becomes an equality, so that each facet is the intersection of $\Gamma(h)$ and $\Gamma(C)$ for some $C \in S$. For each $C \in S$ set $\hat{\Gamma}(C)$ to be the cone generated by $\Gamma(C)$ and $-K$. Since $-K$ is in the interior of $\Gamma(h)$ it follows that Γ is the union of the $\hat{\Gamma}(C)$, $C \in S$.

As above, for any $C \in S$, let $F_{x,C}$ be the member of the pencil corresponding to C passing through x . In the proof of Theorem 4.5 we have seen that $F_{x,C}$ is a curve of best approximation for $-K$, and in the first part of the argument above that $F_{x,C}$ is a curve of best

approximation for all $D \in \Gamma(C)$. By Proposition 4.4(a) we conclude that $F_{x,C}$ is a curve of best approximation for all $D \in \hat{\Gamma}(C)$. The result follows. \square

Note that as part of the proof we have shown that Conjecture 4.2 holds for every point $x \in X$ not on a (-1) -curve.

5. APPENDIX A: GENERATORS OF NEF CONES AND SUBCONES FOR THE CUBIC SURFACE

A version of this appendix, with additional tables and larger font, may be found at [Por]. In all the tables below, the first column is a numerical identifier of the vector in that row. In the tables 1, 2a, and 3 the subsequent columns represent the coefficients of the vector with respect to the basis $\{L, E_1, \dots, E_6\}$ of the Néron-Severi group of X . Thus, vector number 1 in Table 1 is the divisor class $2L - E_1 - E_2 - E_3$. Each of the cones has 99 generators. There is no correspondence or relation between the rows in tables 1, 2a, and 3 with the same numerical identifier.

Table 1, of generators of the nef cone, is reproducing information that has been well known for some time, of course. It was calculated for these tables by finding generators for the cone obtained as the intersection of the half-spaces corresponding to non-negative intersection with each of the 27 lines on the cubic surface.

Table 1: Generators of the nef cone Γ of a smooth cubic surface

#	L	E_1	E_2	E_3	E_4	E_5	E_6	#	L	E_1	E_2	E_3	E_4	E_5	E_6
1	2	-1	-1	-1	0	0	0	51	3	-1	-2	0	-1	-1	-1
2	2	-1	-1	0	-1	0	0	52	3	-1	-1	-2	-1	-1	0
3	2	-1	-1	0	0	-1	0	53	3	-1	-1	-2	-1	0	-1
4	2	-1	-1	0	0	0	-1	54	3	-1	-1	-2	0	-1	-1
5	2	-1	0	-1	-1	0	0	55	3	-1	-1	-1	-2	-1	0
6	2	-1	0	-1	0	-1	0	56	3	-1	-1	-1	-2	0	-1
7	2	-1	0	-1	0	0	-1	57	3	-1	-1	-1	-1	-2	0
8	2	-1	0	0	-1	-1	0	58	3	-1	-1	-1	-1	0	-2
9	2	-1	0	0	-1	0	-1	59	3	-1	-1	-1	0	-2	-1
10	2	-1	0	0	0	-1	-1	60	3	-1	-1	-1	0	-1	-2
11	1	0	0	0	0	0	0	61	3	-1	-1	0	-2	-1	-1
12	3	-2	-1	-1	-1	-1	0	62	3	-1	-1	0	-1	-2	-1
13	3	-2	-1	-1	-1	0	-1	63	3	-1	-1	0	-1	-1	-2
14	3	-2	-1	-1	0	-1	-1	64	3	-1	0	-2	-1	-1	-1
15	3	-2	-1	0	-1	-1	-1	65	3	-1	0	-1	-2	-1	-1
16	3	-2	0	-1	-1	-1	-1	66	3	-1	0	-1	-1	-2	-1
17	1	-1	0	0	0	0	0	67	3	-1	0	-1	-1	-1	-2
18	1	0	-1	0	0	0	0	68	3	0	-2	-1	-1	-1	-1
19	1	0	0	-1	0	0	0	69	3	0	-1	-2	-1	-1	-1
20	1	0	0	0	-1	0	0	70	3	0	-1	-1	-2	-1	-1
21	1	0	0	0	0	-1	0	71	3	0	-1	-1	-1	-2	-1
22	1	0	0	0	0	0	-1	72	3	0	-1	-1	-1	-1	-2
23	2	0	-1	-1	-1	0	0	73	3	-2	-1	-1	-1	-1	-1
24	2	0	-1	-1	0	-1	0	74	3	-1	-2	-1	-1	-1	-1
25	2	0	-1	-1	0	0	-1	75	3	-1	-1	-2	-1	-1	-1
26	2	0	-1	0	-1	-1	0	76	3	-1	-1	-1	-2	-1	-1
27	2	0	-1	0	-1	0	-1	77	3	-1	-1	-1	-1	-2	-1
28	2	0	-1	0	0	-1	-1	78	3	-1	-1	-1	-1	-1	-2
29	2	0	0	-1	-1	-1	0	79	4	-2	-2	-2	-1	-1	-1
30	2	0	0	-1	-1	0	-1	80	4	-2	-2	-1	-2	-1	-1
31	2	0	0	-1	0	-1	-1	81	4	-2	-2	-1	-1	-2	-1
32	2	0	0	0	-1	-1	-1	82	4	-2	-2	-1	-1	-1	-2
33	2	-1	-1	-1	-1	0	0	83	4	-2	-1	-2	-2	-1	-1
34	2	-1	-1	-1	0	-1	0	84	4	-2	-1	-2	-1	-2	-1
35	2	-1	-1	-1	0	0	-1	85	4	-2	-1	-2	-1	-1	-2
36	2	-1	-1	0	-1	-1	0	86	4	-2	-1	-1	-2	-2	-1
37	2	-1	-1	0	-1	0	-1	87	4	-2	-1	-1	-2	-1	-2
38	2	-1	-1	0	0	-1	-1	88	4	-2	-1	-1	-1	-2	-2
39	2	-1	0	-1	-1	-1	0	89	4	-1	-2	-2	-2	-1	-1
40	2	-1	0	-1	-1	0	-1	90	4	-1	-2	-2	-1	-2	-1
41	2	-1	0	-1	0	-1	-1	91	4	-1	-2	-2	-1	-1	-2
42	2	-1	0	0	-1	-1	-1	92	4	-1	-2	-1	-2	-2	-1
43	2	0	-1	-1	-1	-1	0	93	4	-1	-2	-1	-2	-1	-2
44	2	0	-1	-1	-1	0	-1	94	4	-1	-2	-1	-1	-2	-2
45	2	0	-1	-1	0	-1	-1	95	4	-1	-1	-2	-2	-2	-1
46	2	0	-1	0	-1	-1	-1	96	4	-1	-1	-2	-2	-1	-2
47	2	0	0	-1	-1	-1	-1	97	4	-1	-1	-2	-1	-2	-2
48	3	-1	-2	-1	-1	-1	0	98	4	-1	-1	-1	-2	-2	-2
49	3	-1	-2	-1	-1	0	-1	99	5	-2	-2	-2	-2	-2	-2
50	3	-1	-2	-1	0	-1	-1								

Table 2a, of generators of the cone $\Gamma(L_1)$, was generated by using the half-spaces defining Γ in addition to the half-spaces corresponding to the conditions $D \cdot L_1 = \min_{C' \in S} \{D \cdot C'\}$ and $D \cdot L_1 \leq (D \cdot h)/2$ for all $D \in \Gamma$.

Table 2a: Generators D_n of the cone $\Gamma(L_1)$

#	L	E_1	E_2	E_3	E_4	E_5	E_6	#	L	E_1	E_2	E_3	E_4	E_5	E_6
1	4	-3	-1	-1	-1	-1	-1	51	4	-2	-2	-1	-1	0	-1
2	2	-1	-1	0	0	0	0	52	4	-2	-2	-1	0	-1	-1
3	2	-1	0	-1	0	0	0	53	4	-2	-2	0	-1	-1	-1
4	2	-1	0	0	-1	0	0	54	4	-2	-1	-2	-1	-1	0
5	2	-1	0	0	0	-1	0	55	4	-2	-1	-2	-1	0	-1
6	2	-1	0	0	0	0	-1	56	4	-2	-1	-2	0	-1	-1
7	1	0	0	0	0	0	0	57	4	-2	-1	-1	-2	-1	0
8	3	-2	-1	-1	-1	0	0	58	4	-2	-1	-1	-2	0	-1
9	3	-2	-1	-1	0	-1	0	59	4	-2	-1	-1	-1	-2	0
10	3	-2	-1	-1	0	0	-1	60	4	-2	-1	-1	-1	0	-2
11	3	-2	-1	0	-1	-1	0	61	4	-2	-1	-1	0	-2	-1
12	3	-2	-1	0	-1	0	-1	62	4	-2	-1	-1	0	-1	-2
13	3	-2	-1	0	0	-1	-1	63	4	-2	-1	0	-2	-1	-1
14	3	-2	0	-1	-1	-1	0	64	4	-2	-1	0	-1	-2	-1
15	3	-2	0	-1	-1	0	-1	65	4	-2	-1	0	-1	-1	-2
16	3	-2	0	-1	0	-1	-1	66	4	-2	0	-2	-1	-1	-1
17	3	-2	0	0	-1	-1	-1	67	4	-2	0	-1	-2	-1	-1
18	1	-1	0	0	0	0	0	68	4	-2	0	-1	-1	-2	-1
19	2	-1	-1	-1	0	0	0	69	4	-2	0	-1	-1	-1	-2
20	2	-1	-1	0	-1	0	0	70	4	-2	-2	-1	-1	-1	-1
21	2	-1	-1	0	0	-1	0	71	4	-2	-1	-2	-1	-1	-1
22	2	-1	-1	0	0	0	-1	72	4	-2	-1	-1	-2	-1	-1
23	2	-1	0	-1	-1	0	0	73	4	-2	-1	-1	-1	-2	-1
24	2	-1	0	-1	0	-1	0	74	4	-2	-1	-1	-1	-1	-2
25	2	-1	0	-1	0	0	-1	75	5	-2	-2	-2	-1	-1	-1
26	2	-1	0	0	-1	-1	0	76	5	-2	-2	-1	-2	-1	-1
27	2	-1	0	0	-1	0	-1	77	5	-2	-2	-1	-1	-2	-1
28	2	-1	0	0	0	-1	-1	78	5	-2	-2	-1	-1	-1	-2
29	3	-1	-1	-1	-1	0	0	79	5	-2	-1	-2	-2	-1	-1
30	3	-1	-1	-1	0	-1	0	80	5	-2	-1	-2	-1	-2	-1
31	3	-1	-1	-1	0	0	-1	81	5	-2	-1	-2	-1	-1	-2
32	3	-1	-1	0	-1	-1	0	82	5	-2	-1	-1	-2	-2	-1
33	3	-1	-1	0	-1	0	-1	83	5	-2	-1	-1	-2	-1	-2
34	3	-1	-1	0	0	-1	-1	84	5	-2	-1	-1	-1	-2	-2
35	3	-1	0	-1	-1	-1	0	85	5	-3	-2	-2	-1	-1	-1
36	3	-1	0	-1	-1	0	-1	86	5	-3	-2	-1	-2	-1	-1
37	3	-1	0	-1	0	-1	-1	87	5	-3	-2	-1	-1	-2	-1
38	3	-1	0	0	-1	-1	-1	88	5	-3	-2	-1	-1	-1	-2
39	3	-1	-1	-1	-1	-1	0	89	5	-3	-1	-2	-2	-1	-1
40	3	-1	-1	-1	-1	0	-1	90	5	-3	-1	-2	-1	-2	-1
41	3	-1	-1	-1	0	-1	-1	91	5	-3	-1	-2	-1	-1	-2
42	3	-1	-1	0	-1	-1	-1	92	5	-3	-1	-1	-2	-2	-1
43	3	-1	0	-1	-1	-1	-1	93	5	-3	-1	-1	-2	-1	-2
44	4	-1	-1	-1	-1	-1	-1	94	5	-3	-1	-1	-1	-2	-2
45	3	-2	-1	-1	-1	-1	0	95	6	-3	-2	-2	-2	-2	-1
46	3	-2	-1	-1	-1	0	-1	96	6	-3	-2	-2	-2	-1	-2
47	3	-2	-1	-1	0	-1	-1	97	6	-3	-2	-2	-1	-2	-2
48	3	-2	-1	0	-1	-1	-1	98	6	-3	-2	-1	-2	-2	-2
49	3	-2	0	-1	-1	-1	-1	99	6	-3	-1	-2	-2	-2	-2
50	4	-2	-2	-1	-1	-1	0								

We use D_n to refer to the divisor class represented by row n of Table 2a. For any point $x \in X$ not on a (-1) -curve, the unique curve $F := F_{x,L_1}$ in the pencil L_1 passing through x is smooth and irreducible. Row n in Table 2b below is a — very brief! — justification of why F is a Seshadri curve for x with respect to D_n .

Table 2b: Reasons F is a Seshadri curve for D_n

#	Reason	#	Reason	#	Reason	#	Reason
1	$L_1.D_1 = 1$	26	$L_1.D_{26} = 1$	51	$D_{46} + L_2$	76	$L_2 + L_{23} + L_{56}$
2	$L_1.D_2 = 1$	27	$L_1.D_{27} = 1$	52	$D_{47} + L_2$	77	$L_2 + L_{23} + L_{46}$
3	$L_1.D_3 = 1$	28	$L_1.D_{28} = 1$	53	$D_{48} + L_2$	78	$L_2 + L_{23} + L_{45}$
4	$L_1.D_4 = 1$	29	$L + L_{56}$	54	$D_{45} + L_3$	79	$L_3 + L_{23} + L_{56}$
5	$L_1.D_5 = 1$	30	$L + L_{46}$	55	$D_{46} + L_3$	80	$L_3 + L_{23} + L_{46}$
6	$L_1.D_6 = 1$	31	$L + L_{45}$	56	$D_{47} + L_3$	81	$L_3 + L_{23} + L_{45}$
7	$L_1.D_7 = 1$	32	$L + L_{36}$	57	$D_{45} + L_4$	82	$L_4 + L_{23} + L_{46}$
8	$L_1.D_8 = 1$	33	$L + L_{35}$	58	$D_{46} + L_4$	83	$L_4 + L_{23} + L_{45}$
9	$L_1.D_9 = 1$	34	$L + L_{34}$	59	$D_{45} + L_5$	84	$L_5 + L_{23} + L_{45}$
10	$L_1.D_{10} = 1$	35	$L + L_{26}$	60	$D_{46} + L_6$	85	$D_{45} + L_{45}$
11	$L_1.D_{11} = 1$	36	$L + L_{25}$	61	$D_{47} + L_5$	86	$D_{45} + L_{35}$
12	$L_1.D_{12} = 1$	37	$L + L_{24}$	62	$D_{47} + L_6$	87	$D_{45} + L_{34}$
13	$L_1.D_{13} = 1$	38	$L + L_{23}$	63	$D_{48} + L_4$	88	$D_{46} + L_{34}$
14	$L_1.D_{14} = 1$	39	$L_2 + L_{26}$	64	$D_{48} + L_5$	89	$D_{45} + L_{25}$
15	$L_1.D_{15} = 1$	40	$L_2 + L_{25}$	65	$D_{48} + L_6$	90	$D_{45} + L_{24}$
16	$L_1.D_{16} = 1$	41	$L_2 + L_{24}$	66	$D_{49} + L_3$	91	$D_{46} + L_{24}$
17	$L_1.D_{17} = 1$	42	$L_2 + L_{23}$	67	$D_{49} + L_4$	92	$D_{45} + L_{23}$
18	$L_1.D_{18} = 0$	43	$L_3 + L_{23}$	68	$D_{49} + L_5$	93	$D_{46} + L_{23}$
19	$L_1.D_{19} = 1$	44	$L_{23} + L_2 + L_3$	69	$D_{49} + L_6$	94	$D_{47} + L_{23}$
20	$L_1.D_{20} = 1$	45	$L_1.D_{45} = 1$	70	$B_1 + L_2$	95	$L_{23} + L_{46} + L_{56}$
21	$L_1.D_{21} = 1$	46	$L_1.D_{46} = 1$	71	$B_1 + L_3$	96	$L_{23} + L_{45} + L_{56}$
22	$L_1.D_{22} = 1$	47	$L_1.D_{47} = 1$	72	$B_1 + L_4$	97	$L_{23} + L_{45} + L_{46}$
23	$L_1.D_{23} = 1$	48	$L_1.D_{48} = 1$	73	$B_1 + L_5$	98	$L_{23} + L_{34} + L_{56}$
24	$L_1.D_{24} = 1$	49	$L_1.D_{49} = 1$	74	$B_1 + L_6$	99	$L_{23} + L_{24} + L_{56}$
25	$L_1.D_{25} = 1$	50	$D_{45} + L_2$	75	$L_3 + L_{34} + L_{56}$		

As an example, in row 1 of Table 2b, the “Reason” is $L_1.D_1 = 1$, and thus $F.D_1 = L_1.D_1 = 1$. We claim that for the divisors D_n , ϵ_x is always at least one if it is nonzero. Furthermore, if $n \neq 18$ then $\epsilon_x(D_n) \geq 1$ for x not on a (-1) -curve. Granting these claims, since by assumption, x does not lie on any (-1) -curve (and since $D_1 \neq D_{18}$), we have $\epsilon_x(D_1) \geq 1$. The curve F is smooth at x , and the reason given tells us that F has degree 1 with respect to D_1 . Therefore $(D_1 \cdot F)/\text{mult}_x(F) = 1/1 = 1$, hence $\epsilon_x(D_1) = 1$, and F is a Seshadri curve for x with respect to D_1 .

To see the claims, notice that the generators of the nef cone Γ (see Table 1) are all either morphisms to \mathbb{P}^1 corresponding to pencils of conics on the cubic surface, or else morphisms to \mathbb{P}^2 that are the blowing down of six pairwise disjoint (-1) -curves. For a point x not on a (-1) -curve ϵ_x is at least one for the blowdowns to \mathbb{P}^2 , and $\epsilon_x = 0$ for the pencils no matter which point x is. It is straightforward to check that all the generators listed in Table 2a are non-negative integer linear combinations of the generators of the nef cones, and therefore (by Proposition 2.9(b)) enjoy the same property: for any point x , the Seshadri constant $\epsilon_x(D_i)$ is either zero or else is at least one. Furthermore the only generator of $\Gamma(L_1)$ which is morphism to \mathbb{P}^1 is $D_{18} = L_1$, and thus we have $\epsilon_x(D_n) \geq 1$ for all $n \neq 18$.

Similar arguments explain other reasons of the form “ $D_n \cdot L_1 = 1$ ” or “ $=0$ ”.

As a second example of a reason, in row 29 of Table 2b, the comment “ $L + L_{56}$ ” means that the divisor D_{29} represented by that row is the sum of L and L_{56} . Again suppose that x does not lie on a (-1) -curve. Any curve C passing through x that has nonzero intersection with L must have $L.C/\text{mult}_x(C) \geq 1$, since L is an isomorphism away from (-1) -curves. Similarly,

any curve not contracted by L_{56} must also satisfy $L_{56}.C/\text{mult}_x(C) \geq 1$, so any curve not contracted by L_{56} or L must satisfy $(L + L_{56}).C/\text{mult}_x(C) \geq 2$. If C is contracted by L_{56} , then it is either a (-1) -curve, or else it is an element of the divisor class L_{56} itself, in which case it satisfies $(L + L_{56}).C/\text{mult}_x(C) = 2$ by direct calculation. In all cases, since x does not lie on a (-1) -curve, we see that $\epsilon_x(L + L_{56}) \geq 2$. Since $L_1.L = L_1.L_{56} = 1$, we compute that $(L + L_{56}).F/\text{mult}_x(F) = (1 + 1)/1 = 2$ and therefore that $\epsilon_x(L + L_{56}) = 2$. Hence F is a Seshadri curve for x with respect to $D_{29} = L + L_{56}$. Similar arguments explain the other reasons of the form “ $a + b$ ” or “ $a + b + c$ ”.

For these types of arguments, it is useful to know that L_1 (and hence F) has intersection number one with the divisors L , B_1 , L_i for $i \neq 1$, and L_{ij} for $i, j \neq 1$.

Table 3, of generators of the cone $\Gamma(h)$, was generated by using the half-spaces defining Γ in addition to the half-spaces corresponding to the intersection inequalities $(D \cdot h)/2 \leq \min_{C' \in S} \{D \cdot C'\}$ for all $D \in \Gamma$.

Table 3: Generators G_n of the cone $\Gamma(h)$

#	L	E_1	E_2	E_3	E_4	E_5	E_6	Div. Class	#	L	E_1	E_2	E_3	E_4	E_5	E_6	Div. Class
1	8	-3	-3	-3	-3	-3	-3	B_1	51	7	-2	-3	-2	-3	-3	-2	B_2
2	4	-1	-1	-1	-1	-1	-1	L_1	52	7	-2	-3	-2	-3	-2	-3	B_2
3	4	-2	-2	-1	-1	-1	-1	L_1	53	7	-2	-3	-2	-2	-3	-3	B_2
4	4	-2	-1	-2	-1	-1	-1	L_1	54	7	-2	-2	-3	-3	-3	-2	B_3
5	4	-2	-1	-1	-2	-1	-1	L_1	55	7	-2	-2	-3	-3	-2	-3	B_3
6	4	-2	-1	-1	-1	-2	-1	L_1	56	7	-2	-2	-3	-2	-3	-3	B_3
7	4	-2	-1	-1	-1	-1	-2	L_1	57	7	-2	-2	-2	-3	-3	-3	B_4
8	4	-1	-2	-2	-1	-1	-1	L_2	58	6	-3	-2	-2	-2	-2	-1	B_1
9	4	-1	-2	-1	-2	-1	-1	L_2	59	6	-3	-2	-2	-2	-1	-2	B_1
10	4	-1	-2	-1	-1	-2	-1	L_2	60	6	-3	-2	-2	-1	-2	-2	B_1
11	4	-1	-2	-1	-1	-1	-2	L_2	61	6	-3	-2	-1	-2	-2	-2	B_1
12	4	-1	-1	-2	-2	-1	-1	L_3	62	6	-3	-1	-2	-2	-2	-2	B_1
13	4	-1	-1	-2	-1	-2	-1	L_3	63	6	-2	-3	-2	-2	-2	-1	B_2
14	4	-1	-1	-2	-1	-1	-2	L_3	64	6	-2	-3	-2	-2	-1	-2	B_2
15	4	-1	-1	-1	-2	-2	-1	L_4	65	6	-2	-3	-2	-1	-2	-2	B_2
16	4	-1	-1	-1	-2	-1	-2	L_4	66	6	-2	-3	-1	-2	-2	-2	B_2
17	4	-1	-1	-1	-1	-2	-2	L_5	67	6	-2	-2	-3	-2	-2	-1	B_3
18	5	-2	-2	-2	-1	-1	-1	L_1	68	6	-2	-2	-3	-2	-1	-2	B_3
19	5	-2	-2	-1	-2	-1	-1	L_1	69	6	-2	-2	-3	-1	-2	-2	B_3
20	5	-2	-2	-1	-1	-2	-1	L_1	70	6	-2	-2	-2	-3	-2	-1	B_4
21	5	-2	-2	-1	-1	-1	-2	L_1	71	6	-2	-2	-2	-3	-1	-2	B_4
22	5	-2	-1	-2	-2	-1	-1	L_1	72	6	-2	-2	-2	-2	-3	-1	B_5
23	5	-2	-1	-2	-1	-2	-1	L_1	73	6	-2	-2	-2	-2	-1	-3	B_6
24	5	-2	-1	-2	-1	-1	-2	L_1	74	6	-2	-2	-2	-1	-3	-2	B_5
25	5	-2	-1	-1	-2	-2	-1	L_1	75	6	-2	-2	-2	-1	-2	-3	B_6
26	5	-2	-1	-1	-2	-1	-2	L_1	76	6	-2	-2	-1	-3	-2	-2	B_4
27	5	-2	-1	-1	-1	-2	-2	L_1	77	6	-2	-2	-1	-2	-3	-2	B_5
28	5	-1	-2	-2	-2	-1	-1	L_2	78	6	-2	-2	-1	-2	-2	-3	B_6
29	5	-1	-2	-2	-1	-2	-1	L_2	79	6	-2	-1	-3	-2	-2	-2	B_3
30	5	-1	-2	-2	-1	-1	-2	L_2	80	6	-2	-1	-2	-3	-2	-2	B_4

(Table 3 continued)

#	L	E_1	E_2	E_3	E_4	E_5	E_6	Div. Class	#	L	E_1	E_2	E_3	E_4	E_5	E_6	Div. Class
31	5	-1	-2	-1	-2	-2	-1	L_2	81	6	-2	-1	-2	-2	-3	-2	B_5
32	5	-1	-2	-1	-2	-1	-2	L_2	82	6	-2	-1	-2	-2	-2	-3	B_6
33	5	-1	-2	-1	-1	-2	-2	L_2	83	6	-1	-3	-2	-2	-2	-2	B_2
34	5	-1	-1	-2	-2	-2	-1	L_3	84	6	-1	-2	-3	-2	-2	-2	B_3
35	5	-1	-1	-2	-2	-1	-2	L_3	85	6	-1	-2	-2	-3	-2	-2	B_4
36	5	-1	-1	-2	-1	-2	-2	L_3	86	6	-1	-2	-2	-2	-3	-2	B_5
37	5	-1	-1	-1	-2	-2	-2	L_4	87	6	-1	-2	-2	-2	-2	-3	B_6
38	7	-3	-3	-3	-2	-2	-2	B_1	88	5	-2	-2	-2	-2	-2	-1	B_1
39	7	-3	-3	-2	-3	-2	-2	B_1	89	5	-2	-2	-2	-2	-1	-2	B_1
40	7	-3	-3	-2	-2	-3	-2	B_1	90	5	-2	-2	-2	-1	-2	-2	B_1
41	7	-3	-3	-2	-2	-2	-3	B_1	91	5	-2	-2	-1	-2	-2	-2	B_1
42	7	-3	-2	-3	-3	-2	-2	B_1	92	5	-2	-1	-2	-2	-2	-2	B_1
43	7	-3	-2	-3	-2	-3	-2	B_1	93	5	-1	-2	-2	-2	-2	-2	B_2
44	7	-3	-2	-3	-2	-2	-3	B_1	94	3	-1	-1	-1	-1	-1	0	L_1
45	7	-3	-2	-2	-3	-3	-2	B_1	95	3	-1	-1	-1	-1	0	-1	L_1
46	7	-3	-2	-2	-3	-2	-3	B_1	96	3	-1	-1	-1	0	-1	-1	L_1
47	7	-3	-2	-2	-2	-3	-3	B_1	97	3	-1	-1	0	-1	-1	-1	L_1
48	7	-2	-3	-3	-3	-2	-2	B_2	98	3	-1	0	-1	-1	-1	-1	L_1
49	7	-2	-3	-3	-2	-3	-2	B_2	99	3	0	-1	-1	-1	-1	-1	L_2
50	7	-2	-3	-3	-2	-2	-3	B_2									

We use G_n to refer to the divisor class represented by row n of Table 3. The rightmost column of row n is a divisor class $C \in S$ such that G_n is also a generator of the subcone $\Gamma(C)$. From the definition of the cones $\Gamma(C)$ and $\Gamma(h)$, this implies that $G_n \cdot C = (G_n \cdot h)/2$. As explained in the proof of Theorem 4.7, this provides a verification that C_x is a Seshadri curve for x with respect to G_n .

6. APPENDIX B: GEOMETRIC DESCRIPTION OF THE DISTANCE FUNCTIONS

In this section we give a more geometric definition of the distance function, and prove its equivalence to the ones given by the formulae in §2.

Given a variety X defined over k let $\widetilde{X \times X}$ be the blowup of $X \times X$ along the diagonal, with exceptional divisor E . Let v be a place of k extended to \bar{k} (which we also call v), and choose a v -adic metric on $\mathcal{O}_{\widetilde{X \times X}}(E)$, i.e., a (non-trivial) v -adic norm $|\cdot|_v$ on each fibre of $\mathcal{O}_{\widetilde{X \times X}}(E) \otimes k_v$, varying continuously with the points of $(\widetilde{X \times X})(k_v)$ (see e.g., [CS, p. 162] for a discussion of v -adic metrics). We fix a nonzero global section s_E of $\mathcal{O}_{\widetilde{X \times X}}(E)$ with divisor E . Given points $x, y \in X(k)$ with $x \neq y$, then (x, y) is a point of $X \times X$ not on the diagonal, and hence corresponds to a unique point, which we also label (x, y) of $\widetilde{X \times X}$. We then define a distance function $d'_v(\cdot, \cdot)$ by

$$d'_v(x, y) = \begin{cases} |s_E(x, y)|_v & \text{if } x \neq y \\ 0 & \text{if } x = y. \end{cases}$$

Different choices of s_E differ by a scalar, and induce equivalent distance functions. Similarly, different choices of v -adic metrics differ multiplicatively on $\widetilde{X \times X}$ by a bounded function (see [CS, p. 162, Lemma 7.1]), and again induce equivalent distance functions.

The distance functions $d_v(\cdot, \cdot)$ in §2 are obtained by restricting distance functions on \mathbb{P}^r under an embedding $X \hookrightarrow \mathbb{P}_k^r$. We may also view the distance function $d'_v(\cdot, \cdot)$ defined above as being induced from a distance function on projective space, and this will allow us to reduce the problem of showing the equivalence of the two types of distance functions to the case $X = \mathbb{P}^r$. To see this we note that an embedding $X \hookrightarrow \mathbb{P}_k^r$ induces an embedding $X \times X \hookrightarrow \mathbb{P}^r \times \mathbb{P}^r$. Let $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ be the blowup of $\mathbb{P}^r \times \mathbb{P}^r$ along its diagonal $\Delta_{\mathbb{P}^r}$. The proper transform of $X \times X$ in $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ is $\widetilde{X \times X}$. Furthermore, the exceptional divisor E in $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ restricts to the exceptional divisor on $\widetilde{X \times X}$, and a v -adic metric on $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$ restricts to one on the corresponding line bundle on $\widetilde{X \times X}$. Thus the distance function $d'_v(\cdot, \cdot)$ on \mathbb{P}^r restricts to the distance function $d'_v(\cdot, \cdot)$ on X , and to show that $d'_v(\cdot, \cdot)$ and $d_v(\cdot, \cdot)$ are equivalent we may assume that $X = \mathbb{P}^r$.

Let $([Z_0 : \cdots : Z_r], [W_0 : \cdots : W_r])$ be coordinates on $\mathbb{P}^r \times \mathbb{P}^r$. The diagonal of $\mathbb{P}^r \times \mathbb{P}^r$ is cut out by the equations $Z_i W_j - Z_j W_i = 0$ for $0 \leq i < j \leq r$. Set $N = \binom{r+1}{2} - 1$, and let $U_{i,j}$, $0 \leq i < j \leq r$ be coordinates on \mathbb{P}^N (in some chosen order). The blowup $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ is the closure in $\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N$ of the graph of the rational map

$$\begin{array}{ccc} \mathbb{P}^r \times \mathbb{P}^r & \dashrightarrow & \mathbb{P}^N. \\ [Z_0 : \cdots : Z_r] \times [W_0 : \cdots : W_r] & \mapsto & [Z_0 W_1 - Z_1 W_0 : \cdots : Z_{r-1} W_r - Z_r W_{r-1}] \end{array}$$

Among the equations cutting out $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ in $\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N$ are

$$(Z_i W_j - Z_j W_i) U_{\ell m} = (Z_\ell W_m - Z_m W_\ell) U_{ij}$$

for all pairs (i, j) , (ℓ, m) , with $0 \leq i < j \leq r$, and $0 \leq \ell < m \leq r$. These equations simply express that the functions $Z_i W_j - Z_j W_i$ were used to give the rational map to \mathbb{P}^N . On an open subset of $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ where $U_{ij} \neq 0$ and $U_{\ell m} \neq 0$, we can rewrite this as the relation

$$\left. \frac{Z_i W_j - Z_j W_i}{U_{ij}} \right|_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}} = \left. \frac{Z_\ell W_m - Z_m W_\ell}{U_{\ell m}} \right|_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}.$$

Let E be the exceptional divisor of the blow up. The line bundle $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$ is the restriction of $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(1, 1, -1)$ to $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$. On the open set of $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ where $U_{ij} \neq 0$, restricting $(Z_i W_j - Z_j W_i)/U_{ij}$ gives a section of $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(1, 1, -1)|_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}} = \mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$. The equation above shows that these local sections patch together to give a global section s_E of $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$. This section has divisor E , and so we may use it to compute the distance. To give a v -adic metric on $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$, we put v -adic metrics on $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(1, 0, 0)$, $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(0, 1, 0)$, and $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(0, 0, 1)$, by putting an explicit v -adic metric on $\mathcal{O}_{\mathbb{P}^r}(1)$ and $\mathcal{O}_{\mathbb{P}^N}(1)$ as described below, and pull these back from the factors. These then give a v -adic metric on $\mathcal{O}_{\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N}(1, 1, -1)$ which we restrict to get a v -adic metric on $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(E)$. Here is our choice

of v -adic metric on $\mathcal{O}_{\mathbb{P}^r}(1)$ (and similarly for $\mathcal{O}_{\mathbb{P}^N}(1)$). Let V be the k -vector space underlying \mathbb{P}^r so that $\mathbb{P}^r = \mathbb{P}(V)$, and Z_0, \dots, Z_r the coordinates on \mathbb{P}^r (i.e., a chosen basis for V^*). Given any $\tilde{x} \in V$ and section $s \in H^0(\mathbb{P}^r, \mathcal{O}_{\mathbb{P}^r}(1)) = V^*$ we may evaluate $s(\tilde{x})$ to get an element of k_v . Given any $x \in \mathbb{P}^r$ we set

$$|s(x)|_v = \frac{\|s(\tilde{x})\|_v}{\max_{0 \leq i \leq r} (\|Z_i(\tilde{x})\|)}$$

where $\tilde{x} \in V$ is any representative of x . The formula above does not depend on the choice of representative \tilde{x} , and so is well defined; for this reason we will use the notation $s(x)$ and $Z_i(x)$ in further formulae. If two sections s and s' are equal at x , then $|s(x)|_v = |s'(x)|_v$, and hence the formula above puts a v -adic metric on the fibre at x . This metric varies continuously with $x \in X(k_v)$.

Let $x = [x_0 : \dots : x_r]$ and $y = [y_0 : \dots : y_r]$ be points of \mathbb{P}^r , with $x \neq y$, so that (x, y) is a point of $\mathbb{P}^r \times \mathbb{P}^r$ not on the diagonal. The corresponding point on $\widetilde{\mathbb{P}^r \times \mathbb{P}^r}$ (in the coordinates of the embedding in $\mathbb{P}^r \times \mathbb{P}^r \times \mathbb{P}^N$) is $([x_0 : \dots : x_r], [y_0 : \dots : y_r], [x_0 y_1 - x_1 y_0 : \dots : x_{r-1} y_r - x_r y_{r-1}])$. I.e., up to independent scalars in the Z, W , and U variables, $Z_i(x, y) = x_i$, $W_j(x, y) = y_j$, and $U_{ij}(x, y) = x_i y_j - x_j y_i$. Choosing any (ℓ, m) such that $U_{\ell m}(x, y) \neq 0$, so that s_E can be represented near (x, y) by $(Z_\ell W_m - Z_m W_\ell)/U_{\ell m}$ we compute that

$$\begin{aligned} d'_v(x, y) &= |s_E(x, y)|_v = \frac{\max_{0 \leq i < j \leq r} (\|U_{ij}(x, y)\|_v)}{\max_{0 \leq i \leq r} (\|Z_i(x, y)\|_v) \max_{0 \leq j \leq r} (\|W_j(x, y)\|_v)} \cdot \left\| \frac{(Z_\ell W_m - Z_m W_\ell)(x, y)}{U_{\ell m}(x, y)} \right\|_v \\ &= \frac{\max_{0 \leq i < j \leq r} (\|x_i y_j - x_j y_i\|_v)}{\max_{0 \leq i \leq r} (\|x_i\|_v) \max_{0 \leq j \leq r} (\|y_j\|_v)} \cdot \frac{\|x_\ell y_m - x_m y_\ell\|_v}{\|x_\ell y_m - x_m y_\ell\|_v} = \frac{\max_{0 \leq i < j \leq r} (\|x_i y_j - x_j y_i\|_v)}{\max_{0 \leq i \leq r} (\|x_i\|_v) \max_{0 \leq j \leq r} (\|y_j\|_v)}. \end{aligned}$$

Here on the first line the factors of $\max(\|Z_i(x, y)\|_v)$, $\max(\|W_j(x, y)\|_v)$, and $\max(\|U_{ij}(x, y)\|_v)$, come from the construction of the v -adic metric on $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(1, 0, 0)$, $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(0, 1, 0)$, and $\mathcal{O}_{\widetilde{\mathbb{P}^r \times \mathbb{P}^r}}(0, 0, -1)$, respectively, while the last factor is the evaluation of the local representation of s_E at (x, y) .

As the equation shows, for non-archimedean v we have $d_v(x, y) = d'_v(x, y)$. For archimedean v we first consider the case that $k_v = \mathbb{C}$. The issue is then to compare

$$d_v(x, y) = 1 - \frac{|\sum_{i=0}^r x_i \overline{y_i}|^2}{(\sum_{i=0}^r |x_i|^2)(\sum_{j=0}^r |y_j|^2)} = \frac{\sum_{0 \leq i < j \leq r} |x_i y_j - x_j y_i|^2}{(\sum_{0 \leq i \leq r} |x_i|^2)(\sum_{0 \leq j \leq r} |y_j|^2)},$$

with

$$d'_v(x, y) = \frac{\max_{0 \leq i < j \leq r} (|x_i y_j - x_j y_i|^2)}{\max_{0 \leq i \leq r} (|x_i|^2) \max_{0 \leq j \leq r} (|y_j|^2)}.$$

On \mathbb{C}^s with coordinate functions t_1, \dots, t_s , we have

$$\frac{1}{s} (|t_1|^2 + \dots + |t_s|^2) \leq \max(|t_1|^2, \dots, |t_s|^2) \leq |t_1|^2 + \dots + |t_s|^2,$$

so that $|t_1|^2 + \dots + |t_s|^2$ and $\max(|t_1|^2, \dots, |t_s|^2)$ are equivalent functions on \mathbb{C}^s . Applying this equivalence between the max and the sum to the factors in the numerator and denominator of $d_v(\cdot, \cdot)$ shows that $d_v(\cdot, \cdot)$ and $d'_v(\cdot, \cdot)$ are equivalent.

In the case that v is archimedean and $k_v = \mathbb{R}$, the comparison is between

$$d_v(x, y) = \left(\frac{\sum_{0 \leq i < j \leq r} |x_i y_j - x_j y_i|^2}{(\sum_{0 \leq i \leq r} |x_i|^2)(\sum_{0 \leq j \leq r} |y_j|^2)} \right)^{\frac{1}{2}} \quad \text{and} \quad d'_v(x, y) = \frac{\max_{0 \leq i < j \leq r} (|x_i y_j - x_j y_i|)}{\max_{0 \leq i \leq r} (|x_i|) \max_{0 \leq j \leq r} (|y_j|)}$$

which, after squaring the distance functions, reduces to the previous case. Thus in all cases $d_v(\cdot, \cdot)$ and $d'_v(\cdot, \cdot)$ are equivalent distance functions. \square

REFERENCES

- [ADHL] Arzhantsev, I; Derenthal, U.; Hausen, J.; Laface, A., *Cox Rings*, Cambridge studies in advanced mathematics 144, Cambridge University Press, New York, 2015.
- [BG] Bombieri, E.; Gubler, W., *Heights in Diophantine Geometry*, New Mathematical Monographs 4, Cambridge University Press, Cambridge, 2006.
- [CS] Cornell, G.; Silverman, J., *Arithmetic Geometry*, Springer-Verlag, New-York, 1986.
- [D] Demailly, J.-P., *Singular Hermitian metrics on positive line bundles*, Complex Algebraic Varieties (Bayreuth, 1990), Lect. Notes in Math., vol 1507, 1992, pp. 87–104.
- [HS] Hindry, M.; Silverman, J., *Diophantine geometry. An introduction*, Graduate Texts in Mathematics, 201. Springer-Verlag, New York, 2000.
- [La] Lang, S., *Diophantine Geometry*, Interscience Tracts in Pure and Applied Mathematics, No. 11, John Wiley & Sons, New York-London, 1962.
- [Laz] Lazarsfeld, R., *Positivity in Algebraic Geometry I*, Springer-Verlag, 2004.
- [L] Liouville, *Nouvelle démonstration d'un théorème sur les irrationnelles algébriques*, Comptes rendus hebdomadaires des séances de l'Académie des sciences, Tome XVIII, séance de 20 mai 1844, 910–911.
- [McK] McKinnon, D., *A conjecture on rational approximations to rational points*, J. Algebraic Geom., 16 (2007), 257–303.
- [MR] McKinnon, D. and Roth, M., *Seshadri constants, Diophantine approximation, and Roth's theorem for arbitrary varieties.*, Invent. Math. (200), 513–583.
- [Por] McKinnon, D., *Generators of subcones of the nef cone of a cubic surface*, tables computed with help of the package <http://www.iwr.uni-heidelberg.de/groups/comopt/software/PORTA/>. Tables available at <http://www.math.uwaterloo.ca/~dmckinno/Papers/cubictable.pdf>.
- [Se] Serre, J.P., *Lectures on the Mordell-Weil Theorem*, Vieweg, 1997.
- [V] Vojta, P., *Diophantine Approximations and Value Distribution Theory*, Lecture Notes in Mathematics vol. 1239, Springer-Verlag, 1987.

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ON N2L 3G1, CANADA

E-mail address: dmckinnon@uwaterloo.ca

DEPT. OF MATHEMATICS AND STATISTICS, QUEENS UNIVERSITY, KINGSTON, ONTARIO, CANADA

E-mail address: mikeroth@mast.queensu.ca