

## Representations of integers by certain positive definite binary quadratic forms

M. Ram Murty · Robert Osburn

Received: 5 February 2004 / Accepted: 14 February 2005 /

Published online: 18 September 2007

© Springer Science+Business Media, LLC 2007

**Abstract** We prove part of a conjecture of Borwein and Choi concerning an estimate on the square of the number of solutions to  $n = x^2 + Ny^2$  for a squarefree integer  $N$ .

**Keywords** Binary quadratic forms · Genus characters · Rankin-Selberg convolution of L functions · Modular forms

**Mathematics Subject Classification (2000)** Primary 11E25 · 11E45

### 1 Introduction

We consider the positive definite quadratic form  $Q(x, y) = x^2 + Ny^2$  for a squarefree integer  $N$ . Let  $r_{2,N}(n)$  denote the number of solutions to  $n = Q(x, y)$  (counting signs and order). In this note, we estimate

$$\sum_{n \leq x} r_{2,N}(n)^2.$$

A positive squarefree integer  $N$  is called solvable if  $x^2 + Ny^2$  has one form per genus. Note that this means the class number of the form class group of discriminant  $-4N$  equals the number of genera,  $2^t$ , where  $t$  is the number of distinct prime factors of  $N$ . Concerning  $r_{2,N}(n)$ , Borwein and Choi [2] proved the following:

---

M. Ram Murty (✉)

Department of Mathematics & Statistics, Queen's University, Kingston, ON, Canada K7L 3N6  
e-mail: murty@mast.queensu.ca

R. Osburn

Max Planck Institute für Mathematik, Vivatsgasse 7, 53111, Bonn, Germany  
e-mail: osburn@mpim-bonn.mpg.de

**Theorem 1.1** Let  $N$  be a solvable squarefree integer. Let  $x > 1$  and  $\epsilon > 0$ . We have

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{3}{N} \left( \prod_{p|2N} \frac{2p}{p+1} \right) (x \log x + \alpha(N)x) + O(N^{\frac{1}{4}+\epsilon} x^{\frac{3}{4}+\epsilon}),$$

where the product is over all primes dividing  $2N$  and

$$\alpha(N) = -1 + 2\gamma + \sum_{p|2N} \frac{\log p}{p+1} + \frac{2L'(1, \chi_{-4N})}{L(1, \chi_{-4N})} - \frac{12}{\pi^2} \zeta'(2),$$

where  $\gamma$  is the Euler-Mascheroni constant and  $L(1, \chi_{-4N})$  is the L-function corresponding to the quadratic character mod  $-4N$ .

Based on this result, Borwein and Choi posed the following:

**Conjecture 1.2** For any squarefree  $N$ ,

$$\sum_{n \leq x} r_{2,N}(n)^2 \sim \frac{3}{N} \left( \prod_{p|2N} \frac{2p}{p+1} \right) x \log x.$$

Our main result is the following.

**Theorem 1.3** Let  $Q(x, y) = x^2 + Ny^2$  for a squarefree integer  $N$  with  $-N \not\equiv 1 \pmod{4}$ . Let  $r_{2,N}(n)$  denote the number of solutions to  $n = Q(x, y)$  (counting signs and order). Then

$$\sum_{n \leq x} r_{2,N}(n)^2 \sim \frac{3}{N} \left( \prod_{p|2N} \frac{2p}{p+1} \right) x \log x.$$

## 2 Preliminaries

We first discuss two key estimates and a result of Kronecker on genus characters. Then using Kronecker's result, we prove a proposition relating genus characters to poles of the Rankin-Selberg convolution of L-functions. The first estimate is a recent result of Kühleitner and Nowak [13], namely

**Theorem 2.1** Let  $a(n)$  be an arithmetic function satisfying  $a(n) \ll n^\epsilon$  for every  $\epsilon > 0$ , with a Dirichlet series

$$F(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s} = \frac{(\zeta_K(s))^2}{(\zeta(2s))^{m_1} (\zeta_K(2s))^{m_2}} G(s),$$

where  $\Re(s) > 1$  and  $\zeta_K(s)$  is the Dedekind zeta function of some quadratic number field  $K$ ,  $G(s)$  is holomorphic and bounded in some half plane  $\Re(s) \geq \theta$ ,  $\theta < \frac{1}{2}$ , and

$m_1, m_2$  are nonnegative integers. Then for  $x$  large,

$$\begin{aligned} \sum_{n \leq x} a(n) &= \text{Res}_{s=1} \left( F(s) \frac{x^s}{s} \right) + O(x^{\frac{1}{2}} (\log x)^3 (\log \log x)^{m_1+m_2}) \\ &= Ax \log x + Bx + O(x^{\frac{1}{2}} (\log x)^3 (\log \log x)^{m_1+m_2}), \end{aligned}$$

where  $A$  and  $B$  are computable constants.

For an arbitrary quadratic number field  $K$  with discriminant  $d_K$ , let  $\mathcal{O}_K$  denote the ring of integers in  $K$ , and  $r_K(n)$  the number of integral ideals  $\mathcal{I}$  in  $\mathcal{O}_K$  of norm  $N(\mathcal{I}) = n$ . From (4.1) in [13], we have

$$\sum_{n=1}^{\infty} \frac{(r_K(n))^2}{n^s} = \frac{(\zeta_K(s))^2}{\zeta(2s)} \prod_{p|d_K} (1 + p^{-s})^{-1}.$$

Applying Theorem 2.1 with  $m_1 = 1$  and  $m_2 = 0$ , we obtain

**Corollary 2.2** For any quadratic field  $K$  of discriminant  $d_K$  and  $x$  large,

$$\sum_{n \leq x} (r_K(n))^2 = A_1 x \log x + B_1 x + O(x^{\frac{1}{2}} (\log x)^3 \log \log x),$$

with  $A_1 = \frac{6}{\pi^2} L(1, \chi_{d_K})^2 \prod_{p|d_K} \frac{p}{p+1}$  and  $B_1 = A_1 \alpha(N)$  with  $\alpha(N)$  as in Theorem 1.1.

The second estimate is a classical result of Rankin [17] and Selberg [18] which estimates the size of Fourier coefficients of a modular form. Specifically, if  $f(z) = \sum_{n=1}^{\infty} a(n)e^{2\pi i nz}$  is a nonzero cusp form of weight  $k$  on  $\Gamma_0(N)$ , then

$$\sum_{n \leq x} |a(n)|^2 = \alpha \langle f, f \rangle x^k + O(x^{k-\frac{2}{5}}),$$

where  $\alpha > 0$  is an absolute constant and  $\langle f, f \rangle$  is the Petersson scalar product. In particular, if  $f$  is a cusp form of weight 1, then  $\sum_{n \leq x} |a(n)|^2 = O(x)$ . One can adapt their result to say the following. Given two cusp forms of weight  $k$  on a suitable congruence subgroup of  $\Gamma = SL_2(\mathbb{Z})$ , say  $f(z) = \sum_{n=1}^{\infty} a(n)e^{2\pi i nz}$  and  $g(z) = \sum_{n=1}^{\infty} b(n)e^{2\pi i nz}$ , then

$$\sum_{n \leq x} a(n) \overline{b(n)} n^{1-k} = Ax + O(x^{\frac{3}{5}}),$$

where  $A$  is a constant. In particular, if  $f$  and  $g$  are cusp forms of weight 1, then  $\sum_{n \leq x} a(n) \overline{b(n)} = O(x)$ .

We will also use a result of Kronecker on genus characters. Let us first explain some terminology. Let  $K = \mathbb{Q}(\sqrt{d})$  be a quadratic field of discriminant  $d_K$ .  $d_K$  is said to be a prime discriminant if it only has one prime factor. Thus it must be of the form:  $-4, \pm 8, \pm p \equiv 1 \pmod{4}$  for an odd prime  $p$ . Every discriminant can be written

uniquely as a product of prime discriminants, say  $d_K = P_1 \dots P_k$ . Here  $k$  denotes the number of distinct prime factors of  $d_K$ . Thus  $d_K$  can be written as a product of two discriminants, say  $d_K = D_1 D_2$  in  $2^{k-1}$  distinct ways (excluding order). Now, for any such decomposition we define a character  $\chi_{D_1, D_2}$  on ideals by

$$\chi_{D_1, D_2}(\mathfrak{p}) = \begin{cases} \chi_{D_1}(N\mathfrak{p}) & \text{if } \mathfrak{p} \nmid D_1, \\ \chi_{D_2}(N\mathfrak{p}) & \text{if } \mathfrak{p} \nmid D_2, \end{cases}$$

where  $\chi_d(n)$ , is the Kronecker symbol. This is well defined on prime ideals because  $\chi_D(N\mathfrak{a}) = 1$  if  $(\mathfrak{a}, D) = 1$ .  $\chi_{D_1, D_2}$  extends to all fractional ideals by multiplicativity. Hence we have

$$\chi_{D_1, D_2} : I \rightarrow \{\pm 1\},$$

where  $I$  is the group of nonzero fractional ideals of  $\mathcal{O}_K$ . Thus  $\chi_{D_1, D_2}$  has order two, except for the trivial character corresponding to  $d_K = d_K \cdot 1 = 1 \cdot d_K$ . Every such character  $\chi_{D_1, D_2}$  is called the genus character of discriminant  $d_K$ . As these are different for distinct factorizations of  $d_K$  (into a product of two discriminants), we have  $2^{k-1}$  genus characters. Kronecker's theorem (see Theorem 12.7 in [11]) is as follows.

**Theorem 2.3** *The L-function of  $K$  associated with the genus character  $\chi_{D_1, D_2}$  factors into the Dirichlet L-functions,*

$$L(s, \chi_{D_1, D_2}) = L(s, \chi_{D_1})L(s, \chi_{D_2}).l$$

Let  $K = \mathbb{Q}(\sqrt{-N})$ ,  $N$  squarefree,  $I$  as above, and  $P$  the subgroup of  $I$  of principal ideals. For a non-zero integral ideal  $\mathfrak{m}$  of  $\mathcal{O}_K$ , define

$$I(\mathfrak{m}) = \{\mathfrak{a} \in I : (\mathfrak{a}, \mathfrak{m}) = 1\},$$

$$P(\mathfrak{m}) = \{\langle a \rangle \in P : a \equiv 1 \pmod{\mathfrak{m}}\}.$$

A group homomorphism  $\chi : I_{\mathfrak{m}} \rightarrow S^1$  is an ideal class character if it is trivial on  $P(\mathfrak{m})$ , i.e.

$$\chi(\langle a \rangle) = 1$$

for  $a \equiv 1 \pmod{\mathfrak{m}}$ . Thus an ideal class character is a character on the ray class group  $I(\mathfrak{m})/P(\mathfrak{m})$ . Taking the trivial modulus  $\mathfrak{m} = 1$ , we obtain a character on the ideal class group of  $K$ . Note that for  $K = \mathbb{Q}(\sqrt{-N})$ , a genus character is an ideal class character of order at most two.

Let us also recall the notion of the Rankin-Selberg convolution of two L-functions. For squarefree  $N$ , consider two ideal class characters  $\chi_1, \chi_2$  for  $\mathbb{Q}(\sqrt{-N})$ , and their associated Hecke L-series

$$L(s, \chi_1) = \sum_{n=1}^{\infty} \frac{\chi_1(n)}{n^s},$$

$$L(s, \chi_2) = \sum_{n=1}^{\infty} \frac{\chi_2(n)}{n^s}$$

which converge absolutely in some right half-plane. We form the convolution L-series by multiplying the coefficients,

$$L(s, \chi_1 \otimes \chi_2) = \sum_{n=1}^{\infty} \frac{\chi_1(n)\chi_2(n)}{n^s}.$$

The following result describes a relationship between genus characters  $\chi$  and the orders of poles of  $L(s, \chi \otimes \chi)$ .

**Proposition 2.4** *Let  $\chi$  be an ideal class character of  $\mathbb{Q}(\sqrt{-N})$ ,  $-N \not\equiv 1 \pmod{4}$ , and  $L(s, \chi)$ , the associated Hecke L-series. Then  $\chi$  is a genus character if and only if  $L(s, \chi \otimes \chi)$ , has a double pole at  $s = 1$ .*

*Proof* Suppose  $\chi_{D_1, D_2}$  is a genus character of discriminant  $-4N$ , and  $L(s, \chi_{D_1, D_2}) = \sum_{n=1}^{\infty} \frac{b_i(n)}{n^s}$ . By Theorem 2.3 and Exercise 1.2.8 in [14] (see the solution), we have

$$\sum_{n=1}^{\infty} \frac{b_i(n)^2}{n^s} = \frac{L(s, \chi_{D_1}^2)L(s, \chi_{D_2}^2)L(s, \chi_{D_1}\chi_{D_2})^2}{L(2s, \chi_{D_1}^2\chi_{D_2}^2)}.$$

Note that

$$L(s, \chi_{D_1}^2) = \zeta(s) \cdot \prod_{p|D_1} (1 - p^{-s}),$$

$$L(s, \chi_{D_2}^2) = \zeta(s) \cdot \prod_{p|D_2} (1 - p^{-s}),$$

$$L(s, \chi_{D_1}\chi_{D_2})^2 = L(s, \chi_{-4N})^2,$$

and

$$L(2s, \chi_{D_1}^2\chi_{D_2}^2) = \zeta(2s) \cdot \prod_{p|D_1D_2} (1 - p^{-2s}).$$

We have

$$\sum_{n=1}^{\infty} \frac{b_i(n)^2}{n^s} = \frac{\zeta(s)^2 L(s, \chi_{-4N})^2}{\zeta(2s)} \prod_{p|2N} (1 + p^{-s})^{-1}$$

and thus a double pole at  $s = 1$ .

Conversely, let  $\chi$  be an ideal class character of  $K = \mathbb{Q}(\sqrt{-N})$ , and suppose  $L(s, \chi \otimes \chi)$  has a double pole at  $s = 1$ . Now  $\chi$  is an automorphic form on  $GL_1(\mathbb{A}_K)$ . By automorphic induction (see [1]),  $\chi$  is mapped to  $\pi$ , a cuspidal automorphic representation of  $GL_2(\mathbb{A}_{\mathbb{Q}})$ . Note that  $\pi$  is reducible as, otherwise,  $L(s, \pi \otimes \pi)$ , has a simple pole at  $s = 1$  ([1], p. 200). As  $K$  is a quadratic extension of  $\mathbb{Q}$ , we must have  $\pi = \chi_1 + \chi_2$  where  $\chi_i$  are Dirichlet characters. As  $L(s, \chi) = L(s, \pi)$  (see [1]) and thus  $L(s, \chi \otimes \chi) = L(s, \pi \otimes \pi)$ ,

$$L(s, \pi \otimes \pi) = L(s, \chi \otimes \chi) = \frac{L(s, \chi_1^2)L(s, \chi_2^2)L(s, \chi_1\chi_2)^2}{L(2s, \chi_1^2\chi_2^2)}.$$

Now  $L(s, \chi \otimes \chi)$ , has a double pole at  $s = 1$  if and only if either  $\chi_1 = \overline{\chi_2}$ ,  $\chi_2^2 \neq 1$ , and  $\chi_1^2 \neq 1$  or  $\chi_1^2 = 1$ ,  $\chi_2^2 = 1$ , and  $\chi_1 \chi_2 \neq 1$ . The latter implies  $\chi$  is a genus character. We now need to show that the former also implies that  $\chi$  is a genus character. Note that

$$L(s, \chi) = \prod_{\mathfrak{p}} \left(1 - \frac{\chi(\mathfrak{p})}{N(\mathfrak{p})^s}\right)^{-1}$$

and

$$L(s, \chi_1 + \chi_2) = \prod_p \left(1 - \frac{\chi_1(p)}{p^s}\right)^{-1} \prod_p \left(1 - \frac{\chi_2(p)}{p^s}\right)^{-1}.$$

As  $L(s, \chi) = L(s, \pi)$ , and  $L(s, \pi) = L(s, \chi_1 + \chi_2)$ , we compare Euler factors to get

$$\chi_1(p) + \chi_2(p) = \begin{cases} 0 & \text{if } p \text{ is inert in } K, \\ \chi(\mathfrak{p}) + \overline{\chi(\mathfrak{p})} & \text{if } p \text{ splits in } K. \end{cases}$$

For  $p$  inert in  $K$ , this yields  $\chi_1(p) = -\chi_2(p)$ , and so  $\overline{\chi_2(p)} = \chi_1(p) = -\chi_2(p)$ , which implies  $\chi_2^2(p) = -1$  and so  $\chi_2(p) = \pm i$ . Now consider the following equation whose sum sieves the inert primes

$$\frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \left(1 - \left(\frac{-4N}{p}\right)\right) \chi_2^2(p) = -\pi(x).$$

Here  $\pi(x)$  is the number of primes between 1 and  $x$ . Thus

$$\frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \chi_2^2(p) - \frac{1}{2} \sum_{\substack{p \leq x \\ p \text{ prime}}} \left(\frac{-4N}{p}\right) \chi_2^2(p) = -\pi(x).$$

As  $\chi_2^2 \neq 1$ , we have by the prime ideal theorem,  $\sum_{p \leq x} \chi_2^2(p) = o(\pi(x))$ , and so

$$\sum_{p \leq x} \left(\frac{-4N}{p}\right) \chi_2^2(p) \sim \pi(x).$$

This implies  $\left(\frac{-4N}{p}\right) \chi_2^2(p) = 1$ . If  $p$  splits in  $K$ , then  $\chi_2^2(p) = 1$  and so  $\chi_2(p) = \pm 1$ . A similar argument works for  $\chi_1$  and so we also have  $\chi_1(p) = \pm 1$  if  $p$  splits in  $K$ .

Again comparing the Euler factors in  $L(s, \chi)$ , and  $L(s, \pi)$ , the values of  $\chi(\mathfrak{p})$  must coincide with the values of  $\chi_1(p)$ , and  $\chi_2(p)$ , that is,  $\chi(\mathfrak{p}) = \pm 1$ . Now  $\chi(\mathfrak{p}) = \chi([\mathfrak{p}])$ , where  $[\mathfrak{p}]$  is the class of  $\mathfrak{p}$  in the ideal class group of  $K$ . By the analog of Dirichlet's theorem for ideal class characters, we know that in each ideal class  $\mathfrak{C}$  there are infinitely many prime ideals which split. Thus  $\chi(\mathfrak{C}) = \pm 1$  and hence is of order 2. This implies  $\chi$  is a genus character.  $\square$

*Remark 2.5* By Proposition 2.4, if  $\chi$  is a non-genus character, then  $L(s, \chi \otimes \chi)$ , has at most a simple pole at  $s = 1$ .

### 3 Proof of Theorem 1.3

As  $-N \not\equiv 1 \pmod{4}$ , the discriminant of  $K = \mathbb{Q}(\sqrt{-N})$ , is  $-4N$ . We also assume that  $t$  is the number of distinct prime factors of  $N$  and so the discriminant  $-4N$  has  $t + 1$  distinct prime factors.

Given the quadratic form  $Q(x, y) = x^2 + Ny^2$ , we consider the associated Epstein zeta function (see [7, 12, 19], or [20])

$$\zeta_Q(s) = \sum_{x, y \neq 0} \frac{1}{(x^2 + Ny^2)^s} = \sum_{n=1}^{\infty} \frac{r_{2, N}(n)}{n^s}$$

for  $\Re(s) > 1$ . Now for  $K = \mathbb{Q}(\sqrt{-N})$ , we have Dedekind's zeta function

$$\zeta_K(s) = \sum_{\mathfrak{a}} \frac{1}{N(\mathfrak{a})^s} = \sum_{n=1}^{\infty} \frac{a_n}{n^s},$$

where the sum is over all nonzero ideals  $\mathfrak{a}$  of  $\mathcal{O}_K$ . We now split up  $\zeta_K(s)$ , according to the classes  $c_i$  of the ideal class group  $C(K)$ , into the partial zeta functions (see page 458 of [15])

$$\zeta_{c_i}(s) = \sum_{\mathfrak{a} \in c_i} \frac{1}{N(\mathfrak{a})^s}$$

so that  $\zeta_K(s) = \sum_{i=0}^{h-1} \zeta_{c_i}(s)$ , where  $h$  is the class number of  $K$ . In our case  $K = \mathbb{Q}(\sqrt{-N})$  is an imaginary quadratic field and so by [6] (Theorem 7.7, page 137), we may write

$$\zeta_K(s) = \sum_{i=0}^{h-1} \zeta_{Q_i}(s),$$

where  $Q_i$  is a class in the form class group. Note that in this context,  $Q(x, y)$  corresponds to the trivial class  $c_0$  in  $C(K)$ , and so  $\zeta_{c_0}(s) = \zeta_{Q(x, y)}(s)$ . Now let  $\chi$  be an ideal class character and consider the Hecke L-function for  $\chi$ , namely

$$L(s, \chi) = \sum_{\mathfrak{a}} \frac{\chi(\mathfrak{a})}{N(\mathfrak{a})^s},$$

where  $\mathfrak{a}$  again runs over all nonzero ideals of  $\mathcal{O}_K$ . We may now rewrite the Hecke L-function as

$$L(s, \chi) = \sum_{i=0}^{h-1} \chi(c_i) \zeta_{c_i}(s).$$

And so summing over all ideal class characters of  $C(K)$ , we have

$$\sum_{\chi} \overline{\chi}(c_0) L(s, \chi) = \sum_{i=0}^{h-1} \zeta_{c_i}(s) \left( \sum_{\chi} \overline{\chi}(c_0) \chi(c_i) \right).$$

The inner sum is nonzero precisely when  $i = 0$ . As  $\bar{\chi}(c_0) = 1$  we have  $\zeta_{c_0}(s) = \frac{1}{h} \sum_{\chi} L(s, \chi)$ . Thus

$$\zeta_{c_0}(s) = \frac{1}{h} (L(s, \chi_0) + L(s, \chi_1) + \cdots + L(s, \chi_{h-1})).$$

As  $\chi_0$  is the trivial character,  $L(s, \chi_0) = \zeta_K(s)$ . We now compare  $n$ th coefficients, yielding

$$r_{2,N}(n) = \frac{1}{h} (a_n + b_1(n) + \cdots + b_{h-1}(n)),$$

where  $a_n$  is the number of integral ideals of  $\mathcal{O}_K$  of norm  $n$  and the  $b_i$ 's are coefficients of weight 1 cusp forms (see the classical work of Hecke [9, 10] or [3]). From the modern perspective, this is straightforward. Each  $L(s, \chi_i)$ ,  $1 \leq i \leq h-1$ , can be viewed as an automorphic L-function of  $GL_1(\mathbb{A}_K)$ , and by automorphic induction (see [1]) they are essentially Mellin transforms of (holomorphic) cusp forms, in the classical sense. We now have

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{1}{h^2} \left( \sum_{n \leq x} a_n^2 + \sum_{\substack{i \\ n \leq x}} b_i(n)^2 + 2 \sum_{\substack{i \\ n \leq x}} a_n b_i(n) + \sum_{\substack{i \neq j \\ n \leq x}} b_i(n) b_j(n) \right).$$

By the Rankin-Selberg estimate,  $2 \sum_{n \leq x} a_n b_i(n)$ ,  $\sum_{n \leq x} b_i(n) b_j(n)$ , are equal to  $O(x)$ . By Corollary 2.2,

$$\frac{1}{h^2} \sum_{n \leq x} a_n^2 = \frac{1}{h^2} (A_1 x \log x + B_1 x + O(x^{\frac{1}{2}} (\log x)^3 \log \log x)).$$

We now must estimate  $\sum_{n \leq x} b_i(n)^2$ . Let us now assume that the first  $2^t - 1$  terms arise from L-functions associated to genus characters. By Proposition 2.4 and Nowak's proof of Theorem 2.1 (which uses Perron's formula and the residue theorem), we obtain

$$\sum_{n \leq x} b_i(n)^2 = A_1 x \log x + B_1 x + O(x),$$

with  $A_1$  and  $B_1$  as in Corollary 2.2. As this estimate holds for each  $i$  such that  $1 \leq i \leq 2^t - 1$ , the term  $A_1 x \log x$  appears  $2^t$  times in the estimate of  $\sum_{n \leq x} r_{2,N}(n)^2$ . By Remark 2.5, the remaining terms  $\sum_{n \leq x} b_i(n)^2$  for  $2^t - 1 < i \leq h-1$  are all  $O(x)$ . Thus

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{1}{h^2} \left[ \left( 2^t \frac{6}{\pi^2} L(1, \chi_{-4N})^2 \prod_{p|2N} \frac{p}{p+1} \right) x \log x + O(x) \right] + O(x).$$

By (4.11) in [8] (or (8), page 171 in [5]), we have  $L(1, \chi_{-4N}) = \frac{h\pi}{\sqrt{N}}$  and so

$$\sum_{n \leq x} r_{2,N}(n)^2 = \frac{3}{N} \left( \prod_{p|2N} \frac{2p}{p+1} \right) x \log x + O(x).$$

The result then follows.

*Remark 3.1* It should be possible to obtain the second term in the asymptotic formula. By a careful application of the Rankin-Selberg method, one should obtain an error term of the form  $O(x^\theta)$ , with  $\theta < 1$ . The remaining case  $-N \equiv 1 \pmod{4}$  requires more subtle analysis due to the fact that for  $K = \mathbb{Q}(\sqrt{-N})$ ,  $\mathbb{Z}[\sqrt{-N}]$  is not the maximal order of  $K$ . One approach involves the study of L-series attached to orders. Using the techniques in [4] and [12], it should be possible to do this as well as derive sharper error terms. Another way is to use generalized ideal class groups and Hecke L-series attached to them and relate these back to the order. This has now been done by the second author in [16]. Thus the full conjecture of Borwein-Choi is now settled.

## References

1. Arthur, J., Clozel, L.: Simple Algebras, Base Change, and the Advanced Theory of the Trace Formula. Annals of Math. Studies, vol. 120. Princeton University Press, Princeton (1990)
2. Borwein, J., Choi, K.K.: On Dirichlet Series for sums of squares. Ramanujan J., special issues for Robert Rankin **7**, 1–3, 97–130
3. Bump, D.: Automorphic Forms and Representations. Cambridge Studies in Advanced Mathematics, vol. 55. Cambridge University Press (1997)
4. Chapman, R., van der Poorten, A.: Binary quadratic forms and the eta function. In: Number Theory for the Millennium, I (Urbana, IL), pp. 215–227. A.K. Peters, Natick (2002)
5. Cohn, H.: Advanced Number Theory. Dover, New York (1980)
6. Cox, D.: Primes of the Form  $x^2 + ny^2$ . Wiley Inc., New York (1989)
7. Epstein, P.: Zur Theorie allgemeiner Zetafunction. Math. Ann. **56**, 615–644 (1903)
8. Grosswald, E.: Representations of Integers as Sums of Squares. Springer, Berlin (1985)
9. Hecke, E.: Zur theorie der elliptischen modulfunktionen. Math. Ann. **97**, 210–242 (1926)
10. Hecke, E.: Über das verhalten von  $\sum_{m,n} e^{2\pi i t \frac{|m^2 - 2n^2|}{8}}$  und ähnlichen funktionen bei modulustitutionen. J. Reine Angew. Math. **157**, 159–170 (1927)
11. Iwaniec, H.: Topics in Classical Automorphic Forms. Graduate Studies in Mathematics, vol. 17. Am. Math. Soc., Providence (1997)
12. Kani, E.: Epstein Zeta-functions and Hecke L-functions, Preprint
13. Kühleitner, M., Nowak, W.G.: The average number of solutions to the Diophantine equation  $U^2 + V^2 = W^3$  and related arithmetic functions. Acta Math. Hung. **104**, 225–240 (2004)
14. Murty, R.: Problems in Analytic Number Theory. Graduate Texts in Mathematics, vol. 206. Springer, Berlin (2001)
15. Neukirch, J.: Algebraic Number Theory. Grundlehren, vol. 322. Springer, New York (1999)
16. Osburn, R.: A remark on a conjecture of Borwein and Choi. Proc. Am. Math. Soc. **133**, 2903–2909 (2005)
17. Rankin, R.A.: Contributions to the theory of Ramanujan's function  $\tau(n)$  and similar functions. II. The order of the Fourier coefficients of integral modular forms. Proc. Camb. Philos. Soc. **35**, 357–373 (1939)
18. Selberg, A.: Bemerkungen über eine Dirichletsche Reihe, die mit der Theorie der Modulformen nahe verbunden ist. Arch. Math. Natur. B **43**, 47–50 (1940)
19. Selberg, A., Chowla, S.: On Epstein's zeta function. J. Reine Angew. Math. **227**, 86–110 (1967)
20. Siegel, C.L.: Advanced Analytic Number Theory. Tata Institute of Fundamental Research, Bombay (1980)