

Time-varying and state-dependent recovery rates in epidemiological models



Scott Greenhalgh^{a, b, *}, Troy Day^b

^a Department of Mathematics, Siena College, Loudonville, NY, 12211, USA

^b Department of Mathematics and Statistics, Queen's University, Jeffery Hall, Kingston, ON, K7L 3N6, Canada

ARTICLE INFO

Article history:

Received 29 March 2017

Received in revised form 23 September 2017

Accepted 26 September 2017

Available online 14 October 2017

Keywords:

Differential equations

Integral equations

Infectious disease modeling

Waiting time distribution

Duration of infectiousness

SIR epidemic model

ABSTRACT

Differential equation models of infectious disease have undergone many theoretical extensions that are invaluable for the evaluation of disease spread. For instance, while one traditionally uses a bilinear term to describe the incidence rate of infection, physically more realistic generalizations exist to account for effects such as the saturation of infection. However, such theoretical extensions of recovery rates in differential equation models have only started to be developed. This is despite the fact that a constant rate often does not provide a good description of the dynamics of recovery and that the recovery rate is arguably as important as the incidence rate in governing the dynamics of a system. We provide a first-principles derivation of state-dependent and time-varying recovery rates in differential equation models of infectious disease. Through this derivation, we demonstrate how to obtain time-varying and state-dependent recovery rates based on the family of Pearson distributions and a power-law distribution, respectively. For recovery rates based on the family of Pearson distributions, we show that uncertainty in skewness, in comparison to other statistical moments, is at least two times more impactful on the sensitivity of predicting an epidemic's peak. In addition, using recovery rates based on a power-law distribution, we provide a procedure to obtain state-dependent recovery rates. For such state-dependent rates, we derive a natural connection between recovery rate parameters with the mean and standard deviation of a power-law distribution, illustrating the impact that standard deviation has on the shape of an epidemic wave.

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of KeAi Communications Co., Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Compartmental models of infectious-disease transmission have proven to be an invaluable tool for the prediction of disease progression and the evaluation of public health policies and interventions. In general, compartmental models describe how an infectious disease propagates throughout a population by characterizing the rates of transition of a populace from states of susceptible to, infected with, and finally recovered from disease. The rate of transition from the susceptible state to the infected state is often called the force of infection or incidence rate (Liu, Hethcote, & Levin, 1987), and the rate of transition from the state of infected with the disease to recovered from disease is called the recovery rate. While classically the

* Corresponding author. Department of Mathematics, Siena College, 515 Loudon Road, Loudonville, NY, 12211, USA.

E-mail address: scottyg3@gmail.com (S. Greenhalgh).

Peer review under responsibility of KeAi Communications Co., Ltd.

incidence rate is taken as a bilinear term, there has been extensive theoretical work regarding alternative nonlinear incidence rates. These alternative nonlinear incidence rates are normally developed to account for behavioral characteristics, such as the crowding of infected individuals and the avoidance of exposure to infection (Alexander & Moghadas, 2005; Liu et al., 1987; Liu, Levin, & Iwasa, 1986; Ruan & Wang, 2003; van den Driessche & Watmough, 2000) or to account for multi-stage infections (Krylova & Earn, 2013). In fact, formulations of nonlinear incidence rates can be justified through a first-principles derivation (Ponciano & Capistrán, 2011).

Like incidence rates, recovery rates have received significant attention in regards to their formulations in integral equations (Feng, Xu, & Zhao, 2007; Fowler & Hollingsworth, 2015; Hethcote & Tudor, 1980) and stochastic epidemic models (Ball, 1983, 1986, 1991; Britton, 2010; Clancy, 1999, 2014). Recovery rates have also been generalized in ordinary differential equation (ODE) models, mainly through the method of stages, whereby the infected state is broken into multiple stages, where each stage has a constant recovery rate (Krylova & Earn, 2013; Lloyd, 2001a, 2001b). However, theoretical extensions of recovery rates beyond multiple constants in ODEs have only started to be developed, despite the fact that constant recovery rates based on the mean value of an exponentially distributed infectious period are epidemiologically unrealistic (Bailey et al., 1975; Gough, 1977; Keeling & Grenfell, 1999; Keeling & Rohani, 2007; Lloyd, 2001a, 2001b).

Here we derive time-varying and state-dependent recovery rates for ODE models of infectious diseases. To obtain such novel recovery rates, we use the connection between integral equation models, ODEs and survival functions. First we show how an integral equation model for an infectious disease (Hethcote & Tudor, 1980) is related to more typical ODEs. A key component of this relationship is the probability distribution of the time spent in the infectious class. Most traditional models assume this time is exponentially distributed, and so we review this distribution and its connection to stochastic processes. Next, we show how an alternative probability distribution of the time spent in the infectious class leads to different forms of recovery rates in ODE models.

We validate our work by modeling historical measles outbreaks in Iceland (Cliff, Haggett, Ord, & Versey, 1981). First, we develop a compartmental model with a time-varying recovery rate based on infectious periods that follow the family of Pearson distributions (Pearson, 1893, 1895). The family of Pearson distributions span all possible values of skewness and kurtosis, so we evaluate the sensitivity of model predictions of the timing and magnitude of an epidemic peak with such time-varying recovery rates, relative to the uncertainty in the skewness and kurtosis derived from infectious period data.

Finally, we turn to state-dependent recovery rates, where we compare recovery rates based on power-law distributions to constant recovery rate models. Through this comparison, we demonstrate advantages of recovery rates based on a power-law distribution, including the potential for scaling invariance, the finite contribution of infected individuals to disease spread and the potential for deterministic disease burnout.

2. Methods

We now outline how to obtain state-dependent and time-varying recovery rates η in ODE models. To accomplish this, we illustrate the connection between integral equation models to survival functions and finally ODE models.

2.1. Integral equation models of infectious diseases

We considered a population divided into the proportion of susceptible (s), infected (i) and recovered (r) individuals, with $s + i + r = 1$, which follow an integral equation formulation of an SIR compartmental model:

$$\begin{aligned} s &= 1 - (1 - s_0)\exp(-\tilde{b}t) - \int_{t_0}^t \tilde{\lambda}(x)s(x)\exp(-\tilde{b}(t-x))dx, \\ i &= i_0P(t, t_0)\exp(-\tilde{b}t) + \int_{t_0}^t \tilde{\lambda}(x)s(x)P(t, x)\exp(-\tilde{b}(t-x))dx, \\ r &= (r_0 + i_0 - i_0P(t, t_0))\exp(-\tilde{b}t) + \int_{t_0}^t \tilde{\lambda}(x)s(x)(1 - P(t, x))\exp(-\tilde{b}(t-x))dx. \end{aligned} \tag{1}$$

Here $\tilde{\lambda}$ is the force of infection, \tilde{b} is the population birth rate minus the population death rate, and $P(t, x)$ is the waiting-time distribution for the time spent in the infectious class before recovery, where x is the time of infection.

As $P(t, x)$ characterizes the time waiting in the infectious class before recovery, it also may be interpreted as a survival function. Thereby, $P(t, x)$ possesses all the properties of a survival function, including $1 - P(t, x)$ corresponding to cumulative distribution function, $-\frac{d}{dt} \ln P(t, x)$ being a hazard function, and $\int_x^\infty P(t, x)dt$ as the average waiting-time (Arino & van den Driessche, 2006; Rodriguez, 2007).

Note, as $r = 1 - s - i$, we omit entirely the integral equation, integro-differential equation (IDE) and ODE formulations of r for the ease of presentation.

2.2. The probability of remaining infected

To generalize constant recovery rates, we now provide insight into the formulation of $P(t, t_0)$. We begin by expressing $P(t, t_0)$ in terms of the recovery rate. Taking $\phi(t, t + \Delta t)$ to be the probability of recovery during the time interval $[t, t + \Delta t]$, it follows that

$$\phi(t, t + \Delta t) = \eta(i(t), s(t), t)\Delta t,$$

where $\eta(i(t), s(t), t)$ is a time-dependent and/or state-dependent recovery rate. Thus, if we want to know $P(t + \Delta t, t_0)$, where Δt is some small increment of time, then

$$P(t + \Delta t, t_0) \approx P(t, t_0)(1 - \phi(t, t + \Delta t)), \quad (2)$$

If $\eta(i(t), s(t), t)$ is constant and time invariant, then $\phi(t, t + \Delta t) = \gamma\Delta t$, and

$$P(t + \Delta t, t_0) \approx P(t, t_0)(1 - \gamma\Delta t), \quad (3)$$

where γ is the constant recovery rate.

By rearranging the terms in (3) and letting $\Delta t \rightarrow 0$, the ODE formulation of $P(t, t_0)$ is obtained:

$$\frac{d}{dt}P(t, t_0) = -\gamma P(t, t_0), \quad P(t_0, t_0) = 1$$

and thus,

$$P(t, t_0) = \exp(-\gamma(t - t_0)). \quad (4)$$

Alternatively, if $\phi(t, t + \Delta t)$ is dependent on factors, such as the state of the epidemic, or time, then $\phi(t, t + \Delta t) = \eta(i, s, t)\Delta t$ where $\eta(i, s, t)$ is the recovery rate at time t . It once again follows from (2) that

$$P(t + \Delta t, t_0) \approx P(t, t_0)(1 - \eta(i, s, t)\Delta t).$$

Rearranging terms and letting $\Delta t \rightarrow 0$ yields the ODE formulation of $P(t, t_0)$:

$$\frac{d}{dt}P(t, t_0) = -\eta(i, s, t)P(t, t_0), \quad P(t_0, t_0) = 1 \quad (5)$$

and thus,

$$P(t, t_0) = \exp\left(-\int_{t_0}^t \eta(i(z), s(z), z)dz\right). \quad (6)$$

Equation (6) is not the only alternative to the waiting-time distribution (4). For instance, taking

$$P(t, t_0) = \begin{cases} 1 & \text{if } t \in [t_0, \omega + t_0] \\ 0 & \text{otherwise} \end{cases}$$

where ω is the constant waiting-time in the infected state, is a common assumption used to motivate delay-differential equations (Arino & van den Driessche, 2006).

2.3. Differential equation models of infectious diseases

To obtain ODE models of infectious diseases, we start by differentiating (1) to obtain the IDEs:

$$\begin{aligned} \frac{ds}{dt} &= \tilde{b}(1 - s_0)\exp(-\tilde{b}(t - t_0)) + \tilde{b} \int_{t_0}^t \tilde{\lambda}(x)s(x)\exp(-\tilde{b}(t - x))dx - \tilde{\lambda}s, \\ \frac{di}{dt} &= \frac{d}{dt} \left(i_0 P(t, t_0)\exp(-\tilde{b}(t - t_0)) \right) + \int_{t_0}^t \tilde{\lambda}(x)s(x) \frac{d}{dt} \left(P(t, x)\exp(-\tilde{b}(t - x)) \right) dx + \tilde{\lambda}s. \end{aligned} \tag{7}$$

From system (7), we obtain ODEs by imposing assumptions on $P(t, x)$, and then using system (1) to eliminate the integral terms.

2.3.1. Constant recovery rates

To derive ODE models with a constant recovery rate from integral equation models, we make use of the standard assumptions that $P(t, t_0) = \exp(-\gamma(t - t_0))$. Thus it follows that $\frac{d}{dt}P(t, t_0) = -\gamma \exp(-\gamma(t - t_0)) = -\gamma P(t, t_0)$. Thereby system (7) simplifies to

$$\begin{aligned} \frac{ds}{dt} &= \tilde{b}(1 - s_0)\exp(-\tilde{b}(t - t_0)) + \tilde{b} \int_{t_0}^t \tilde{\lambda}(x)s(x)\exp(-b(t - x))dx - \tilde{\lambda}s, \\ \frac{di}{dt} &= -(\gamma + \tilde{b}) \left(i_0 P(t, t_0)\exp(-\tilde{b}(t - t_0)) + \int_{t_0}^t \tilde{\lambda}(x)s(x)P(t, x)\exp(-\tilde{b}(t - x))dx \right) + \tilde{\lambda}s. \end{aligned} \tag{8}$$

From system (8), the standard ODE formulation of an SIR compartmental model is obtained by further substituting system (1), which yields:

$$\begin{aligned} \frac{ds}{dt} &= \tilde{b} - \tilde{b}s - \tilde{\lambda}s, \\ \frac{di}{dt} &= \tilde{\lambda}s - \gamma i - \tilde{b}i, \end{aligned} \tag{9}$$

2.3.2. State-dependent and time-varying recovery rates

Alternative assumptions on $P(t, t_0)$ leads to formulation of other recovery rates. For instance, $P(t, t_0)$ as in (5). To obtain state-dependent and time-varying recovery rates, first note that (6) reduces the IDEs from system (7) to

$$\begin{aligned} \frac{ds}{dt} &= \tilde{b} - \tilde{b}s - \tilde{\lambda}s, \\ \frac{di}{dt} &= -(\eta(i, s, t) + \tilde{b}) \left(i_0 \exp\left(-\int_{t_0}^t \eta(i(z), s(z), z)dz - \tilde{b}(t - t_0)\right) + \int_{t_0}^t \tilde{\lambda}(x)s(x)\exp\left(-\int_x^t \eta(i(z), s(z), z)dz - \tilde{b}(t - x)\right)dx \right) + \tilde{\lambda}s. \end{aligned} \tag{10}$$

The IDEs from system (10), through the substitution of system (1), reduce to the ODEs,

$$\begin{aligned} \frac{ds}{dt} &= \tilde{b} - \tilde{b}s - \tilde{\lambda}s, \\ \frac{di}{dt} &= \tilde{\lambda}s - \eta(i, s, t)i - \tilde{b}i, \end{aligned} \tag{11}$$

with recovery rate $\eta(i, s, t)$.

2.4. The reproductive numbers

The next-generation method is a standard technique to determine the basic reproduction number, R_0 (Diekmann, Heesterbeek, & Metz, 1990; Heffernan, Smith, & Wahl, 2005; Liu et al., 1987; van den Driessche & Watmough, 2000, 2002). Traditionally, the next-generation method is applied to system (10), with $\tilde{\lambda} = \beta i$ and $\eta(i, s, t) = \gamma$, linearizing the flow of newly infected individuals, $\mathcal{F} = \beta is$, and the flow from the infected state $\mathcal{V} = \gamma i + bi$, at the disease-free equilibrium $E_0 = \{s = 1, i = 0, r = 0\}$. From the linearized flows,

$$F = \left. \frac{d}{di} \mathcal{F} \right|_{E_0} = \left. \frac{d}{di} \tilde{\lambda} \right|_{E_0} = \beta \text{ and } V = \left. \frac{d}{di} \mathcal{V} \right|_{E_0} = \left. \frac{d}{di} (\gamma i + bi) \right|_{E_0} = \gamma + b.$$

The largest eigenvalue of the next-generation matrix, $G = FV^{-1}$ is taken as R_0 , which in this case corresponds to $\frac{\beta}{\gamma+b}$.

In general, system (9) with arbitrary $\tilde{\lambda}$ and η need not be linearizable (Greenhalgh, Galvani, & Medlock, 2015), which impedes the use of the next-generation method. Thus, we consider computing the effective reproduction number directly as

$$Re(i, s) = \mathcal{F}\mathcal{V}^{-1} = \frac{\tilde{\lambda}s}{\eta i + bi},$$

and the basic reproduction number as,

$$R_0 = Re(i, s)|_{s=1}.$$

If the assumption of a constant R_0 is still imposed, it follows that the force of infection is of the form:

$$\tilde{\lambda} = R_0(\eta|_{s=1} + b)i. \tag{12}$$

For the force of infection (12) to be biologically meaningful (Liu et al., 1987), it follows that

$$\lim_{i \rightarrow 0^+} \tilde{\lambda} = 0 \Leftrightarrow \lim_{i \rightarrow 0^+} i \cdot \eta|_{s=1} = 0.$$

Such a condition ensures the infectious period is well-posed near the disease-free equilibrium and provides a natural connection to the null probability of transmission in the absence of infection (Korobeinikov & Maini, 2005; Ponciano & Capistrán, 2011). It also follows that

$$\frac{d\tilde{\lambda}}{di} \geq 0 \Leftrightarrow i \left. \frac{d\eta}{di} \right|_{s=1} + \eta|_{s=1} + b \geq 0,$$

and

Table 1
Parameter values.

Definition	Parameter	Point estimate	Reference
Transmission rate (time-varying)	β_{TV}	0.2374/day	Fit
Rate of demographic turnover	b	0.027/year	Statistics Iceland
Average infectious period	μ	8.00 days	(Cliff et al., 1981; Simpson, 1952)
Standard deviation of infectious period	σ	2.26 days	(Cliff et al., 1981; Simpson, 1952)
Skewness (μ_3) and kurtosis (μ_4) of infectious period distribution: Estimates from data	μ_3	0.39 days	(Cliff et al., 1981; Simpson, 1952)
	μ_4	2.92 days	
Best fits of Pearson distributions			
Type I	μ_3	0.23 days	
	μ_4	2.61 days	
Type III	μ_3	02.1 days	
	μ_4	09.6 days	
Type IV	μ_3	00.1 days	
	μ_4	3.03 days	
Type V	μ_3	0.66 days	
	μ_4	3.82 days	
Type VI	μ_3	4.13 days	
	μ_4	46.5 days	

$$\frac{d^2 \tilde{\lambda}}{di^2} \leq 0 \Leftrightarrow t \frac{d^2 \eta}{di^2} \Big|_{s=1} + 2 \frac{d\eta}{di} \Big|_{s=1} \leq 0.$$

Here, such conditions placed on the force of infection amount to bounds on how quickly η changes relative to the infected proportion.

3. Novel recovery rates

To demonstrate model predictions with time-varying and state-dependent recovery rates, we model measles outbreaks in Reykjavik Iceland. To do this, we use a classical data set on the infectious period of measles (Simpson, 1952), time series data of measles incidence in Reykjavik Iceland (Cliff et al., 1981) from 1924 to 1928, and parameter estimates from the literature (Table 1).

First, we apply our methodology to obtain time-varying recovery rates that capture the first four statistical moments (mean, standard deviation, skewness and kurtosis) of infectious period data (Schuster, 2016, pp. 83–197). To do this, we assume the infectious period distribution belongs to the family of Pearson distributions. The family of Pearson distributions consists of types I, III, IV, V, and VI, which encompass nearly all classical continuous probability distributions, including normal, Student's t , exponential, gamma, beta, inverse gamma, beta prime, and F distributions. The family of Pearson distributions also has the distinction of being the first to fit the higher statistical moments of skewness and kurtosis arbitrarily well. Importantly, this ability implies that one can specify any mean, standard deviation, skewness and kurtosis and find a Pearson distribution that describes that probability density (Pearson, 1893, 1895; Rhind, 1909) (Fig. 1).

Using the family of Pearson distributions, we develop recovery rates that account for any mean, standard deviation, skewness and kurtosis of an infectious period distribution. We evaluate how uncertainty in these moments impacts predictions on the timing and magnitude of the epidemic peak through a variance-based sensitivity analysis (Sobol, 2001). To do this, we make use of a data set on the infectious period of measles (Simpson, 1952), and determine a sampling region for skewness and kurtosis based on the best fits of each member of the family of Pearson distributions (Fig. 2) to the observed values of skewness and kurtosis from the data.

Next, we demonstrate how to obtain state-dependent recovery rates based on a power-law distribution, outlining how such a recovery rate leads to a finite contribution to the spread of disease from infected individuals. We then demonstrate how different power-law exponents affect the trajectory of disease, through a comparison to the traditional model with a constant recovery rate.

3.1. Time-varying recovery rates associated to the Pearson distribution

To obtain time-varying recovery rates for ODEs, we take $\tilde{\lambda} = \beta i$ and impose that the infectious period distribution $Q(t)$ is distributed according to the family of Pearson distributions, which are given by the relation:

$$\left(\frac{dQ}{dt}\right)^{-1} \frac{d^2 Q}{dt^2} = -\frac{t - \lambda + a}{b_2(t - \lambda)^2 + b_1(t - \lambda) + b_0}, \tag{13}$$

where




Fig. 1. Square of skewness and kurtosis plane. The region for each member from the family of Pearson distributions is labelled, in addition to the impossible zone ($\mu_4 \geq \mu_3^2 + 1$) and sampling region for the variance-based sensitivity analysis.




Fig. 2. Best fit to Pearson distribution. Least-squares fit of the family of Pearson distributions to infectious period data of measles.

$$b_0 = \sigma^2 \frac{4\mu_4 - 3\mu_3}{10\mu_4 - 18 - 12\mu_3}, b_1 = a = \sigma \frac{\sqrt{\beta_1}(\mu_4 + 3)}{10\mu_4 - 18 - 12\mu_3}, b_2 = \frac{2\mu_4 - 3\mu_3 - 6}{10\mu_4 - 18 - 12\mu_3},$$

and λ is a scale parameter. Note, the parameters σ , μ_3 and μ_4 correspond to the standard deviation, square of skewness and kurtosis of the distribution respectively.

Each member of the family of Pearson distributions corresponds to particular assumptions placed on parameters a , b_0 , b_1 and b_2 . For each case of these assumptions we fit the resulting cumulative distribution function $(1 - Q(t))$, through a least-squares procedure, to infectious period data of measles (Fig. 2). After obtaining the best fit $Q(t)$ for each member of the family of Pearson distributions, we estimate the conditional waiting-time distribution given time t_N by

$$\Pr\{T > \tau | t = t_N\} = \frac{\sum_{j=0}^N I(t_j) Q(\tau + t_N - t_j)}{\sum_{j=0}^N I(t_j) Q(t_N - t_j)}, \tag{14}$$

where T is a random variable for the time spent in the infectious class, $\tau \geq 0$ is the time spent in the infectious class and $I(t_j)$ corresponds to the bi-weekly measles incidence data from Reykjavik Iceland.

As equation (14) is a survival function, integrating τ over all durations of infectiousness yields a series of constant residual waiting-times (Finkelstein, 2002):

$$m(t_N) = \int_0^\infty \Pr\{T > \tau | t = t_N\} d\tau.$$

Approximating $m(t)$ for all t through linear interpolation between each (t_N, t_{N+1}) , we finally determine $\eta(t)$ from (Finkelstein, 2002)

$$\eta(t) = \frac{m'(t) + 1}{m(t)}. \tag{15}$$

3.1.1. The sensitivity of model predictions to an infectious period distribution

To evaluate the sensitivity of model predictions to the uncertainty in the mean, standard deviation, skewness and kurtosis from an infectious period distribution, we calculate the variance-based sensitivity indices for predicting the time to, and magnitude of the measles epidemic peak.

We now outline the procedure to conduct the analysis. First, we estimate a skewness of 0.39 and a kurtosis of 2.92 directly from the infectious period data. We sample the ball of radius $r \approx 9.49$ centered at $(0.39^2, 2.92)$ in the $\mu_3 - \mu_4$ plane (Table 1). The value of r is the average distance from $(0.39^2, 2.92)$ to the point corresponding to the best fit parameters of each Pearson distribution type (Table 1, Fig. 1). Additional details of the sample distributions and parameters are available in Table 1. Next, for evaluating the sensitivity of model predictions relative to the uncertainty in model inputs, we measure 1) the predicted time to the measles epidemic peak, 2) the predicted magnitude of the measles epidemic peak, and 3) the combined time to, and magnitude of, the measles epidemic peak, as given by:

$$D = \sqrt{\left(I_{peak} - M \cdot \max_{t \in [0,365]} i(t)\right)^2 + \left(t_{peak} - \operatorname{argmax}_{t \in [0,365]} i(t)\right)^2},$$

where $I_{peak} = 831$, $t_{peak} = 302$ days and $M = 24134$ is the total population size.

The calculation of the variance-based sensitivity indices yields that uncertainty in the skewness of the infectious period distribution is the largest contributor to sensitivity in both predictions of the magnitude of the measles epidemic and the combined scenario (Table 2).

3.2. State-dependent recovery rates associated to the power-law distribution

To obtain state-dependent recovery rates we follow a similar approach to one that connects constant rates to exponential distributions. In particular, one assumes that no inflow into the infectious class occurs given a certain infectious proportion i_0 at some initial time t_0 (Martcheva, 2015, pp. 9–31). Under such conditions with an exponentially distributed infectious period, the infectious proportion from system (11) behaves according to a linear ODE with coefficient $-\gamma$. The solution to this ODE corresponds to the exponential decay, and $i = i_0 \exp(-\gamma(t - t_0)) \rightarrow 0$ as $t \rightarrow \infty$, with $i > 0$ for any $t < \infty$. The waiting-time distribution stems from normalizing i by i_0 , which leads to $P(t, t_0) = \exp(-\gamma(t - t_0))$. Note, due to the exponential decay $P(t, t_0) > 0$ for any $t \in [t_0, \infty)$. Physically speaking, having $P(t, t_0) > 0$ for all finite infectious periods is unrealistic, even in the most extreme of conditions.

Under the same scenario of no inflow into the infectious class, an alternative to the exponential distribution that avoids the pitfall of $P(t, t_0) > 0$ for all finite infectious periods is taking $P(t, t_0)$ to correspond to the power-law distribution,

$$P(t, t_0) = \begin{cases} (1 - c(t - t_0))^{\frac{1}{\xi}} & t < t_0 + \frac{1}{c}, \\ 0 & t \geq t_0 + \frac{1}{c}. \end{cases} \tag{16}$$

Here $c > 0$ and $\xi > 0$. Once again, making use of the assumption $i = i_0 P(t, t_0)$ we obtain

$$i = i_0 (1 - c(t - t_0))^{\frac{1}{\xi}}.$$

In addition, the mean and standard deviation for such a $P(t, t_0)$, are:

$$\mu = \frac{\xi}{c(1 + \xi)}, \quad \sigma^2 = \frac{\xi^2}{c^2(1 + \xi)^2(2\xi + 1)}. \tag{17}$$

Similarly to the exponential distribution case we have $P(t, t_0) \rightarrow 0$ as $t \rightarrow \infty$. However, now $P(t, t_0) = 0$ for $t \geq t_0 + \frac{1}{c}$.

To determine the recovery rate $\eta(i)$ that corresponds to this choice of $P(t, t_0)$, we isolate $P(t, t_0) = (1 - c(t - t_0))^{\frac{1}{\xi}}$ for $t - t_0$ yielding,

$$P(t, t_0)^{\xi} - 1 = -c(t - t_0).$$

Making the substitution $P(t, t_0) = \frac{i}{i_0}$ and rearranging the factor i_0^{ξ} yields

$$i^{\xi} - i_0^{\xi} = -c i_0^{\xi} (t - t_0). \tag{18}$$

Through differentiation and simplification, it follows that

Table 2
First-order sensitivity indices.

Scenario	Mean	Standard deviation	Skewness	Kurtosis
Magnitude of epidemic	0.013	0.013	0.244	0.071
Timing of epidemic	0.008	0.006	0.287	0.091
Timing and magnitude of epidemic	0.066	0.019	0.241	0.138

$$\frac{di}{dt} = -\frac{c}{\xi} \left(\frac{i_0}{i}\right)^\xi i = -\eta(i)i,$$

where the state-dependent recovery rate is

$$\eta(i) = \frac{c}{\xi} \left(\frac{i_0}{i}\right)^\xi. \tag{19}$$

3.2.1. The epidemic wave from an infectious period that is power-law distributed

We now demonstrate how a power-law distributed infectious period affects the shape of the predicted epidemic wave. As we are only concerned with effects of a single epidemic wave, we take $\tilde{b} = 0$ with an average infectious period of $\mu = 8.00$ days (Table 1). It follows from the formulas for the mean and standard deviation (17) that

$$c = \frac{\mu^2 - \sigma^2}{\mu(\mu^2 + \sigma^2)}, \quad \xi = \frac{1}{2} \left(\frac{\mu^2}{\sigma^2} - 1\right),$$

and

$$\eta(i) = \frac{2}{\mu} \frac{\sigma^2}{\mu^2 + \sigma^2} i^{\frac{1}{2} \frac{\mu^2 - \sigma^2}{\sigma^2} - \frac{1}{2}} \left(\frac{\mu^2}{\sigma^2} - 1\right).$$

For $\eta(i)$ to be biologically valid requires $\sigma \in \left[\frac{1}{\sqrt{3}}\mu, \mu\right]$. Violation of this interval results in 1) $\frac{di}{dt}$ being undefined at $i = 0$ when $\sigma < \frac{1}{\sqrt{3}}\mu$, or 2) the associated survival function (16) taking positive values before the initial moment of infection t_0 , as $\sigma > \mu$ forces $c < 0$. In addition, for a state-dependent recovery rate to be well-posed over the course of an entire epidemic the parameters of the power-law distribution need to be selected to account for the largest occurring infected proportion i^* , instead of just any infectious proportion i_0 at some time t_0 .

This proportion is precisely the endemic equilibrium value of i , or when no such equilibrium exists, the value of i along a trajectory when $s = \frac{1}{R_0}$ and thus $\frac{di}{dt} = 0$.

Finally, taking $\tilde{\lambda}$ as in (12), yields

$$\tilde{\lambda} = R_0 \eta(i)i = R_0 \frac{2}{\mu} \frac{\sigma^2}{\mu^2 + \sigma^2} i^{\frac{1}{2} \frac{\mu^2 - \sigma^2}{\sigma^2} - \frac{1}{2}} \left(\frac{\mu^2}{\sigma^2} - 1\right) i.$$

Because demographic factors are not included it follows that

$$\frac{di}{ds} = -1 + \frac{1}{R_0 s} \Leftrightarrow i(s) = i_0 + s_0 - \frac{1}{R_0} \left(R_0 s + \ln\left(\frac{s_0}{s}\right)\right).$$

As the peak of infection occurs for $s = \frac{1}{R_0}$, as this enforces $\frac{di}{dt} = \frac{di}{ds} = 0$, the magnitude of the peak of infection is

$$i^* = i_0 + s_0 - \frac{1}{R_0} (1 + \ln(s_0 R_0)). \tag{20}$$

Further imposing $\sigma = \frac{1}{\sqrt{3}}\mu$ reduces system (11) to a linear ODE system. From the solution of the linear ODE system we determine the time of the peak of infection,

$$t_{peak} - t_0 = 2\mu \frac{\ln(s_0 R_0)}{i^* R_0}. \tag{21}$$

With (20)–(21) and the measles incidence in Reykjavik Iceland (Cliff et al., 1981) from 1924 to 1928, we determine i^* so that the predicted epidemic peak corresponds to the peak of infection for $R_0 \in (1, 10]$ and standard deviations of an infectious period that is power-law distributed (Fig. 3).

4. Discussion

We derive infectious disease models with both time-varying and state-dependent recovery rates. To do this, we illustrate how integral equations reduce to ODE models through their connection with survival functions. We first demonstrated the novelty of time-varying recovery rates by quantifying the contribution of uncertainty in an infectious period distribution's moments to the sensitivity in predicting the epidemic peak of Iceland's 1924 measles outbreak. Our findings show that skewness had a sensitivity index at least double the other moments (Table 2). This suggests that parameter estimation techniques that rely on fitting infectious period distributions should pay close attention to the resulting value of skewness if the prediction of epidemic peaks is the ultimate goal. Secondly, we demonstrated state-dependent recovery rates derived from a power-law distribution. For such rates, we illustrated how the parameters associated with the infectious-period distribution influence the shape of an epidemic wave.

To explore the impact of uncertainty on model predictions, we used the family of Pearson distributions because it spans all possible skewness and kurtosis values (Pearson, 1893, 1895). However, alternative distribution families, such as the exponentially generalized Beta distribution, which is often used in estimation of hazard functions, or distributions associated with nonstandard parametric survival functions (Lawless, 2002) also provide further avenues for exploring the overall effects of uncertainty on model predictions.

An important extension of our approach in developing time-varying and state-dependent rates is in the evaluation of deterministic models that compare treatment outcomes, such as drug treatments for malaria (Greenhalgh, Ndeffo, Galvani, & Parikh, 2015; Tarning et al., 2012) or vaccine interventions. These models often reduce collected treatment and vaccination data to mean values in order to determine rate parameters. While models that compare treatment outcomes in such a fashion are useful, incorporating higher moments, like skewness and kurtosis, yields models that use more information and thus stand to provide more powerful predictions.

Many physically motivated functions merit use as a state-dependent recovery rate. We demonstrate how to obtain a state-dependent recovery rate from a power-law distribution, however other distributions with inverse cumulative density functions, such as the Weibull and Kumaraswamy distributions (Jones, 2009), would naturally lead to novel state-dependent recovery rates. Specifically, state-dependent recovery rates based on distributions that capture the recovery process, such as those with a finite infectious period or those that more accurately capture the concentration around the mean infectious period, provide a more powerful description of how infected individuals contribute to the spread of disease. Such state-dependent recovery rates would improve the modeling of quarantine, treatment or other policies that curtail the ability of infected individuals to spread disease.

Another benefit of state-dependent and time-varying recovery rates is the potential for new exactly solvable systems. Exactly solvable systems, such as the typical Susceptible-Infected models, are exceptionally rare and exceptionally useful in disease modeling. Thus, any new such system achieved from state-dependent and time-varying recovery rates would open the door to a plethora of new analytical insights.

The development of time-varying and state-dependent recovery rates stands to benefit the study of disease elimination and eradication. Disease elimination and eradication are currently important issues to world health authorities (Hopkins,




Fig. 3. The shape of the epidemic wave. Measles time series (black dashed line) and measles incidence for $i^* = 831/24134$, $R_0 \in (1, 10]$ with a) $\sigma = \mu$, b) $\sigma = \frac{1}{2}(1 + \frac{1}{\sqrt{3}})\mu$ and c) $\sigma = \frac{1}{\sqrt{3}}\mu$.

2013), particularly because of the partial success of the Global Malaria Eradication program (Nájera, González-Silva, & Alonso, 2011) and the successful eradication of both smallpox (Fenner, Henderson, Arita, Jezek, & Ladnyi, 1988) and rinderpest (Normile, 2008). For diseases slated for eradication, such as measles, yaws and polio, models that accurately capture the physical process of disease spread stand to provide quantitative information on policies and interventions designed to expedite the elimination and eradication process. As time-varying and state-dependent recovery rates better account for the contribution of infected individuals to disease spread and also include the potential for disease burnout (Greenhalgh, Galvani, et al., 2015), they thereby provide invaluable information absent from more traditional modeling formulations.

5. Conclusions

Approaches to accurately account for non-constant rates in deterministic epidemic models is recognized as an open challenge in infectious disease modeling (Roberts, Andreasen, Lloyd, & Pellis, 2015). Here, we provided an approach to account for time-varying and state-dependent rates developed from infectious period data. By basing time-varying and state-dependent recovery rates on infectious period distributions derived from such data, ODE models of infectious diseases are able to more accurately account for the dynamics of transmission. While we developed ODE models of infectious diseases with non-constant rates based on such infectious period data, our approach is easily generalizable to novel rates from alternate sources, including viral load, latent period and pharmacokinetic data. In summary, our work enables disease and compartmental modellers to shed the shackles imposed by exponentially distributed waiting times and constant rates by enabling the discovery and use of new non-constant rates that more accurately capture the dynamics of disease.

Acknowledgements

The authors would like to express thanks to the anonymous reviewers whose quality critiques helped to improve and clarify this manuscript.

References

- Alexander, M. E., & Moghadas, S. M. (2005). Bifurcation analysis of an SIRS epidemic model with generalized incidence. *SIAM Journal on Applied Mathematics*, 65(5), 1794–1816. <https://doi.org/10.1137/040604947>.
- Arino, J., & van den Driessche, P. (2006). Time delays in epidemic models. In *Delay differential equations and applications* (pp. 539–578). Dordrecht: Springer Netherlands. https://doi.org/10.1007/1-4020-3647-7_13.
- Bailey, N. T. J., & others. (1975). *The mathematical theory of infectious diseases and its applications*. 5a Crendon Street, High Wycombe, Bucks HP13 6LE: Charles Griffin & Company Ltd.
- Ball, F. (1983). The threshold behaviour of epidemic models. *Journal of Applied Probability*, 20(2), 227. <https://doi.org/10.2307/3213797>.
- Ball, F. (1986). A unified approach to the distribution of total size and total area under the trajectory of infectives in epidemic models. *Advances in Applied Probability*, 18(2), 289. <https://doi.org/10.2307/1427301>.
- Ball, F. G. (1991). Dynamic population epidemic models. *Mathematical Biosciences*, 107(2), 299–324. [https://doi.org/10.1016/0025-5564\(91\)90011-7](https://doi.org/10.1016/0025-5564(91)90011-7).
- Britton, T. (2010). Stochastic epidemic models: A survey. *Mathematical Biosciences*, 225(1), 24–35. <https://doi.org/10.1016/j.mbs.2010.01.006>.
- Clancy, D. (1999). Outcomes of epidemic models with general infection and removal rate functions at certain stopping times. *Journal of Applied Probability*, 36(3), 799–813. Retrieved from <http://www.jstor.org/stable/3215442>.
- Clancy, D. (2014). SIR epidemic models with general infectious period distribution. *Statistics & Probability Letters*, 85, 1–5. <https://doi.org/10.1016/j.spl.2013.10.017>.
- Cliff, A. D., Haggett, P., Ord, J. K., & Versey, G. R. (1981). *Spatial diffusion: An historical geography of epidemics in an island community*. Cambridge University Press.
- Diekmann, O., Heesterbeek, J. A. P., & Metz, J. A. J. (1990). On the definition and the computation of the basic reproduction ratio R_0 in models for infectious diseases in heterogeneous populations. *Journal of Mathematical Biology*, 28(4). <https://doi.org/10.1007/BF00178324>.
- van den Driessche, P., & Watmough, J. (2000). A simple SIS epidemic model with a backward bifurcation. *Journal of Mathematical Biology*, 40(6), 525–540. <https://doi.org/10.1007/s002850000032>.
- van den Driessche, P., & Watmough, J. (2002). Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical Biosciences*. [https://doi.org/10.1016/S0025-5564\(02\)00108-6](https://doi.org/10.1016/S0025-5564(02)00108-6).
- Feng, Z., Xu, D., & Zhao, H. (2007). Epidemiological models with non-exponentially distributed disease stages and applications to disease control. *Bulletin of Mathematical Biology*, 69(5), 1511–1536. <https://doi.org/10.1007/s11538-006-9174-9>.
- Fenner, F., Henderson, D. A., Arita, I., Jezek, Z., & Ladnyi, I. D. (1988). *Smallpox and its eradication*. Geneva.
- Finkelstein, M. S. (2002). On the reversed hazard rate. *Reliability Engineering & System Safety*, 78(1), 71–75. [https://doi.org/10.1016/S0951-8320\(02\)00113-8](https://doi.org/10.1016/S0951-8320(02)00113-8).
- Fowler, A. C., & Hollingsworth, T. D. (2015). Simple approximations for epidemics with exponential and fixed infectious periods. *Bulletin of Mathematical Biology*, 77(8), 1539–1555. <https://doi.org/10.1007/s11538-015-0095-3>.
- Gough, K. J. (1977). The estimation of latent and infectious periods. *Biometrika*, 64(3), 559–565.
- Greenhalgh, S., Galvani, A. P., & Medlock, J. (2015). Disease elimination and re-emergence in differential-equation models. *Journal of Theoretical Biology*, 387, 174–180. <https://doi.org/10.1016/j.jtbi.2015.09.037>.
- Greenhalgh, S., Ndeffo, M., Galvani, A. P., & Parikh, S. (2015). The epidemiological impact of HIV antiretroviral therapy on malaria in children. *AIDS*, 29(4), 473–482. <https://doi.org/10.1097/qad.0000000000000550>.
- Heffernan, J. M., Smith, R. J., & Wahl, L. M. (2005). Perspectives on the basic reproductive ratio. *Journal of the Royal Society, Interface/the Royal Society*, 2, 281–293. <https://doi.org/10.1098/rsif.2005.0042>.
- Hethcote, H. W., & Tudor, D. W. (1980). Integral equation models for endemic infectious diseases. *Journal of Mathematical Biology*, 9(1), 37–47. <https://doi.org/10.1007/BF00276034>.
- Hopkins, D. R. (2013). Disease eradication. *New England Journal of Medicine*, 368(1), 54–63. <https://doi.org/10.1056/NEJMr1200391>.
- Jones, M. C. (2009). Kumaraswamy's distribution: A beta-type distribution with some tractability advantages. *Statistical Methodology*, 6(1), 70–81. <https://doi.org/10.1016/j.stamet.2008.04.001>.
- Keeling, M., & Grenfell, B. (1999). Stochastic dynamics and a power law for measles variability. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 354(1384), 769–776. <https://doi.org/10.1098/rstb.1999.0429>.
- Keeling, M. J., & Rohani, P. (2007). *Modeling infectious diseases in humans and animals*. Princeton University Press.

- Korobeinikov, A., & Maini, P. K. (2005). Non-linear incidence and stability of infectious disease models. *Mathematical Medicine and Biology: A Journal of the IMA*, 22(2), 113–128. <https://doi.org/10.1093/imamm/dqi001>.
- Krylova, O., & Earn, D. J. D. (2013). Effects of the infectious period distribution on predicted transitions in childhood disease dynamics. *Journal of The Royal Society Interface*, 10(84). <https://doi.org/10.1098/rsif.2013.0098>, 20130098–20130098.
- Lawless, J. F. (2002). *Statistical models and methods for lifetime data*. Hoboken, NJ, USA, NJ, USA: John Wiley & Sons, Inc. <https://doi.org/10.1002/9781118033005>.
- Liu, W. M., Hethcote, H. W., & Levin, S. A. (1987). Dynamical behavior of epidemiological models with nonlinear incidence rates. *Journal of Mathematical Biology*, 25(4), 359–380. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3668394>.
- Liu, W. M., Levin, S. A., & Iwasa, Y. (1986). Influence of nonlinear incidence rates upon the behavior of SIRS epidemiological models. *Journal of Mathematical Biology*, 23(2), 187–204. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/3958634>.
- Lloyd, A. L. (2001a). Destabilization of epidemic models with the inclusion of realistic distributions of infectious periods. *Proceedings of the Royal Society B: Biological Sciences*, 268(1470), 985–993. <https://doi.org/10.1098/rspb.2001.1599>.
- Lloyd, A. L. (2001b). Realistic distributions of infectious periods in epidemic models: Changing patterns of persistence and dynamics. *Theoretical Population Biology*, 60(1), 59–71. <https://doi.org/10.1006/tpbi.2001.1525>.
- Martcheva, M. (2015). *Introduction to epidemic modeling*. https://doi.org/10.1007/978-1-4899-7612-3_2.
- Nájera, J. A., González-Silva, M., & Alonso, P. L. (2011). Some lessons for the future from the global malaria eradication programme (1955–1969). *PLoS Medicine*, 8(1), e1000412. <https://doi.org/10.1371/journal.pmed.1000412>.
- Normile, D. (2008). Rinderpest. Driven to extinction. *Science (New York, N.Y.)*, 319(5870), 1606–1609. <https://doi.org/10.1126/science.319.5870.1606>.
- Pearson, K. (1893). Contributions to the mathematical theory of evolution. *Proceedings of the Royal Society of London*, 54(326–330), 329–333. <https://doi.org/10.1098/rspl.1893.0079>.
- Pearson, K. (1895). Contributions to the mathematical theory of evolution. II. Skew variation in homogeneous material. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 186, 343–414. <https://doi.org/10.1098/rsta.1895.0010>.
- Ponciano, J. M., & Capistrán, M. A. (2011). First principles modeling of nonlinear incidence rates in seasonal epidemics. *PLoS Computational Biology*, 7(2), e1001079. <https://doi.org/10.1371/journal.pcbi.1001079>.
- Rhind, A. (1909). Tables to facilitate the computation of the probable errors of the chief constants of skew frequency distributions. *Biometrika*, 7(1–2), 127–147. <https://doi.org/10.1093/biomet/7.1-2.127>.
- Roberts, M., Andreasen, V., Lloyd, A., & Pellis, L. (2015). Nine challenges for deterministic epidemic models. *Epidemics*, 10, 49–53. <https://doi.org/10.1016/j.epidem.2014.09.006>.
- Rodriguez, G. (2007). *Lecture notes on generalized linear models*. Princeton University. Retrieved from <http://data.princeton.edu/wws509/notes/>.
- Ruan, S., & Wang, W. (2003). Dynamical behavior of an epidemic model with a nonlinear incidence rate. *Journal of Differential Equations*, 188(1), 135–163. [https://doi.org/10.1016/S0022-0396\(02\)00089-X](https://doi.org/10.1016/S0022-0396(02)00089-X).
- Schuster, P. (2016). *Distributions, moments, and statistics*. https://doi.org/10.1007/978-3-319-39502-9_2.
- Simpson, R. E. H. (1952). Infectiousness of communicable diseases in the household (measles, chickenpox, and mumps). *Lancet (London, England)*, 2(6734), 549–554. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12981903>.
- Sobol, I. (2001). Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates. *Mathematics and Computers in Simulation*, 55(1–3), 271–280. [https://doi.org/10.1016/S0378-4754\(00\)00270-6](https://doi.org/10.1016/S0378-4754(00)00270-6).
- Tarning, J., Zongo, I., Somé, F. A., Rouamba, N., Parikh, S., Rosenthal, P. J., ... Lindegardh, N. (2012). Population pharmacokinetics and pharmacodynamics of piperazine in children with uncomplicated falciparum malaria. *Clinical Pharmacology and Therapeutics*, 91(3), 497–505. <https://doi.org/10.1038/clpt.2011.254>.